UNDER THE FEE OF INSEUTICIDE PUNGICIDE AND PUTCHTURE ACT FOR CONOMIC . U.UN REGISTER. 10/103 ED LUNDER NO.

# **PROKIL** Perthane Malathion 10-4 Dust

### SUGGESTED USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSE-QUENCE OF LOCAL WEATHER OR CONDITIONS; WE **RECOMMEND REFERENCE TO AGRICULTURAL EXPERI-**MENT STATIONS ON SPECIFIC USAGE

#### BURN BAGERMANN LATELY WHEN PLAFTY CTAY OUT F CLOKE.

**PROKIL** Perthane-Malathion 10-4 Dust is suitable for application through a ground duster or by airplane.

#### DIRECTIONS

Use 25 pounds per acre unless otherwise specified Thorough coverage is essential. Repeat application as necessary.

CABBAGE, BRUSSELS SPROUTS, CAULIFLOWER: Cabbage Worm. Cabbage Looper, Aphids - Do not apply within 7 days of harvest

8 71 114- PROOF

CALIFORNIA (Plant) P. O. Box 406 340 East Main Street Calipatria, Calif. 92233 Telephone: 348-5150

Active Ingredients

Diethyi Dipheriyl Dichloroethune

Related Compounds

Mclathion\*\*

Inerr Ingredients

\*\*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

## READ THE LABEL CAUTION **KEEP OUT OF REACH OF CHILDREN**

Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs Do not apply under conditions involving possible drift to food, forage or other plantings that night be damaged or the crop thereof rendered unfit for sale, use or consumption.

SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS!

## NET WEIGHT 50 LBS.



<b>t</b> .	Ŵ	by	%
8	8.		
2	1.		
0	4.		

86.0

100.0

LETTUCE: Cabbage Looper, Leafhoppers, Importea Cabbage Worm, Aphids — Apply when insects appear. Repeat at weekly or 10-day intervals as necessary. Do not apply within 7 days of harvest on head lettuce or 14 days on leaf lettuce.

SPINACH: Leafhoppers, Cabbage Looper, Aphids, --- Apply at weekly to 10-day intervals. Do not apply within 7 days of harvest.

This product is toxic to fish. Keep out of lakes streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not cipply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal or wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to treat ment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

#### NOTICE ON CONDITIONS OF SALE

والمرجع والمرجع العمروق متعاور والمعاقب المراجع المراجع الأرجان and the residue. The appendix product to the stress of the second recommendation, · · · - the stand with a - Press of the street on story in the work area The Elevency t ••• and the second n n rai stan vin

USDA Reg. No. 10163-37 Calif. Reg. No. 10163-X

> ARIZONA Route 1, Box 33

Somerton, Arizona 85350

Telephone: 627-2479