

8-30-73  
 UNDER THE FEDERAL INSECTICIDE ACT  
 FOR ECONOMIC PEST CONTROL REGISTERED  
 UNDER NO. 10163-40

# PROKIL Cryolite 75 Dust

ACTIVE INGREDIENTS	% by Wt.
SODIUM FLUOALUMINATE*	72.0
INERT INGREDIENTS	28.0
	100.0
*Flourine expressed as elemental	40.6

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
 Avoid Inhalation of Dust

THIS PRODUCT HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**NET WEIGHT 50 LBS.**

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USES.

KEEP PESTICIDE IN ORIGINAL CONTAINER!

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

Use at rate of 17 to 40 pounds per acre depending on plant size and insect population. Repeat application as necessary. Remove residues remaining at harvest by washing, brushing or other effective means. Do not apply under conditions involving possible drift to food, forage or other planting that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

6-72-113A-1M

EPA Reg. No. 10163-40 AA

BEANS: Mexican Bean Beetles, Bean Leafroller—Repeat at two week intervals.

CABBAGE, CAULIFLOWER, BROCCOLI: Flea Beetles, Cabbage Caterpillars — DO NOT APPLY to Broccoli after the edible parts start to form. DO NOT APPLY to Cabbage after the heads start to form. DO NOT APPLY to Cauliflower after the seedling stage.

CARROTS: Vegetable Weevil.

CUCUMBERS: Striped Cucumber Beetle, Twelve-Spotted Cucumber Beetle, Flea Beetle — Dust both sides of foliage at first signs of Beetles.

EGGPLANT: Flea Beetle — Dust thoroughly.

LETTUCE: Climbing Cutworms, Alfalfa Looper, Corn Earworm — Apply when worms first appear. DO NOT APPLY after the seedling stage.

MELONS: Cucumber Beetle — Dust both sides of foliage. Repeat if necessary.

PEPPERS: Pepper Weevil — Dust thoroughly at first set of fruit. Repeat every 5 days until about 6 applications have been made.

STRAWBERRIES: Leaf Beetle — Dust both sides of leaves.

TOMATOES: Flea Beetle — Apply when plants are first set out and repeat as necessary. If applied after fruit starts to set, remove residue at harvest by washing, brushing or other effective means.

### NOTICE ON CONDITIONS OF SALE


Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse or such as when combined with other materials.

CALIFORNIA (Plant)  
 P. O. Box 406  
 340 East Main Street  
 Calipatria, Calif. 92233  
 Telephone: 348-5150

# The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA  
 Route 1, Box 33  
 Somerton, Arizona 85350  
 Telephone: 627-2479



# OKIL Cryolite 75 Dust

PERCENTS	By Wt.
DIPOCALUMINATE*	72.0
PERCENTS	28.0
purine expressed as elemental	100.0
	40.6

## CAUTION

OUT OF REACH OF CHILDREN  
Avoid Inhalation of Dust

VERY TOXIC TO BEES exposed to treatment or residues on  
attention may be obtained from your Cooperative Agricultural

NET WEIGHT 50 LBS.

USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARN-  
INGS, AND WITH APPLICABLE STATE AND FEDERAL REGULA-  
TIONS. DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL  
CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICUL-  
TURAL AGENCIES CONCERNING SPECIFIC USES.

ORIGINAL CONTAINER!

DESTROY WHEN EMPTY.

40 pounds per acre depending on plant size and insect  
population as necessary. Remove residues remaining at  
harvest by washing or other effective means. Do not apply under  
conditions of possible drift to food, forage or other planting that might  
be rendered unfit for sale, use or consumption.

EPA REG. NO. 10103-46

BEANS: Mexican Bean Beetle, Bean Leaf-roller—Repeat at two week intervals.

CABBAGE, CAULIFLOWER, BROCCOLI: Flea Beetles, Cabbage Caterpillars — DO  
NOT APPLY to Broccoli after the edible parts start to form. DO NOT APPLY  
to Cabbage after the heads start to form. DO NOT APPLY to Cauliflower after  
the seedling stage.

CARROTS: Vegetable Weevil.

CUCUMBERS: Striped Cucumber Beetle, Twelve-Spotted Cucumber  
Beetle, Flea Beetle — Dust both sides of foliage at first signs of  
Beetles.

EGGPLANT: Flea Beetle — Dust thoroughly.

LETTUCE: Climbing Cutworms, Alfalfa Looper, Corn Earworm — Apply when  
worms first appear. DO NOT APPLY after the seedling stage.

MELONS: Cucumber Beetle — Dust both sides of foliage. Repeat if necessary.

PEPPERS: Pepper Weevil — Dust thoroughly at first set of fruit. Repeat every 5  
days until about 6 applications have been made.

STRAWBERRIES: Leaf Beetle — Dust both sides of leaves.

TOMATOES: Flea Beetle — Apply when plants are first set out and repeat as  
necessary. If applied after fruit starts to set, remove residue at harvest by  
washing, bushing or other effective means.

### NOTICE ON CONDITIONS OF SALE

Conditions of sale for this product are based upon tests believed to be  
representative. The user assumes full responsibility for the control of the manufacturer, no  
warranty is made for the results of such use. The results to be  
obtained are dependent upon the user's application of this product in accordance with the  
label directions and good agricultural practices. The user assumes full responsibility for any  
damage, resulting from its use.

# The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA

Route 1, Box 33

Somerton, Arizona 85350

Telephone: 627-2479