

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTRATION
ED UNDER NO. 10163-41

PROKIL Cryolite 96

ACTIVE INGREDIENTS	% by Wt.
Sodium Fluoaluminate*	96
INERT INGREDIENTS	4
	<hr/> 100

*Fluorine expressed as elemental 50%

CAUTION

KEEP OUT OF REACH OF CHILDREN
AVOID INHALATION OF DUST

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NET WEIGHT 50 LBS.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USES.

KEEP PESTICIDE IN ORIGINAL CONTAINER!

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE or BURNING. STAY AWAY FROM SMOKE or FUMES.

This product is formulated for application by ground or air equipment.

Use at rate of 25 to 50 pounds per acre depending on plant size and insect population. Repeat application as necessary. Remove residues remaining at harvest by washing, brushing or other effective means. Do not apply under conditions involving possible drift to food, forage or other planting that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

7-72—164 PROOF

EPA Reg. No. 10163-41 AA

BEANS—Mexican bean beetle, Bean leaf beetle, bean leaf roller, corn earworm, cucumber beetles.

CABBAGE, COLLARDS, MUSTARD, TURNIPS, BROCCOLI, RADISHES, and CAULIFLOWER—Imported cabbage worm, cabbage looper, diamond-back moth caterpillar, yellow striped armyworm, climbing cut worms, corn earworm, flea beetles.

CANTALOUPE, WATERMELON, CUCUMBER and SQUASH—Melon worms, pickle-worm, cucumber beetles.

TOMATO—Hornworm, fruitworm, pinworm, blister beetle.

APPLES, PEACHES, PEARS, SHADE TREES, and ORNAMENTALS—Flea weevil, codling moth, gypsy moth caterpillar, Fuller's rose beetle, plum curculio.

GRAPEFRUIT, LEMONS, ORANGES—Orange Tortrix.

CRANBERRY—Cranberry weevil, gypsy moth caterpillar, false army worm.

LETTUCE—Cabbage looper.

GRAPES—Grape leaf folder.

STRAWBERRIES—Strawberry leaf roller.

PEPPERS—Pepper weevil.

LIMITATIONS—Do not use in combination with lime or compounds containing free lime. Do not apply immediately before harvest. Do not apply to bush and vine fruits after fruit starts to form. Do not apply to cabbage, broccoli or grapes after heads or edible parts start to form. Do not apply to collards, mustard, cauliflower or lettuce after the seedling stage or to turnips after seedling stage if tops are to be used for food.

Remove excess residues on edible portions of fruit and vegetables by washing, brushing or other effective means.

NOTICE ON CONDITIONS OF SALE


Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practices. The buyer must assume all responsibility, including injury or damage, resulting from misuse as such, or in combination with other materials.

CALIFORNIA (Plant)
P. O. Box 406
340 East Main Street
Calipatria, Calif. 92233
Telephone: 348-5150

The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA
Route 1, Box 33
Somerton, Arizona 85350
Telephone: 627-2479



PROKIL Cryolite 96

ACTIVE INGREDIENTS	% by Wt.
Sodium Fluoroaluminate*	96
INERT INGREDIENTS	4
	<hr/> 100
*Fluorine expressed as elemental	50%

CAUTION

KEEP OUT OF REACH OF CHILDREN
AVOID INHALATION OF DUST

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NET WEIGHT 50 LBS.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USES.

KEEP PESTICIDE IN ORIGINAL CONTAINER!

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE or BURNING. STAY AWAY FROM SMOKE or FUMES.

This product is formulated for application by ground or air equipment.

Use at rate of 25 to 50 pounds per acre depending on plant size and insect population. Repeat application as necessary. Remove residues remaining at harvest by washing, brushing or other effective means. Do not apply under conditions involving possible damage to food, forage or other planting that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

7-72-164 PROOF

EPA Reg. No. 10163-41 AA

BEANS—Mexican bean beetle, bean leaf beetle, bean leaf weevil, cucumber beetles.

CABBAGE, COLLARDS, MUSTARD, TURNIPS, BROCCOLI, RUCIFLOWER—Imported cabbage worm, cabbage looper, diamond borer, yellow striped armyworm, climbing cut worms, corn ear worm, cucumber beetles.

TOMATO—Horn worm, fruitworm, pinworm, blister beetle. APPLES, PEACHES, PEARS, SHADE TREES, and ORNAMENTALS—Imported cabbag moth caterpillar, Fuller's rose beetle, plum borer.

GRAPEFRUIT, LEMONS, ORANGES—Orange Tortrix. CRANBERRY—Cranberry weevil, gypsy moth caterpillar, false codling moth.

LETTUCE—Cabbage looper.

GRAPES—Grape leaf folder.

STRAWBERRIES—Strawberry leaf roller.

PEPPERS—Pepper weevil.

LIMITATIONS—Do not use in combination with lime or complete fertilizer. Do not apply immediately before harvest. Do not apply to fruits after fruit starts to form. Do not apply to cabbage, cauliflower or lettuce after the seedling stage or to turnips after heads or edible parts start to form. Do not apply to cauliflower or lettuce after the seedling stage or to turnips after heads or edible parts start to form.

Remove excess residues on edible portions of fruit and vegetable by washing, brushing or other effective means.


NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon reliable information. The use of this product being beyond the control of the manufacturer, no guarantee is made as to the effects of use. Buyer must assume all responsibility, including injury or damage to crops, animals or other materials, resulting from the use of this product.

CALIFORNIA (Plant)
P. O. Box 406
340 East Main Street
Colipatria, Calif. 92233
Telephone: 348-5150

The Dune Company
AGRICULTURAL CHEMICALS

Somerton,
Telep



PROKIL Cryolite 96^T

	% by Wt.
	96
	4
	100
as elemental	50%

CAUTION
 OUT OF REACH OF CHILDREN
 AVOID INHALATION OF DUST
 TOXIC TO BEES exposed to treatment or residues on
 may be obtained from your Cooperative Agricultural

WEIGHT 50 LBS.
 STRICTLY IN ACCORDANCE WITH CAUTIONS, WARN-
 WITH APPLICABLE STATE AND FEDERAL REGULA-
 OSAGE MAY VARY AS A CONSEQUENCE OF LOCAL
 WE RECOMMEND REFERENCE TO STATE AGRICUL-
 CONCERNING SPECIFIC USES.

CONTAINER!
 LINER. DESTROY IT BY BURYING WITH WASTE or
 OR SMOKE or FUMES.

Application by ground or air equipment.
 Pounds per acre depending on plant size and insect
 as necessary. Remove residues remaining at
 or other effective means. Do not apply under
 to food, forage or other planting that might
 be rendered unfit for sale, use or consumption.

EPA Reg. No. 10163-41 AA

- BEANS— Mexican bean beetle, bean leaf beetle, bean leaf roller, corn earworm, cucumber beetles.
- CABBAGE, COLLARDS, MUSTARD, TURNIPS, BROCCOLI, RADISHES, and CAULIFLOWER— Imported cabbage worm, cabbage looper, diamond-back moth caterpillar, yellow striped armyworm, climbing cut worms, corn earworm, flea beetles.
- CANTALOUPE, WATERMELON, CUCUMBER and SQUASH— Melon worms, pickle-worm, cucumber beetles.
- TOMATO— Horn worm, fruitworm, pinworm, blister beetle.
- APPLES, PEACHES, PEARS, SHADE TREES, and ORNAMENTALS— Flea weevil, coal-ing moth, gypsy moth caterpillar, Fuller's rose beetle, plum curculio.
- GRAPEFRUIT, LEMONS, ORANGES— Orange Tortrix.
- CRANBERRY— Cranberry weevil, gypsy moth caterpillar, false army worm.
- LETTUCE— Cabbage looper.
- GRAPES— Grape leaf folder.
- STRAWBERRIES— Strawberry leaf roller.
- PEPPERS— Pepper weevil.

LIMITATIONS— Do not use in combination with lime or compounds containing free lime. Do not apply immediately before harvest. Do not apply to bush and vine fruits after fruit starts to form. Do not apply to cabbage, broccoli or grapes after heads or edible parts start to form. Do not apply to collards, mustard, cauliflower or lettuce after the seedling stage or to turnips after seedling stage if tops are to be used for food.
 Remove excess residues on edible portions of fruit and vegetables by washing, brushing or other effective means.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based on data believed to be reliable. The use of this product beyond the limits of the label is not a guarantee expressed or implied. No liability is assumed for any loss or damage obtained if not used in accordance with the label. The buyer must assume all responsibility, including injury to persons, animals or crops, or to such, or to other property with other persons.

The Dune Company
 AGRICULTURAL CHEMICALS

ARIZONA
 Route 1, Box 33
 Somerton, Arizona 85350
 Telephone: 627-2475