

1/3 10163-41 PM-16

05 NOV 1987

305  
10  
198,852  
2

Ms. Betsey M. Bowman  
Registration Specialist  
Gowan Company  
P.O. Box 5696  
Yuma, AZ 85364

Dear Ms. Bowman:

Subject: Prokil Cryolite 96  
EPA Registration No. 10163-41  
Your Amendment Dated April 13, 1987

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

Submit five (5) copies of your final printed labeling incorporating the following correction before you release the product for shipment.

The residue data referenced on your data matrix for grapes were found acceptable to support the use of the subject product on raisin, wine, and table grapes in California with application of up to 30 days before harvest. Refer to our letter of September 30, 1985 regarding our review of these data.

Accordingly, your proposed label directions for grapes (grown for raisins, wine, and/or table) must specify for use in California only.

94258-I-Nautz-N-13-KENCO-10/13/87-10/22/87-NocCoc-VO-JH-SONJA  
CONCURRENCES

SYMBOL							
SURNAME							
DATE							


2/3

-2-

If this condition is not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of this condition.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,



William H. Miller  
Product Manager (16)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767C)

Enclosure

3/5

# PROKIL Cryolite 96

ACTIVE INGREDIENT	% by Wt.
Sodium Fluoroaluminate*	96
INERT INGREDIENTS	4
Total	100
*Fluorine expressed as elemental ...	50%

## DIRECTIONS FOR USE

READ entire label. Use strictly in accordance with cautions and directions, and with applicable State and Federal regulations. Uses, timing and dosage may vary as a consequence of local weather or conditions. We recommend reference to State Agricultural Extension Service concerning specific uses.

This product is formulated for application by ground or air equipment. Unless otherwise specified, use at rate of 25 to 50 pounds per acre depending on plant size and insect population. Repeat application as necessary. Do not apply under conditions involving possible drift to food, forage or other planting that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN, wash with soap and water.  
IF IN EYES, flush eyes with plenty of clear water for at least 15 minutes.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

HARMFUL IF SWALLOWED, ABSORBED THROUGH SKIN, OR INHALED. CAUSES MODERATE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist or dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not enter treated areas without protective clothing until sprays have dried or dusts have settled. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and oral warnings must include the following information: CAUTION—Area treated with Cryolite (date of application). Do not enter without appropriate protective clothing until sprays have dried or dusts have settled. In case of accidental exposure, follow instructions under Statement of Practical Treatment section.

### STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

NOV 5 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 10163-41

# Gowan Company

AGRICULTURAL CHEMICALS

EPA Reg. No. 10163-41 AA  
EPA Est. No. 48370-DN-01

P.O. Box 5696  
Yuma, AZ 85364

NET WEIGHT 50 LBS.

BEANS—Mexican Bean Beetle, Bean Leaf Beetle, Bean Leaf Roller, Corn Earworm, Cucumber Beetles.  
CABBAGE, COLLARDS, MUSTARD, TURNIPS, BROCCOLI, RADISHES, and CAULIFLOWER—Imported Cabbage Worm, Cabbage Looper, Diamond-back Moth Caterpillar, Yellow Striped Armyworm, Climbing Cutworms, Corn Earworm, Flea Beetles.  
CANTALOUPE, WATERMELON, CUCUMBER and SQUASH—Melon Worms, Pickleworm, Cucumber Beetles.  
TOMATO—Tomworm, Fruitworm, Pinworm, Blister Beetle  
APPLES, PEACHES, PEARS, SHADE TREES, and ORNAMENTALS—Flea weevil, Codling Moth Caterpillar, Gypsy Moth Caterpillar, Fuller's Rose Beetle, Plum Curculio.  
GRAPEFRUIT, LEMONS, ORANGES—Orange Tortrix, <sup>Fuller's Rose</sup> Beetle  
CRANBERRY—Cranberry Weevil, Gypsy Moth Caterpillar, False Army Worm.  
LETTUCE—Cabbage Looper.

STRAWBERRIES—Strawberry Leaf Roller.  
PEPPERS—Pepper Weevil.

For above crops—  
LIMITATIONS—Do not use in combination with lime or compounds containing free lime. Do not apply immediately before harvest. Do not apply to bush and vine fruits after fruit starts to form. Do not apply to cabbage, broccoli after heads or edible parts start to form. Do apply to collards, mustard, cauliflower or lettuce after the seedling stage or to turnips after seedling stage if tops are to be used for food. Remove excess residues on edible portions of fruit and vegetables by washing, brushing or other effective means.

GRAPES: Grape leaf folder - Apply 25 to 50 lbs. per acre depending on plant size and insect population. Repeat application as necessary. Do not apply after fruit starts to form.

GRAPES (Grown for Raisins, Wine and/or Table (fresh market)): Grape leaf folder, Omnivorous Leafroller, and Grape Leaf Skeletonizer - Apply 0 to 8 lbs. per acre by ground equipment using an amount of water sufficient for adequate coverage. For application by air, use 8 lbs. Cryolite in 20 gallons of water. Apply as needed, but no more than two treatments per crop year. Do not apply later than 30 days prior to harvest.

Limitations and Precautions: Growers must inform the processors that Prokil Cryolite 96 has been used on the crop and that raisins must be washed.

Precaution for use on Table Grapes: Physical residues may be visible at harvest when application is made on maturing fruit, especially on the darker varieties. Local discretion should be used.

### NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practices. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

rev/app 10/3/85

Insecticide or Fungicides, Agricultural, Dry, N.O.I.B.N.