

PM 14 1063 --

# PROKIL Cryolite 96

ACTIVE INGREDIENT	% by Wt.
Sodium Fluoroaluminate	96
Aluminafluoride	4
<b>INERT INGREDIENTS</b>	<b>100</b>
Total	100
*Fluorine expressed as elemental	50%

## KEEP OUT OF REACH OF CHILDREN CAUTION

**STATEMENT OF PRACTICAL TREATMENT**  
 IF ON SKIN, wash with soap and water  
 IF IN EYES, flush eyes with plenty of clear water for at least 15 minutes.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

**HARMFUL IF SWALLOWED, ABSORBED THROUGH SKIN, OR INHALED. CAUSES MODERATE EYE IRRITATION.** Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist or dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (i.e. swamps, bays, marshes and estuaries). Do not contaminate water in an opening of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not enter treated areas without protective clothing until sprays have dried or dusts have settled. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and oral warnings must include the following information: CAUTION—Area treated with Cryolite on (date of application). Do not enter without appropriate protective clothing until sprays have dried or dusts have settled. In case of accidental exposure, follow instructions under Statement of Practical Treatment section.

Do not apply this product through any type of irrigation system.

#### STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

#### DIRECTIONS FOR USE

READ entire label. Use strictly in accordance with cautions and directions, and with applicable State and Federal regulations. Use, timing

and dosage may vary as a consequence of local weather or conditions. We recommend reference to State Agricultural Extension Service concerning specific uses.

This product is formulated for application by ground or air equipment. Unless otherwise specified, use at rate of 25 to 50 pounds per acre depending on plant size and insect population. Repeat application as necessary. Do not apply under conditions involving possible drift to food, forage or other planting that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

- BEANS**—Mexican Bean Beetle, Bean Leaf Beetle, Bean Leaf Roller, Corn Earworm, Cucumber Beetle.
- CABBAGE, COLLARDS, MUSTARD, TURNIPS, BROCCOLI, RADISHES, and CAULIFLOWER**—Imported Cabbage Worm, Cabbage Looper, Diamond-back Moth Caterpillar, Yellow Striped Armyworm, Climbing Cutworms, Corn Earworm, Flea Beetles.
- CANTALOUPE, WATERMELON, CUCUMBER and SQUASH**—Moth Worms, Pickleworm, Cucumber Beetle.
- TOMATO**—Hornworm, Fruitworm, Pinworm, Earler Beetle.
- APPLES, PEACHES, PEARs, SHADE TREES, and ORNAMENTALS**—Flea weevil, Codling Moth Caterpillar, Gypsy Moth Caterpillar, Fuller's Rose Beetle, Plum Curculio.
- GRAPEFRUIT, LEMONS, ORANGES**—Orange Tortrix, Fuller's Rose Beetle.
- CRANBERRY**—Cranberry Weevil, Gypsy Moth Caterpillar, False Army Worm.
- LETTUCE**—Cabbage Looper.
- GRAPES**—Grape Leaf Folder.
- STRAWBERRIES**—Strawberry Leaf Roller.
- PEPPERS**—Pepper Weevil.

**KIMIFRUIT**—(California only) Quinceleafroller - Apply 10 TBS. per acre with ground equipment only using 200 gallon of spray per acre. Apply up to 4 applications per season on a 15-30 day interval between applications. Do not apply until 30 days of harvest.

**LIMITATIONS FOR ABOVE CROPS**—Do not use in combination with lime or compounds containing free lime. Do not apply immediately before harvest. Do not apply to bush and vine fruits after fruit starts to form. Do not apply to cabbage or broccoli after heads or edible parts start to form. Do not apply to collards, mustard, cauliflower or lettuce after the seedling stage or to turnips after seedling stage if tops are to be used for food. Remove excess residues on edible portions of fruit and vegetables by washing, brushing or other effective means.

**GRAPES**—Grape leaf folder—Apply 25 to 50 lbs. per acre depending on plant size and insect population. Repeat application as necessary. Do not apply after fruit starts to form.

**GRAPES (California Only)**—(Grown for Raisins, Wine and/or Table (fresh market))—Grape leaf folder, Quinceleafroller, and Grape Leaf Skeletonizer—Apply 6 to 8 lbs. per acre by ground equipment using an amount of water sufficient for adequate coverage. For application by air, use 3 lbs. Cryolite in 20 gallons of water. Apply as needed, but no more than two treatments per crop year. Do not apply later than 30 days prior to harvest.

**Limitations and Precautions:** Growers must inform the processors that Prokil Cryolite 96 has been used on the crop and that raisins must be washed.

**Precaution for use on Table Grapes:** Physical residues may be visible at harvest when application is made on maturing fruit, especially on the darker varieties. Local discretion should be used.

#### NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse or such or in combination with other materials.

Insecticide or Fungicides, Agricultural, Dry, N.O.I.B.N.

# Gowan Company

EPA Reg No 10163-41 AA  
 EPA Est No 48370-DN-01

**NET WEIGHT 50 LBS.**

**BEST AVAILABLE COPY**

5569  
 Box 8090  
 Phoenix, AZ 85066  
 85366

ACCEPTED

DEC 14 1969

10163-41