

9-2-71  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC PEST REGISTRATION  
BY UNDER NO. 10163-42

# PROKIL Perthane Malathion 6-10 Dust

## SUGGESTED USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS; WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

PROKIL Perthane-Malathion 6-10 Dust is suitable for application through a ground duster or by airplane.

### DIRECTIONS

Thorough coverage is essential. Repeat application as necessary.

**BRUSSELS SPROUTS:** Cabbage Worm, Cabbage Looper, Aphids—Apply 25 lbs. per acre. Do not apply within 7 days of harvest.

**CABBAGE, CAULIFLOWER:** Cabbage Worm, Cabbage Looper, Aphids—Apply 25-30 lbs. per acre. Do not apply within 7 days of harvest.

8-71 114A—Proof USDA Reg. No. 10163-42

Active Ingredients	By Wt.
Diethyl Dimethyl Dichloroethane	5.28%
Related Compound	0.72%
Malathion**	10.00%
Inert Ingredients	84.00%
	100.00%

\*\*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

## READ THE LABEL

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered unfit for sale, use or consumption.

SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS!

## NET WEIGHT 50 LBS.

**LETTUCE:** Cabbage Looper, Leafhoppers, Imported Cabbage Worm, Aphids—Apply 25 to 40 lbs. per acre. Repeat at weekly or 10-day intervals as necessary. Do not apply within 7 days of harvest on head lettuce or 14 days on leaf lettuce.

**SPINACH:** Leafhoppers, Cabbage Looper, Aphids, —Apply 20 lbs. per acre at weekly to 10-day intervals. Do not apply within 7 days of harvest.

This product is toxic to fish. Keep out of lakes streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

### NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based on tests believed to be reliable. The use of this product is beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use. The product should be used in accordance with the label or established practice. The purchaser is responsible for proper storage, including injury or damage, including fire, theft, or loss, such as in combination with other materials.

Calif. Reg. No. 10163 50034 AA

CALIFORNIA (Plant)  
P. O. Box 406  
340 East Main Street  
Calipatria, Calif. 92233  
Telephone: 348-5150

# The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA  
Route 1, Box 33  
Somerton, Arizona 85350  
Telephone: 627-2479