

BEST AVAILABLE COPY

NOTE TO PHYSICIAN

ANTIDOTE: If swallowed, Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY! If On Skin, Remove contaminated clothing and immediately wash skin with soap and water

NOTE TO PHYSICIAN: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply under conditions involving possible drift to food, storage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

CRUSH CONTAINER, BURY 18 INCHES DEEP AWAY FROM WATER SUPPLY (WHEN EMPTY). NEVER REUSE.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS, AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. REFER TO STATE AGRICULTURAL EXPERIMENT STATION FOR THE TIMING OF APPLICATIONS.

STORE IN A COOL, DRY PLACE.

KEEP PESTICIDE IN ORIGINAL CONTAINER DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION—Vegetable and Field Crops—20 to 250 gals. Deciduous Fruit Crops—40 to 800 gals. Citrus (Conventional Type Sprayers)—100 to 3,000 gals. (On Young Fruit Plantings—Use a minimum of 20 gals. water per acre.)

Where Recommended for Air Application—Field Crops—5 to 10 gals. Small Fruits—7 to 20 gals. Citrus (mature)—10 to 20 gals. Tree Crops (Mature)—10 to 20 gals.

APPLES: EARLY DELAYED DORMANT ONLY—Green and Reddy Apple Aphids, European Red Mite, San Jose Scale, Fruit Tree Leafhopper—¼ pt plus 2 gals. of a heavy sec. oil (1½ gals. of a very heavy 140 sec. oil in Western States) per 100 gals. water as full coverage sprays. Do not combine oil sprays with Sulfur, Captain or Folpet

PRE-BLOSSOM and POST-BLOSSOM FIRST BROOD COVER SPRAYS—First Brood Codling Moth, Leafhoppers, Green and Reddy Apple Aphids, Leaf Miners—¼ pt per 100 gals. water. Repeat at 7 to 10 day intervals (14 to 21 day intervals in the West). Repeat Application to Prevent Aphid Buildup—1/8 pt. per 100 gals. water. Do not use Phosphamidon on Apples within 30 days of harvest

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

PHOSPHAMIDON 8 SPRAY

Active Ingredients	By Wt.
Phosphamidon	75.5%
Related Compounds	2.5%
Inert Ingredients	22%
	100%

2-Chloro-2-dimethylcarbamoyl-1-methylvinyl dimethyl phosphite
Contains 8 lbs. Phosphamidon insecticide per gallon. U.S. Patent No. 2,989,888. **PROXIL** Phosphamidon 8 Spray is a liquid systemic insecticide for the control of both sucking and chewing insects. It may be applied as a dilute spray or as a concentrate spray.



DANGER - POISON



KEEP OUT OF REACH OF CHILDREN

DO NOT INHALE—DO NOT GET ON SKIN

DO NOT TAKE INTERNALLY

Poisonous if Swallowed, Inhaled or Absorbed Through Skin. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for phosphamidon protection. Respirator Absorbed Through Skin! Do not get in eyes, on skin or clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and water before re-use. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vapored areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Do not use on Maple Trees, Peaches, Plums or Sweet Cherries as injury may occur, avoid drift to these

DO NOT STORE OR USE AROUND THE HOME

NET CONTENTS _____ GALLON LIQUID

GOWAN COMPANY

AGRICULTURAL CHEMICALS

ARIZONA

P.O. Box 9099

12200 E. County 8th St

Telephone 783-8844

Tucson, AZ 85724

EPA Reg No 10163-45 AA

EPA Est No 10163-AZ-1

ACCEPTED
APR 1 1980
Under the Federal Insecticide, and Fungicide Act, as amended, for the pesticide registered under EPA Reg. No. 10163-45

BROCCOLI, CAULIFLOWER: Aphids, Imported Cabbageworm—½ to 1 pint l cover one acre. Apply every 10 to 14 days in a preventative program for aph. Do not apply within 3 days of harvest.

CANTALOUPE, WATERMELON: Aphids, Leaf Miner, Leafhopper—¼ to 1 acre in water to cover. Do not apply within 3 days of harvest.

COTTON: Early Buds (2 through 10 leaf stage)—Aphids, Thrips, Fleahopper, Texas and Oblongate, Spider Mite—Apply 2 or 3 applications of 3 ounce with ground equipment. Spray directly on young plants in sufficient water. Make first application at the 2 true leaf stage. Mid and Late Season—Ly Beetles, Leafhoppers—¼ ounces per acre in water to cover by ground or air. 1 ounce per acre by ground or air. Repeat as necessary. Do not use within 2 harvest.

CUCUMBERS: Aphids, Leaf Miner, Leafhopper—¼ to ½ pt per acre in water. Do not apply within 3 days of harvest

TANGERINES (California and Arizona, Only) ORANGES, GRAPEFRUIT, Apples, Thrips—¼ to ½ pt. per 100 gals. water. (Ground Application)—1 pts. per acre in water to cover; Air Application—½ to ¾ pt per acre in water. Repeat as necessary. Apply during periods of aphid attack on new flush or not apply within 15 days of harvest

PEPPERS: Aphids, Leaf Miners, White Fly—½ pt. per acre in water to cc insects appear. European Corn Borer—1 pt per acre in water to cover when C infestation is expected. Do not apply within 8 days of harvest.

POTATOES: Aphids, Colorado Potato Beetles, Leafhoppers, Leaf Miner, Flea Stinkbug—½ pint per acre in water to cover. Repeat as necessary. Tubers ½ to 1 pt per acre in water to cover by ground (10 gals. by air). Apply prior closing or at first sign of leaf mining. Repeat every 10 to 12 days. Do not app 7 days of harvest

TOMATOES: Aphids, Colorado Potato Beetles, Leaf Miner, White Fly, Flea Stinkbug—½ pt per acre in water to cover. Repeat as necessary. Do not app 10 days of harvest

WALNUTS: Aphids, Codling Moth—1 pt per acre (Ground—400 gals. water. A 20 gals. water). Walnut Nuts Fly—1 pt per acre (Ground—200 to 300 gal. Air—10 to 20 gals. water), applying after egg stings first occur in the husks use within 7 days of harvest

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests below reliable. The use of this product being beyond the control of the manufacturer, expressed or implied, is made as to the effects of such or the result obtained if not used in accordance with directions or established safe practice. Buyer must assume all responsibility, including injury or damage, resulting misuse as such, or in combination with other materials