

LECTICIDE

ER OF CHILDREN

on skin, eyes or clothing. Use water diluting concentrate. May be fatal if eaten. Do not use in areas where children play.

MADE IN U.S.A. REGISTERED TRADEMARK OF THE MANUFACTURER.

ANTIDOTAL. 2-PAM IS ALSO ADMINISTERED IN CONJUNCTION

Company
CHEMICALS

12300 E. 8th Street
P.O. Box 5696

Telephone 783-8844
Tucson, AZ. 85364

PTED

1980

46346

Loopers, Nitidulids, Leafhoppers - 1 pt. per acre in water to cover. Do not use on plants with large leaves. Large Loopers, Stink Bugs - 1 pt. per acre in water to cover.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE AND COLLARDS: Imported Cabbageworm, Diamond Back Caterpillars, Aphids - 1 pt. per acre. Looper - 2 pts. per acre. May be applied within 1 day of harvest on Broccoli, Cauliflower, Flower, Brussels Sprouts and Kale.

COTTON: Lygus, Flea Perforator, Fleahoppers, Armyworms, Bollworms, Loopers and for Suppression of Mites - 1 pt. per acre. Apply at 7 to 10 day intervals when pests persist. For Heavy Insect or Egg Population of any Listed Pest - Use 1 pt. per acre plus residual insecticides for best results. Do not apply within 1 day of hand picking.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Loopers, Thrips, Armyworms - 1 pt. per acre in water to cover. Pickleworms, Leaf Miners - 1 pt. per acre in water to cover. Aerial Applications - Use minimum of 10 gals. water per acre. May be applied up to 1 day of harvest.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miners, Whitefly - 1 pt. per acre. May be applied up to 1 day of harvest.

WALNUTS: Aphids - 1 pt. to 100 gals. water. Do not exceed 10 pts. per acre. Do not apply within 10 days of harvest.

GRAPES: Drosophila, Leafhoppers - 1 to 2 pt. per 100 gals. water. 100 to 150 gal. diluted spray per acre depending on size of vines. Use only 1 pt. per 100 gals. water on Muscat varieties. Do not use on table varieties as fruit injury may occur. Do not use within 4 days of harvest.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm - 1 pt. per acre. Do not use within 1 day of harvest.

CELERY, SPINACH, CHARD, TURNIP GREENS: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners - Use ground application only with fine mist spray. 1 pt. per 100 gals. water per acre on plants less than 4" grown. On larger plants use 1 pt. per 100 gals. water per acre. May be applied up to 1 day of harvest on spinach, Chard and Turnip Greens.

MELONS: Cantaloupes, Honeydew, Muskmelons, Watermelons, PUMPKINS, WINTER SQUASH - Loopers, Armyworms, Leafhoppers, Spider Mites, Leafminers - when temperature is over 80° F. On Honeydew and smooth skin melons, use 1 to 2 pints to 100 gals. water. Do not exceed 2 pints per acre. Do not apply within 1 day of harvest. Do not apply if it rains or are from spray application.

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: Aphids, Soft Brown Scale Crawlers, Citrus Caterpillars, Leaf Roller (Tortrix) - 1 pt. to 100 gals. water. Thrips (Bloom) - 1 pt. per 100 gals. water. Do not apply within 7 days of harvest. May be applied up to 1 day of harvest in California and Arizona.

PEACHES: Western States Only: Lygus, Suppression of mites, peach twig Borer, Oriental Fruit Moth (California Only) - 1 pt. to 100 gals. water per acre. Do not exceed 2 pts. per acre. Do not use within 10 days of harvest. Do not apply when temperature is over 90° F.

RICE: Grasshoppers - 1 to 2 pt. per acre. Apply at first signs of insects. Do not make more than 3 applications. Do not use within 2 days of harvest.

ROSES: (Field Grown): Aphids, Mites, Tent Caterpillars, Thrips, Leaf Rollers - 1 pt. to 100 gals. water.

STRAWBERRIES: Leaf Roller, Spider Mites, Aphids, Omnivorous Leaf Tier, Spittlebugs (Northwest) - 1 pt. per acre. May be applied up to 1 day of harvest.

SUGAR BEETS: Spider Mites, Leafhoppers - 1 pt. per acre. Do not apply within 5 days of harvest.

TOBACCO: Hornworm - 1 pt. to 100 gals. water, using up to 200 gals. diluted spray per acre. Flea Beetles - 1 pt. per acre in water to cover. Apply at first sign of insects. Repeat as necessary.

TOMATOES: (Field) Fruit Flies (Drosophila) - 1 pt. in no less than 10 gals. water per acre with ground sprayer. 1 to 10 gals. per acre. Make first application 5 to 7 days before first picking and continue at 5 to 7 day intervals until harvest. (Field and Greenhouse) Tomato Fruit Worm, Hornworms, Leafminers, Flea Beetles, Mites - 1 pt. to 100 gals. water. Spray plants thoroughly. For Field - Use up to 200 gals. diluted spray per acre. Do not apply when temperature is over 90° F. May be applied up to 1 day of harvest.

GREENHOUSE VAPOR TREATMENT OF ROSES, OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS, Spider Mites, Adult Whiteflies, Aphids, Leafrollers, Mealybugs

1 fluid ounce per 10,000 cu. ft. when plants are dry. Apply undiluted to pipes by means of a plastic squeeze bottle or other dispenser. Apply continuously or to intermittent sections of pipe and on 1 or more of the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 160° F. In steam heated houses, close vents for at least 1 hour after pipes are hot. With hot water systems, close house for at least 3 hours. Houses may remain closed overnight if necessary. Wear goggles and gloves to hot pipes, wear goggles and approved clothing to hot pipes. Do not use in areas where workers are present. Wear respirator when entering greenhouses. Do not use in areas where children are present. Do not use in houses ventilated by fans. Do not use in houses where there is a fire hazard. Do not use in houses where there is a fire hazard. Do not use in houses where there is a fire hazard.

SPIDER MITES: Repeat as necessary. Repeat as necessary. Repeat as necessary.

APHIDS: Apply when aphids appear. Repeat as necessary. **WARNING:** Avoid contact with eyes. Do not apply to plants as injury may result. White Butterfly and other pests may be present. Do not use on plants with yellow flowers. Green Wax on Dew. Do not use on plants with yellow flowers. Do not use on plants with yellow flowers.

FRUIT FLIES: Drosophila spp. in and around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas, Cider Mills and Wineries - Use 2 ounces in 20 gals. water. 5 pints to 100 gals. Apply as a coarse spray to walls, floors, doorways, windows, refuse and cull piles where insects congregate. Apply every 5 to 7 days as long as necessary to maintain control. Do not apply to cull fruit or refuse piles to be fed to livestock. Avoid contamination of pens, feedstuffs and food processing machinery. **DO NOT** use in the edible products areas of food processing plants or other areas where food is commercially prepared or processed.

ADULT MOSQUITO, GNAV, STABLEFLY, DOG FLIES, AND HOUSEFLY CONTROL: RESIDENTIAL AREAS, CROPS LISTED ON LABEL, WOODLANDS, LIVESTOCK PASTURES, FEEDLOTS, AND PASTURES INCLUDING DAIRY CATTLE: It is not necessary to use this product on areas and dairy farms. Make applications during peak of insect activity and repeat as necessary. When using over crop areas clear of the fruit trees and pasture areas specified on the label. Do not use to kill and stallish. Avoid use where they will be used. **AIRCRAFT:** East of Mississippi - Apply 1 to 4 fluid ounces per acre. West of Mississippi - Apply 1 to 4 fluid ounces per acre diluted with water. Apply 1 to 2 quarts of diluted spray per acre. **MIST BLOWER:** Use 1 to 2 pints per 100 gals. water to achieve recommended rate of travel and output, to apply 0.1 to 0.25 lb. per acre. No seed per acre.

CORRAIS, HOLDING PENS: Dairy and Beef Cattle, Hogs, Sheep, Horses, Houseflies, Mosquitoes - Airplane Application - Use 1 to 4 fluid ounces per acre. Do not use in areas where animals are present. Repeat as necessary.

GRASSHOPPERS: Field Areas - crops listed on label - 1 to 2 pints per acre in water to cover. Young Grasshoppers - 1 to 2 pints per acre. Armyworms - 1 to 2 pints per acre. Apply with ground sprayer or backpack. Use thorough coverage of entire area. Repeat as necessary. Do not use in areas where children are present. Do not use in areas where children are present. Do not use in areas where children are present.

NOTICE ON CONDITIONS OF SALE: Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such, or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility regarding injury or damage resulting from its misuse, as such, or its combination with other products.

CORROSIVE LIQUID, N.O.S. - CORROSIVE MATERIAL

**PRECAUTIONARY STATEMENTS
DANGER**

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension or Service.

This Product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do Not apply where runoff is likely to occur. Do Not apply when weather conditions favor drift from areas treated. Do Not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and Crabs may be killed at application rates recommended on this label. Do Not apply where these are important resources. Apply this product only as specified on this label. **DO NOT APPLY** in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do Not use in serving areas while food is exposed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label. Use strictly in accordance with cautions, warnings and directions, and with applicable state and federal regulations. Timing may vary as a consequence of local weather or conditions; we recommend reference to Agricultural Experiment Stations on specific timing.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

PROKIL NALED is a liquid insecticide for the control of both sucking and chewing insects. It may be applied as a conventional diluted spray or as a concentrate spray according to directions. The normal requirements for the following application methods are as follows: Vegetable and Field Crops: Air 5-15 gals. per acre. Ground 20-200 gals. per acre. Small Fruits: Air 7-15 gals. per acre. Ground 20 to 200 gals. per acre. Tree Crops (mature): Air 10-20 gals. per acre. Ground 40-200 gals. per acre. Citrus (mature): Air 10-20 gals. per acre. Ground 100-1500 gals. per acre.

May be applied up to 4 days before harvest, grazing or hand picking unless otherwise noted. Begin application at first sign of insects. Repeat as necessary. Do not use with highly alkaline materials such as lime or bordeaux. **MIXING DIRECTIONS:** Add to spray tank when 1/4 to 1/2 full with agitator running and maintain agitation until spraying is completed.

USE DESIGNATED RATE BY GROUND EQUIPMENT AS A THOROUGH COVERAGE SPRAY. ALFALFA (Seed): Lygus, Mite Suppression [1 application] - 1 pt. per acre. Mites - 1 1/2 pints per acre. 2 applications at 7 day intervals. Thrips - 1/2 pt. per acre. Combine with recommended chemicals for residual control. **DO NOT** apply within 4 days of harvest.

BEANS (Dry or Field), **SOYBEANS**, **LIMA BEANS** AND **PEAS** for Processing: Alfalfa Looper, Aphids, Spider Mites, Lygus, Leaf Miner; **GREEN BEANS:** Aphids, Spider Mites,

**PROKIL
NALED INSECTICIDE**

DANGER

KEEP OUT OF REACH OF CHILDREN

DANGER.

Concentrate causes skin and eye damage. Do not get on skin, eyes or clothing. Use water-proof gloves and face shield or goggles when handling concentrate. May be fatal if swallowed or absorbed through the skin. Do not breathe spray mist. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Avoid contamination of food, foodstuffs and drinking water. **DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.** Do not use on Nectarines, ornamental Cherries, Liquidambar or Chrysanthemums as injury may occur.

NOTE TO PHYSICIAN: ATROPINE IS ANTIDOTAL. 2-PAM IS ALSO ANTIDOTAL AND MAY BE ADMINISTERED IN CONJUNCTION WITH ATROPINE.

Gowan Company

AGRICULTURAL CHEMICALS

12300 E. 8th Street
P.O. Box 5696
Phoenix, Arizona 85044
Telephone 783-8844

EPA Reg. No. 10163-46 AA
EPA Est. No. 10163-AZ-1

ACCEPTED
SEP 25 1980
Under Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10163-46

Loopers, Nitidulids, Leafhoppers, etc. **Large Loopers:** BROCCOLI, CABBAGE, COULARDS: Imported Looper - 2 pts. per acre. Flower, Brussels Sprouts, COTTON: Lygus, Leafhopper, Mite Suppression of Mites, Heavy Insect or Egg Spray for best results. CUCUMBERS, SUMMERS: Thrips, Armyworms per acre in water to be applied up to 1 day before harvest. **EGGPLANT, PEPPER:** Whitefly - 1/2 pt. per acre. **WALNUTS:** Aphids apply within 10 days. **GRAPES:** Drosophila spray per acre depends on varieties. Do not use before harvest. **HOPS:** Aphids, Spider Mites within 4 days of harvest. **CELERY, SPINACH:** Marsh Caterpillars, Leafhoppers. 50 total gals. water per acre. **GREENS:** 50 total gals. water per acre.

MELONS (Cantaloupe, Watermelon): Looper, Aphids. Do not use if temperature is over 90° F. Do not exceed 2 pints per acre to prevent injury from spray. **ORANGES, LEMONS:** Citrus Caterpillars, Leafhoppers. 100 gals. water. Do not use in California and Arizona.

PEACHES (Western): Fruit Moth (California). To avoid fruit spotting, do not use over 90° F.

RICE: Grasshoppers, Leafhoppers. More than 3 applications. 100 gals. water.

ROSES (Field Grown): Aphids, Spider Mites. 100 gals. water.

STRAWBERRIES: 1/2 pt. per acre. (Northwest) - 1 pt. per acre.

SUGAR BEETS: Spider Mites. 100 gals. water.

TOBACCO: Hornworm, Flea Beetles - 1 pt. per acre. 100 gals. water.

TOMATOES: (Field) with ground sprayer, and continue at 5 to 7 day intervals. For Field culture, use thoroughly. For Field culture, use over 90° F.

TICIDE

CHILDREN

es or clothing. Use water-ventrate. May be fatal if mist. In case of contact, s with plenty of water; for stuffs and drinking water TEN FLAME. Do not use nuns as injury may occur

AL. 2-PAM IS ALSO IN CONJUNCTION

any 33

12300 E. 8th Street
P.O. Box 5696
Telephone 783-8844
Tuluma, AZ. 85364

Loopers, Nitidulids, Leafhoppers—1 pt. per acre in water to cover. 100 gals. per acre sprayable. Large Loopers, Stink Bugs—1 1/2 pts. per acre in water to cover.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE AND COLLARDS: Imported Cabbageworm, Diamond Back Caterpillars, Aphids—1 pt. per acre. Looper—2 pts. per acre. May be applied within 1 day of harvest on Broccoli, Cabbage, Cauliflower, Brussels Sprouts and Kale.

COTTON: Lygus, Leaf Perforator, Fleahoppers, Armyworms, Bollworms, Loopers and for Suppression of Mites—1 pt. per acre. Apply at 7 to 10 day intervals while pests persist. For Heavy Insect or Egg Population of any Listed Pest—Use 1/2 pt. per acre plus residual materials for best results. Do not apply within 4 days of hand picking.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Loopers, Thrips, Armyworms—1 pt. per acre in water to cover. Pickleworms, Leaf Miners—1 1/2 pts. per acre in water to cover. Aerial Applications—Use minimum of 10 gals. water per acre. May be applied up to 1 day of harvest.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miners, Whitefly—1 pt. per acre. May be applied up to 1 day of harvest.

WALNUTS: Aphids—1/2 to 1 pt. to 100 gals. water. Do not exceed 10 pts. per acre. Do not apply within 10 days of harvest.

GRAPES: Drosophila, Leafhoppers—1/2 to 1 pt. per 100 gals. water, (100 to 130 gal. diluted spray per acre depending on size of vines). Use only 1/2 pt. per 100 gals. water on Muscat varieties. Do not use on Italia varieties as fruit injury may occur. Do not use within 4 days of harvest.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm—1 pt. per acre. Do not use within 4 days of harvest.

CELERY, SPINACH, CHARD, TURNIP GREENS: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners—Use ground application only with fine mist spray. 1 pt. in 50 total gals. water per acre on plants less than 1/4 grown. On larger plants use 1 1/2 pts. in 100 total gals. water per acre. May be applied up to 1 day of harvest on Spinach, Chard and Turnip Greens.

MELONS (Cantaloupes, Honeydew, Muskmelons, Watermelons), PUMPKINS, WINTER SQUASH: Loopers, Armyworms, Leafhoppers, Spider Mites, Leafminers—when temperature is over 90° F. On Honeydew and smooth skin melons, use 1 to 2 pints to 100 gals. water. Do not exceed 2 pints per acre. Do not apply within 1 day of harvest. Do not apply dust to prevent injury from spray application.

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: Aphids, Soft Brown Scale Crawlers, Citrus Cutworms, Leaf Roller [Tortrix]—1 pt. to 100 gals. water. Thrips [Bloom]—1/2 pt. per 100 gals. water. Do not apply within 7 days of harvest. May be applied up to 1 day of harvest in California and Arizona.

PEACHES (Western States Only): Lygus, Suppression of mites, peach twig Borer, Oriental Fruit Moth [California Only]—1/2 pt. to 100 gals. water per acre. Do not exceed 8 pts. per acre. To avoid fruit spotting, do not use within 30 days of harvest. Do not apply when temperature is over 90° F.

RICE: Grasshoppers—1 to 4 pt. per acre. Apply at first signs of insects. Do not make more than 3 applications. Do not use within 2 days of harvest.

ROSES (Field Grown): Aphids, Mites, Tent Caterpillars, Thrips, Leaf Rollers—1 pt. to 100 gals. water.

STRAWBERRIES: Leaf Roller, Spider Mites, Aphids, Omnivorous Leaf Tier, Spittlebugs [Northwest]—1 pt. per acre. May be applied up to 1 day of harvest.

SUGAR BEETS: Spider Mites, Leafhoppers—1 pt. per acre. Do not apply within 5 days of harvest.

TOBACCO: Hornworm—1 pt. to 100 gals. water, using up to 200 gals. diluted spray per acre. Flea Beetles—1 pt. per acre in water to cover. Apply at first sign of insects. Repeat as necessary.

TOMATOES: (Field) Fruit Flies [Drosophila]—1 pt. in no less than 40 gals. water per acre with ground sprayer. 7 to 10 gals. by air. Make first application 5 to 7 days before first picking and continue at 5 to 7 day intervals until harvest. (Field and Greenhouse): Tomato Fruit Worm, Hornworms, Leafminers, Flea Beetles, Mites—1 pt. in 100 gals. water. Spray plants thoroughly. For Field—Use up to 2 1/2 gals. diluted spray per acre. Do not apply when temperature is over 90° F. May be applied up to 1 day of harvest.

GREENHOUSE VAPOR TREATMENT OF ROSES; OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mites, Adult Whiteflies, Aphids, Leafrollers, Mealybugs—1 fluid ounce per 10,000 cu. ft. when plants are dry. Apply undiluted to pipes by means of a plastic squeeze bottle or other dispenser. Apply continuously or to intermittent sections of pipe and on 1 or more of the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 100° F. In steam heated houses, close vents for at least 1 hour after pipes are hot. With hot water systems, close house for at least 3 hours. Houses may remain closed overnight following treatment. When application is made to hot pipes wear goggles and approved respirator. Ventilate house before workers re-enter. Wear respirator when entering treated houses to ventilate. Protect bees of bees in cucumber and other houses by covering with plastic or rubber sheet during treatment and until house is ventilated. Do not apply to tomatoes or cucumbers closer than 1 day before harvest. Spider Mites [Resistant]—1 No. per acre. Make 3 to 4 treatments at 3 or 4 day intervals. Repeat as necessary. Aphids—Apply when aphids appear. Repeat as necessary. **WARNING:** Avoid direct contact and direct application to plants as injury may result. White Butterfly and Golden Rapture, Roses, Pans, Champagne, Chrysanthemums, Green Wandering Jew, Wincemans and Duchmans Pipe may be injured by NALED vapor treatment.

FRUIT FLIES (Drosophila sp.) in and around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas, Cider Mills and Wineries: Use 2 ounces in 2 1/2 gals. water (5 pints to 100 gals.) Apply as a coarse spray to walls, floors, doorways, windows, refuse and cull piles where insects congregate. Apply every 5 to 7 days as long as necessary to maintain control. Do not apply to cull fruit or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs and food processing machinery. DO NOT use in the edible products areas of food processing plants or other areas where food is commercially prepared or processed.

ADULT MOSQUITO, GNAT, STABLEFLY (Dog Flies), AND HOUSEFLY CONTROL—RESIDENTIAL AREAS, CROPS LISTED ON LABEL, WOODLANDS, LIVESTOCK PASTURES, FEED LOTS, AND PASTURES INCLUDING DAIRY CATTLE: It is not necessary to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. When using over crop areas observe the limitations and restrictions specified for that crop. Poisonous to fish and shellfish. Avoid use where they will be endangered. **AIRCRAFT—East of Mississippi:** Apply 2 1/2 to 4 fluid ounces per acre diluted with water. **West of Mississippi:** Apply 1.6 to 4 fluid ounces per acre diluted with water. Apply 2 to 8 quarts of diluted spray per acre. **MIST BLOWER:** Use 2 1/2 gals. per 100 gals. water. Calibrate equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical Naled per acre.

CORRALS, HOLDING PENS (Dairy and Beef Cattle, Hogs, Sheep, Horses), HOUSES, Mosquitoes—Airplane Application: Use 1.6 to 4 fluid ounces per acre. Dilute 1 to 2 pts. in 10 gals. water. Apply 1 gal. diluted spray per acre. Apply over areas with animals present. Repeat as necessary.

GRASSHOPPERS: Field Areas (crops listed on label)—1 to 4 pints in sufficient water to cover. 1 acre. Young Grasshoppers—1 pint per acre. Armyworms—1 pint per acre. Apply when hoppers hatch is completed. Use thorough cover and contact spray. Repeat as necessary. When using over crop areas observe the limitations and restrictions specified for that crop. Animals may be present during treatment. Forage plants or pastures or field areas should not be cut for hay to be sold or shipped interstate within 1 day of application.

NOTICE ON CONDITIONS OF SALE
Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility including injury or damage, resulting from its misuse as such, or in combination with other materials.

CORROSIVE LIQUID, N.O.S.—CORROSIVE MATERIAL 34