

# PROKIL

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE FOR PROPER TIMING AND NUMBER OF APPLICATIONS. KEEP PESTICIDE IN ORIGINAL CONTAINER!**

### DIRECTIONS

**DO NOT RE-USE EMPTY DRUMS. RETURN TO DRUM RECONDITIONER OR DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.**

When applying Prokil Toxaphene 8 EC by ground use the following dilutions with water. Ground sprayers 20-200 gals. per acre; aircraft 5-15.

**SEED ALFALFA:** For Lygus Bug, use 1-1/2 pt. For **Alfalfa Webworm, Alfalfa Leafhopper and Thrips**, use 2-1/4 pt. For **Armyworm, Grasshopper, Climbing Cutworm, and Alfalfa Looper**, apply 2-1/4 to 3-1/2 pts. per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not use any part of treated crop for food, feed or forage purposes.

**COTTON:** For **Fleahopper and Thrips**, use 3 pints per acre. For **Lygus Bug, Leafworm, Bollworm, Cutworm, and Armyworm**, use 4 pints per acre. Do not graze dairy animals or animals being finished for slaughter in treated fields. Do not apply after bolls open at rates above 4 pints per acre.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** For **Thrips**, use 1-1/2 pts. per acre. **Fall Armyworm**, use 1.6 pts. per acre. For **Imported Cabbageworm**, use 2 pts. per acre. For **Cabbage Looper, Diamond-backed Moth, and Corn Earworm**, apply 2-3 pints per acre. Do not apply after heads or edible parts start to form or within 30 days of harvest.

**CELERY:** **Serpentine Leaf Miner, Thrips and Fall Armyworm**, apply 1.6 pints per acre. For **Imported Cabbageworm**, apply 1-1/2-2 pints per acre. **Cutworms (climbing and surface)** apply 2-5 pints per acre. Do not apply after celery starts to bunch or is half grown.

**EGGPLANT, PEPPERS:** **Armyworm, Blister Beetle, Colorado Potato Beetle, Cucumber Beetle, Flea Beetle, Leafhopper, Serpentine Leaf Miner**. Apply 2 to 3 pints per acre. Do not apply within 5 days of harvest.

## TOXAPHENE 8 EC

### ACTIVE INGREDIENTS:

Toxaphene\* ..... 72.30%  
Aromatic petroleum derivatives ..... 23.20%

INERT INGREDIENTS: ..... 4.50%

TOTAL ..... 100.00%

\*Technical Chlorinated Camphene containing 67-69% chlorine.

Contains 8 lbs. Technical Toxaphene per Gallon

## WARNING

### Keep Out of Reach of Children

### NOT FOR USE OR STORAGE IN OR AROUND HOME

### DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

**May be fatal if swallowed, inhaled or absorbed through skin or eyes. Do not breathe spray or mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid storage near food and feed products, heat or open flame.**

NET CONTENTS.....GALLONS

**LETTUCE:** **Armyworm**, apply 2 pints per acre. **Cutworm and Grasshopper**, use 4 pints per stage on Leaf Lettuce. Do not apply to Head Lettuce.

**CORN, SORGHUM:** **Armyworm**, use 2-1/2 pints per acre. **Climbing Cutworm, and Flea Beetle**, apply 2-3 pints within 40 days of harvest or treat more than 40 days before harvest. Do not feed forage to dairy animals, or animals treated fields. Do not ensile treated sorghum.

**TOMATOES:** **Cabbage Looper, Cutworm (climbing), Thrips, Tomato Hornworm, Tomato Fruitworm, Mite, Pinworm, Colorado Potato Beetle, Armyworm**. Apply 2-5 pints per acre, not later than 15 days before harvest. Do not apply within 5 days of harvest.

**STRAWBERRIES:** **Crown Borer, Leafroller, 1-spotted Weevil**. Apply 2-3 pints per acre. Do not apply within 14 days of harvest.

Do not allow this product to drift onto adjacent crops which may contaminate these crops making them unfit for consumption. Keep clear of food or dairy animals and slaughter.

Toxic to fish and wildlife. Birds feeding in trees and shrubs near lakes, streams or ponds. Do not apply when weather conditions favor drift. Do not apply when weather conditions favor drift. Do not contaminate any body of water by direct application or disposal of wastes and containers. Do not apply to crops not specifically indicated on this label.

This product is toxic to bees and should not be applied to crops visited by bees.

### NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based on tests believed to be reliable. The manufacturer's liability beyond the control of the manufacturer or implied, is made as to the effects of this product obtained if not used in accordance with the directions on the safe practice. The buyer must assume responsibility for injury or damage, resulting from its misapplication with other materials.

ACCEPTED EPA Reg. No. 10163-47 AA

MAY 15 1975

CALIFORNIA  
P.O. Box 458

The Duro Company

# P R O K I L

## TOXAPHENE 8 EC

**ACTIVE INGREDIENTS:**

Toxaphene*	72.30%
Aromatic petroleum derivatives	23.20%
INERT INGREDIENTS:	4.50%
<b>TOTAL</b>	<b>100.00%</b>

\*Technical Chlorinated Camphene containing 67-69% chlorine.  
Contains 8 lbs. Technical Toxaphene per Gallon

### WARNING

**Keep Out of Reach of Children**

**NOT FOR USE OR STORAGE IN  
OR AROUND HOME**

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT  
OR OPEN FLAME**

**May be fatal if swallowed, inhaled or absorbed through skin or eyes. Do not breathe spray or mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid storage near food and feed products, heat or open flame.**

NET CONTENTS.....GALLONS

**STRICTLY IN ACCORDANCE WITH DIRECTIONS, AND WITH APPLICATIONS. TIMING MAY VARY AS LOCAL WEATHER CONDITIONS. WE STATE AGRICULTURAL EXTENSION AND NUMBER OF APPLICATIONS. CONTAINER!**

**DIRECTIONS  
RETURN TO DRUM RECONDITIONING OR CRUSHING AND BURYING**

EC by ground use the following dilutions gals. per acre; aircraft 5-15.

1/2 pt. For Alfalfa Webworm, Alfalfa Leaf-roller, Armyworm, Grasshopper, Climbing Cutworm, 1/4 to 3-1/2 pts. per acre. Do not feed animals being finished for slaughter. Do not feed or forage purposes.

use 3 pints per acre. For Lygus Bug, Leaf-roller, Armyworm, use 4 pints per acre. Do not graze animals being finished for slaughter in treated fields. Do not use 4 pints per acre.

**BROCCOLI, CAULIFLOWER:** For Thrips, use 1-1/2 pints per acre. For Imported Cabbageworm, use 1.6 pts. per acre. For Imported Cabbageworm, Cabbage Looper, Diamond-backed Moth, use 4 pints per acre. Do not apply after heads or edible parts are ready for harvest.

Spring and Fall Armyworm, apply 1.6 pints per acre. For Cutworms, apply 1-1/2-2 pints per acre. Do not apply after celery starts to grow.

Colorado Potato Beetle, Colorado Potato Beetle, Cucumber Beetle, Serpentine Leaf Miner. Apply 2 to 3 pints per acre. Do not apply after heads or edible parts are ready for harvest.

**LETTUCE:** Armyworm, apply 2 pints per acre. For Cabbage Looper, Climbing Cutworm and Grasshopper, use 4 pints per acre. Do not apply after seedling stage on Leaf Lettuce. Do not apply to Head Lettuce after heads begin to form.

**CORN, SORGHUM:** Armyworm, use 2-1/2 pints per acre. For Grasshopper, Climbing Cutworm, and Flea Beetle, apply 2-3 pints per acre. Do not apply to Sorghum within 40 days of harvest or treat more than once after heads start to form. Do not feed forage to dairy animals, or animals being finished for slaughter in treated fields. Do not ensile treated sorghum.

**TOMATOES:** Cabbage Looper, Cutworm (climbing and surface), Flea Beetle, Thrips, Tomato Hornworm, Tomato Fruitworm, Serpentine Leaf Miner, Russet Mite, Pinworm, Colorado Potato Beetle, Stink Bug, Corn Earworm, Beet Armyworm. Apply 2-5 pints per acre, not less than 10 gallons water for air application. Do not apply within 5 days of harvest.

**STRAWBERRIES:** Crown Borer, Leafroller, Tarnished Plant Bug, Strawberry Weevil. Apply 2-3 pints per acre. Do not apply after fruit starts to form.

Do not allow this product to drift onto adjoining food or feed crops as residues may contaminate these crops making them unfit for sale, feed, or human consumption. Keep clear of food or dairy animals, or animals being finished for slaughter.

Toxic to fish and wildlife. Birds feeding in treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes or containers. Do not apply this product to any area or any crop not specifically indicated on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

### NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

**CEP** Reg. No. 10163-47 AA

MAY 15 1975

The Federal Insecticide, Fungicide, and Rodenticide Act, and the pesticide laws of the State of Arizona.

# The Dune Company

**ARIZONA**  
P.O. Box 5696  
2760 W. Cty. 15th St.  
Yuma, Ariz. 85364