

PROKIL

DIMETHOATE W-25 INSECTICIDE

ACTIVE INGREDIENT
Dimethoate, CC dimethyl-S-(N-methylcarbamoyl)methyl
phosphorodithioate 25.0 %
INERT INGREDIENTS 75.0 %
TOTAL 100.0 %

By Wt
25.0 %
75.0 %
100.0 %

CAUTION

KEEP OUT OF REACH OF CHILDREN
— NOT FOR USE OR STORAGE IN OR AROUND HOME —

HARMFUL if Swallowed, Inhaled or Absorbed through Skin
Avoid Breathing dust or spray
Avoid contact with Skin and Eyes
Wash thoroughly after handling
DO NOT contaminate food or feed products

NOTE TO PHYSICIANS: This product upon use may cause cholinesterase inhibition. Atropine or atropine Hydrochloride (2-FAM PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

This product is TOXIC TO WILDLIFE. Birds feeding on treated area may be killed. Keep out of lakes, streams or ponds. DO NOT apply where runoff is likely to occur. DO NOT apply when weather conditions favor drift from treated areas. DO NOT contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

See Back Label for Directions for Use

NET CONTENTS LBS.

CALIFORNIA (Plant)
P.O. Box 459
340 East Main St.
Telephone: 348-2282
Calipatria, CA 92233

The Dune Company
AGRICULTURAL CHEMICALS

ARIZONA
P.O. Box 5696
2760 W. Cty. 15 St.
Telephone 344-0040
Yuma, AZ 85364

This product is TOXIC TO BEES. Restrict application to the period after late when bees are inside the hive or in the early morning before bees are foraging in the field. Protective information may be obtained from your cooperative agricultural extension service.

BAG DISPOSAL: Destroy empty bags by burning. Stay away from smoke and fumes. If burning can not be done safely then render bag unusable and deliver it to an approved disposal site.

— DIRECTIONS FOR USE —

Do Not use this product for any uses other than those specified on this label.

Repeat application, as necessary. Do Not apply when weather conditions favor drifts of spray from areas treated. Consult state experimental station or state extension service for proper timing of applications.

APPLY PROKIL DIMETHOATE W-25 INSECTICIDE in sufficient amount of water to provide sufficient coverage. Conventional ground sprayers—Small fruit 20-200 gals per acre. Tree crops (mature) 40-600 gals per acre. Concentrate Ground Sprayers—Small Fruits and Tree Crops (mature)—30-40 gals per acre.

APPLES, PEARS — Aphids, Leafhoppers, Pear psylla, Mites (except rust mites): Apply 2-4 lb. per 100 gal. water. Do Not apply within 28 days of harvest. Do Not apply during the bloom period.

GRAPE: For raisin and wine grapes and table and canning grapes grown in California to control grape leafhopper and acitic spider mite—apply 6-8 lbs per acre. DO NOT apply within 28 days of harvest. DO NOT apply to canning or table grapes after berries reach 1/4 inch in diameter.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility including injury or damage resulting from its misuse or such, or in combination with other materials.

EPA Reg. No. 10163-CA
8-74-194—PROOF SET

EPA Reg No. 10163-55

