PROKIL DIMETHOATE W-25 INSECTICIDE

ACTIVE INGREDIENT:	% By Wt.
Dimethoate O,O-dimethyl-S-(N-(methylcarbamoyl)methyl)	
phosphorodithioate	25.0%
INERT INGREDIENTS:	75.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

- NOT FOR USE OR STORAGE IN OR AROUND HOME -

HARMFUL if Swallowed, Inhaled or Absorbed through Skin Avoid Breathing Dust or Spray Avoid contact with Skin and Eves

Wash thoroughly after handling

DO NOT contaminate food or feed products

NOTE TO PHYSICIANS: This product upon use may cause cholinesterase inhibition. Atropine is anticottic Pralidoxime chloride (2-PAM; PROTO-PAM chloride) may be energive as an adjunct to atropine. Use according to label directions f


This product is TOXIC TO WILDLIFE. Birds feeding on treated area may be killed. Keep out of labor streams, or ponds. DO NOT apply where runoff it likely to pockur. DO: NET apply when weather conditi. s favor drift from areas treated 1 DO MOT contaminate water by cleaning of equipment on disposed by wester Apply this product only as specified on this label. Function of the transferred to the period after This product of the birst or in the early morning before bees

This produce if the K. TO BEES. Restrict application to the period after dark when the maide the hive or in the early morning before bees are foraging in the field. Protective information may be obtained from your cooperative agricultural extension service.

BAG DISPOSAL: Destroy empty bags by burning. Stay away from smoke and fumes. If burning can not be done safely then render bag unusable and deliver it to an approved disposal site.



AGRICULTURAL CHEMICALS



12900 E. 8th Street P.O. Box 5696 Yuma, Arisona 85964

- DIRECTIONS FOR USE -

Required clothing and equipment for application:

All applicators, including homeowners, flaggers, and all personnel involved with the mixing, loading and transferring operations must wear the protective clothing and equipment enumerated below. Pilots are exempt from this requirement. The protective clothing and equipment to be worn is as follows:

- a) Impermeable gloves (rubber or plastic covered reinforced)
- b) Boots or boot covers
- c) long sleeve shirt and long pants
- d) wide brimmed hats
- e) Respirators must be worn by flaggers and mixer/loaders

Do Not use this product for any uses other than those specified on this label.

Repeat applications as necessary. Do Nct apply when weather conditions favor drifts of spray from areas treated. Consult state experimental station or state extension service for proper timing of applications.

APPLY PROKIL DIMETHOATE W-25 INSECTICIDE BY GROUND ONLY in sufficient amount of water to provide sufficient coverage. Conventional ground sprayers—Small fruit 20-200 gals. per acre. Tree crops—(mature) 40-600 gals. per acre. Concentrate Ground Sprayers—Small Fruits and Tree Crops (mature)—30-40 gals. per acre.

APPLES, PEARS – Aphids, Leafhoppers, Pear psylla, Mites (except rust mites). Apply 1-2 lbs. per 100 gals. water. Do Not apply within 28 days of harvest. Do not apply during the bloom period. (2-lbs. maximum)

GRAPES: For raisin and wine grapes and table and canning grapes grown in California. To control grape leafhopper and pacific spider mite—apply 6-8 lbs. per acre. DO NOT apply within 28 days of harvest. DO NOT apply to canning or table grapes after berries reach ¹/₄ inch in diameter.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility including injury or damage, resulting from its misuse as such, or in combination with other materials.

Y