

DIRECTIONS FOR USE

IMPORTANT: The directions on this label are based on tests and field experience relating to (a) effectiveness; and (b) possible injury to plants and animals; and (c) residues in food, feed, meat and milk. Follow directions for most effective and safest use.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information as the timing will vary with local conditions.

COTTON INSECT CONTROL

Recommended dosages refer to quarts of Prokil Sevimol per acre. Molasses formulation intended for improved bollworm control.

Use 1/2 to 1 quart for early-season thrips, flea beetles, cotton fleahopper, striped blister beetle and cotton leafworm.

Use 1 to 2 quarts for mid-season and late season boll weevil, bollworm, fall armyworm, cotton leaf perforator, leaf rollers, leathoppers, and tarnished plant bug. Also for light-to-moderate infestations of Lygus bug in Western irrigated cotton. Applications should be made every 5 to 7 days as long as control is necessary.

Use 1 1/2 to 2 1/2 quarts for pink bollworm on a 5 to a 7 day schedule depending on the size of the cotton and degree of infestation.

Use 2 quarts for stink bug and salt-marsh caterpillar.

Use 1/2 to 1 1/2 quarts for grasshoppers. The highest dosage should be used when grasshoppers are mature or material is applied on partly defoliated plants.

Aphid populations will be suppressed by repeated application of this insecticide. Prokil Sevimol does not control spider mites. Where infestations are encountered use a recommended miticide.

To avoid staining lint, do not use Prokil Sevimol after bolls open.

FORAGE AND FIELD CROP INSECT CONTROL

Recommended dosages refer to quarts of Prokil Sevimol per acre. Apply when insects or their damage appear and repeat 7 to 14 days later if necessary. Apply sufficient gallonage to obtain full coverage but not less than one gallon or finished spray per acre. See specific directions below for grasshopper control.

SUGAR BEETS

Use 1 to 2 quarts for armyworms, leathoppers and webworms. Do not apply within 14 days of harvest.

GRASSHOPPERS

Apply 1/2 to 1 1/2 quarts as often as necessary to control grasshoppers on the listed forage crops. Follow pre-harvest and grazing use limitations listed for each of the listed crops. A 1/2 to 1 quart rate is suggested for nymphs on small plants or sparse vegetation in wasteland, rangeland, ditchbanks and borders.

A 1 to 1 1/2 quart dosage should be used when grasshoppers are mature or material is applied to crops requiring greater coverage. This information applies to the following forage crops: Alfalfa, soybeans, sorghums and sugarbeets.

ALFALFA AND SOYBEANS

Recommended dosages refer to quarts of Prokil Sevimol per acre.

Use 1/2 to 1 quart for blister beetle and Mexican bean beetle.

Use 1 quart for alfalfa caterpillar, bean leaf beetle, cucumber beetle, gray cloverworm, Japanese beetle, leathoppers, three-cornered alfalfa hopper, thrips and velvet bean caterpillar.

Use 1 to 1 1/2 quarts for armyworms, cutworms, corn earworm (bollworm), stink bugs and webworms.

Use 1 1/2 quarts for control of alfalfa weevil larvae in the spring if pre-treatment damage is extensive, cut the alfalfa and apply to stubble.

Use 1 1/2 quarts for cutworm control on alfalfa.

To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

WARNING! Do not apply a combination of Prokil Sevimol and 2, 4-DB herbicide to soybeans.

SORGHUMS (Including MILO AND GRAIN SORGHUM)

Use 1 to 2 quarts for armyworms, corn earworm, stink bugs and webworms. Use 1 1/2 quarts for sorghum midge and Southwestern corn borer.

Use 2 quarts for cutworm control. For optimum insect control on grain sorghum direct spray into the forming heads. Treat for sorghum midge 3 to 4 days after heads have emerged from boot.

Do not apply within 21 days of harvest of grain. No time limitation on sorghums used for forage.

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND ROBOTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 12163-60 SUBJECT TO ATTACHED COMMENTS

PROKIL Sevimol^(R) 4

CARBARYL INSECTICIDE (Contains 4 Pounds Carbaryl Per Gallon)

ACTIVE INGREDIENT:

Carbaryl

(1-naphthyl methycarbamate) 40.38% by wt

INERT Ingredients 59.62% by wt

100.00% Total

(R) SEVIMOL is the trade mark of Union Carbide Corporation for an insecticide containing Sevin (R) carbaryl and molasses.

U.S. Patent nos. 2,903, 478 and 3,009,855.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful If Inhaled Or Swallowed. Avoid breathing of Spray. Do Not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. DO NOT use 2-PAM, opiates or cholinesterase-inhibiting drugs.

This product may be toxic to Fish, Birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do Not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do Not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to BEES and should not be applied when bees are actively visiting the area. Not to be stored in areas where temperatures frequently exceed 100°F.

Container Disposal: Do Not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke and fumes.

See Back Panel for Directions for Use.

NET CONTENTS _____ GALLONS

The Dune Company
AGRICULTURAL CHEMICALS

CALIFORNIA (Plant)
P.O. Box 458
340 East Main St.
Calipatria, CA 92233
Telephone: 348-2262

ARIZONA
P.O. Box 5696
2750 W. City 15 St.
Yuma, AZ. 85364
Telephone 344-0040

TOBACCO INSECT CONTROL

Special Instructions: When using conventional ground equipment, dilute with water and use 25-40 gallons of finished spray per acre. Apply full coverage.

Use 1 to 2 quarts for budworm, flea beetle, Japanese beetle, June bug and hornworm. Use lower rate on young plants up to knee high; higher rate for mature plants. Begin treatments when worms are small. Avoid excessive application in plant bud as injury may result.

Treated fields may be entered immediately after foliage has dried.

If late applications are necessary, allow 3 days before pruning or cutting.

VEGETABLE INSECT CONTROL

Suggested dosages refer to quarts of Prokil Sevimol per acre. Where a dosage range is indicated, use lower rate on young plants, higher rate on mature plants. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7-to-10-day intervals or as necessary unless a shorter interval is specified below. Apply sufficient gallonage to obtain full coverage but not less than one gallon of finished spray per acre. See specific directions below for grasshopper control.

CORN

Use 1 1/2 quarts for corn earworm, corn rootworm adults, grasshoppers, European corn borer, Southwestern corn borer, fall armyworm, flea beetle, Japanese beetle, sap beetle and leathoppers. Use 2 quarts for Western bean cutworm. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at least every 2 or 3 days starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation; timing and good coverage are absolutely essential for effective control.

Application of Prokil Sevimol to the tassel region of corn during the pollination period will seriously reduce bee populations. Use 2 quarts in at least 15 gallons of water for cutworm control. Spray 12" band over corn row.

Use 2 quarts in at least 15 gallons of water for cutworm control. Spray 12" band over corn row.

BEANS

(Green beans, lima beans, snap beans, cowpeas and black-eyed peas)

Use 1/2 quart for Mexican bean beetle.

Use 1 quart for bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leathoppers, velvet bean caterpillar and Western bean cutworm.

Use 1 to 1 1/2 quarts for armyworms, cutworms, corn earworm, stink bug and tarnished plant bug.

In California, use 2 quarts for corn earworm, lima bean pod borer, Lygus and stink bugs.

POTATO, TOMATO AND PEPPER

Use 1/2 to 1 quart for Colorado potato beetle, flea beetle and leathoppers.

Use 1 to 2 quarts for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bug. Use 2 quarts for cutworm control.

GRASSHOPPERS

Apply 1/2 to 1 1/2 quarts as often as necessary to control grasshoppers on the listed vegetable crops; corn, beans, potato, tomato, and peppers.

SMALL FRUIT INSECT CONTROL

Recommended dosages refer to quarts of Prokil Sevimol per acre. Use sufficient spray gallonage to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7-to-10-day intervals or as necessary.

CRANBERRIES

Use 2 quarts for cutworms, fireworms, fruitworms, Japanese beetle and leathoppers. Apply in late bloom and as needed at 7-to-10-day intervals. A dilute spray of 300 gallons per acre is suggested.

Do not apply within 1 day of harvest.

STRAWBERRIES

Use 1 to 2 quarts for strawberry leaf roller. A dilute spray of 100 to 200 gallons per acre is suggested.

Do not apply within 1 day of harvest.

FEDERAL INSECTICIDE ACT
 10163-60 SUBJECT
 COMMENTS.

PROKIL Sevimol^(R) 4

CARBARYL INSECTICIDE (Contains 4 Pounds Carbaryl Per Gallon)

ACTIVE INGREDIENT:

Carbaryl	(1-naphthyl methycarbamate)	40.38% by wt
INERT Ingredients		59.62% by wt
		100.00% Total

(R) SEVIMOL is the trade mark of Union Carbide Corporation for an insecticide containing Sevin (R) carbaryl and molasses.
 U.S. Patent nos. 2,903,478 and 3,009,855.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful If Inhaled Or Swallowed. Avoid breathing of Spray. Do Not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. DO NOT use 2-PAM, opiates or cholinesterase-inhibiting drugs.

This product may be toxic to Fish, Birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do Not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do Not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to BEES and should not be applied when bees are actively visiting the area. Not to be stored in areas where temperatures frequently exceed 100°F.

Container Disposal: Do Not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke and fumes.

See Back Panel for Directions for Use.

NET CONTENTS _____ GALLONS

The Dune Company
 AGRICULTURAL CHEMICALS

CALIFORNIA (Plant)
 P.O. Box 458
 340 East Main St.
 Calipatria, CA 92233
 Telephone: 348-2282

ARIZONA
 P.O. Box 5696
 2760 W. Cty. 15 St.
 Yuma, AZ. 85364
 Telephone 344-0040

TOBACCO INSECT CONTROL

Special Instructions: When using conventional ground equipment dilute with water and use 25-40 gallons of finished spray per acre. Apply for full coverage.

Use 1 to 2 quarts for budworm, flea beetle, Japanese beetle, June beetle and hornworm. Use lower rate on young plants up to knee high; higher rate for mature plants. Begin treatments when worms are small. Avoid excessive application in plant bud as injury may result. Treated fields may be entered immediately after foliage has dried. If late applications are necessary, allow 3 days before priming or cutting.

VEGETABLE INSECT CONTROL

Suggested dosages refer to quarts of Prokil Sevimol per acre. Where a dosage range is indicated, use lower rate on young plants; higher rate on mature plants. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7-to-10-day intervals or as necessary unless shorter interval is specified below. Apply sufficient gallonage to obtain full coverage but not less than one gallon of finished spray per acre. See specific directions below for grasshopper control.

CORN

Use 1 1/2 quarts for corn earworm, corn rootworm adults, grasshoppers, European corn borer, Southwestern corn borer, fall armyworm, flea beetle, Japanese beetle, sap beetle and leafhoppers. Use 2 quarts for Western bean cutworm. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at least every 2 or 3 days starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation; timing and good coverage are absolutely essential for effective control.

Application of Prokil Sevimol to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

Use 2 quarts in at least 15 gallons of water for cutworm control. Spray in 12" band over corn row.

BEANS

(Green beans, lima beans, snap beans, cowpeas and black-eyed peas)

Use 1/2 quart for Mexican bean beetle. Use 1 quart for bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvet bean caterpillar and Western bean cutworm.

Use 1 to 1 1/2 quarts for armyworms, cutworms, corn earworm, stink bug and tarnished plant bug.

In California, use 2 quarts for corn earworm, lima bean pod borer, Lygus and stink bugs.

POTATO, TOMATO AND PEPPER

Use 1/2 to 1 quart for Colorado potato beetle, flea beetles and leafhoppers. Use 1 to 2 quarts for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs.

Use 2 quarts for cutworm control.

GRASSHOPPERS

Apply 1/2 to 1 1/2 quarts as often as necessary to control grasshoppers on the listed vegetable crops, corn, beans, potato, tomato, and peppers.

SMALL FRUIT INSECT CONTROL

Recommended dosages refer to quarts of Prokil Sevimol per acre. Use sufficient spray gallonage to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7-to-10 day intervals or as necessary.

CRANBERRIES

Use 2 quarts for cutworms, fireworms, fruitworms, Japanese beetle and leafhoppers. Apply in late bloom and as needed at 7-to-10-day intervals. A dilute spray of 300 gallons per acre is suggested.

Do not apply within 1 day of harvest.

STRAWBERRIES

Use 1 to 2 quarts for strawberry leaf roller. A dilute spray of 100 to 200 gallons per acre is suggested.

Do not apply within 1 day of harvest.

IMPORTANT

1. Clean equipment. Flush tank and lines with clear water before and after spraying. Screens should be 50 mesh or coarser. Provide moderate agitation while adding Prokil Sevimol to water in spray or mix tank. Recirculate at intervals to maintain a uniform suspension.

Prokil Sevimol (R) is a suspension of Sevin (R) carbaryl insecticide in molasses for dispersion in water and low volume application in hydraulic sprayers, mist blowers, low gallonage ground equipment and aircraft. Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when Prokil Sevimol is used in accordance with label directions. If Prokil Sevimol (R) is used in accordance with label directions, forage or feed crops may be grazed or harvested for use as feed for dairy or meat animals without resulting residues in meat or milk. **READ THIS LABEL. USE IN STRICT ACCORDANCE WITH DIRECTIONS AND CAUTIONS.**

DIRECTIONS FOR EQUIPMENT CARE, SPRAY PREPARATION AND CONTAINER DISPOSAL

EQUIPMENT CARE:

1. Flush entire spray system with clear water to remove other chemicals before using Prokil Sevimol. DO NOT FLUSH WITH KEROSENE OR DIESEL FUEL.
2. Use 50 mesh or coarser screens in entire system.
3. Nozzles should be fan type, number 3 or larger. Cone nozzles may be used but applying at least 3 gallons of finished spray per acre.
4. After spraying, flush tank and lines with clear water.

SPRAY PREPARATION:

1. Prokil Sevimol may be applied undiluted on cotton. For other crops, see directions 2 through 5.
2. Add required amount of water to spray or mix tank and provide moderate agitation while adding Prokil Sevimol. Refer to specific crop headings for directions on amount of water to use.
3. For combination sprays Prokil Sevimol must be added to at least an equal volume of water and mixed thoroughly before another pesticide is added. (See General Cautions.)
4. Reapply spray mixtures at frequent intervals after mixing to maintain a uniform suspension.
5. Mix only as much spray as can be applied on day of mixing.

STORAGE OF UNUSED MATERIAL AND CONTAINER DISPOSAL:

1. Store unused Prokil Sevimol in original container only, in cool, dry area out of reach of children and animals.
2. Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

GENERAL CAUTIONS

1. Physical compatibility of Prokil Sevimol with other pesticides is not fully known. Incompatible mixtures curdle or precipitate and may injure crops. A compatibility test can be made by adding water in your locality to known herbicides and other formulation ingredients. Do not apply if the mixture is incompatible.
2. Do not mix Prokil Sevimol with nutrients or adjuvants or other pesticides unless an experience operator that the mixture is safe and effective. If unsure, prepare a small amount of the mixture and test on a representative portion of the crop.
3. Irritable under highly alkaline conditions. Not effective if used with alkalis, urea, urea-ammonium, urea-sulfur and urea-lime spreads.
4. To avoid possible injury on tender foliage do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

NOTICE ON CONDITIONS OF SALE

Directions and instructions for use of this product are based upon the best information available to the manufacturer. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury to persons or property.

7-74-191

EPA Est. 10163-CA-1
 Calif Reg No 10163-50054 AA
 EPA Reg. 10163-60

PROKIL Sevimol^(R) 4

CARBARYL INSECTICIDE (Contains 4 Pounds Carbaryl
Gallon)

ACTIVE INGREDIENT:

Carbaryl
1-naphthyl methycarbamate) 40.38% by wt
Inert Ingredients 59.62% by wt
100.00% Total

SEVIMOL is the trade mark of Union Carbide Corporation for an insecticide containing Sevin (R) carbaryl molasses.

Patent nos. 2,903, 478 and 3,009,855.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Dangerous if Inhaled Or Swallowed. Avoid breathing of spray. Do Not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Rinse to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

WARNING FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. DO NOT use 2-PAM, opiates or cholinesterase-inhibiting drugs.

This product may be toxic to Fish, Birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do Not apply in weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do Not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to BEES and should not be applied when bees are actively visiting the area.

Store to be stored in areas where temperatures frequently exceed 100°F.

Container Disposal: Do Not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke and fumes.

See Back Panel for Directions for Use.

NET CONTENTS _____ GALLONS

The Dune Company

AGRICULTURAL CHEMICALS

CALIFORNIA (Plant)
P.O. Box 458
East Main St.
Yupatonia, CA 92233
Telephone: 348-2262

ARIZONA
P.O. Box 5696
2760 W. Cty. 15 St.
Yuma, AZ. 85364
Telephone 344-0040

TOBACCO INSECT CONTROL

Special Instructions: When using conventional ground equipment, dilute with water and use 25-40 gallons of finished spray per acre. Apply for full coverage.

Use 1 to 2 quarts for budworm, flea beetle, Japanese beetle, June beetle and hornworm. Use lower rate on young plants up to knee high, higher rate for mature plants. Begin treatments when worms are small. Avoid excessive application in plant bud as injury may result.

Treated fields may be entered immediately after foliage has dried.

If late applications are necessary, allow 3 days before pruning or cutting.

VEGETABLE INSECT CONTROL

Suggested dosages refer to quarts of Prokil Sevimol per acre. Where a dosage range is indicated, use lower rate on young plants, higher rate on mature plants. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7-to-10-day intervals or as necessary unless shorter interval is specified below. Apply sufficient gallonage to obtain full coverage but not less than one gallon of finished spray per acre. See specific directions below for grasshopper control.

CORN

Use 1½ quarts for corn earworm, corn rootworm adults, grasshoppers, European corn borer, Southwestern corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetle and leafhoppers. Use 2 quarts for Western bean cutworm. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at least every 2 or 3 days starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation; timing and good coverage are absolutely essential for effective control.

Application of Prokil Sevimol to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

Use 2 quarts in at least 15 gallons of water for cutworm control. Spray in 12" band over corn row.

BEANS

(Green beans, lima beans, snap beans, cowpeas and black-eyed peas)

Use ½ quart for Mexican bean beetle.

Use 1 quart for bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvet bean caterpillar and Western bean cutworm.

Use 1 to 1½ quarts for armyworms, cutworms, corn earworm, stink bugs and tarnished plant bug.

In California, use 2 quarts for corn earworm, lima bean pod borer, Lygus and stink bugs.

POTATO, TOMATO AND PEPPER

Use ½ to 1 quart for Colorado potato beetle, flea beetles and leafhoppers.

Use 1 to 2 quarts for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs.

Use 2 quarts for cutworm control.

GRASSHOPPERS

Apply ½ to 1½ quarts as often as necessary to control grasshoppers on the listed vegetable crops: corn, beans, potato, tomato, and peppers.

SMALL FRUIT INSECT CONTROL

Recommended dosages refer to quarts of Prokil Sevimol per acre. Use sufficient spray gallonage to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7-to-10 day intervals or as necessary.

CRANBERRIES

Use 2 quarts for cutworms, fireworms, fruitworm, Japanese beetle and leafhoppers. Apply in late bloom and as needed at 7-to-10-day intervals. A dilute spray of 300 gallons per acre is suggested.

Do not apply within 1 day of harvest.

STRAWBERRIES

Use 1 to 2 quarts for strawberry leaf roller. A dilute spray of 100 to 200 gallons per acre is suggested.

Do not apply within 1 day of harvest.

IMPORTANT

Use clean equipment. Flush tank and lines with clear water before and after spraying. Screens should be 50 mesh or coarser. Provide moderate agitation while adding Prokil Sevimol to water in spray or mix tank. Resuscitate at intervals to maintain a uniform suspension.

Prokil Sevimol (R) is a suspension of Sevin (R) carbaryl insecticide in molasses for dispersion in water and low volume application in hydraulic sprayers, mist blowers, low gallonage ground equipment and aircraft. Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when Prokil Sevimol is used in accordance with label directions. If Prokil Sevimol (R) is used in accordance with label directions, forage or feed crops may be grazed or harvested for use as feed for dairy or meat animals without resulting residues in meat or milk. READ THIS LABEL. USE IN STRICT ACCORDANCE WITH DIRECTIONS AND CAUTIONS.

DIRECTIONS FOR EQUIPMENT CARE, SPRAY PREPARATION AND CONTAINER DISPOSAL

EQUIPMENT CARE:

- 1) Flush entire spray system with clear water to remove other chemicals before using Prokil Sevimol. DO NOT FLUSH WITH KEROSENE OR DIESEL FUEL.
- 2) Use 50 mesh or coarser screens in entire system.
- 3) Nozzles should be fan type, number 3 or larger. Cone nozzles may be used if applying at least 3 gallons of finished spray per acre.
- 4) After spraying, flush tank and lines with clear water.

SPRAY PREPARATION:

- 1) Prokil Sevimol may be applied undiluted on cotton. For other crops, see directions 2 through 5.
- 2) Add required amount of water to spray or mix tank and provide moderate agitation while adding Prokil Sevimol. Refer to specific crop headings for directions on amount of water to use.
- 3) For combination sprays Prokil Sevimol must be added to at least an equal volume of water and mixed thoroughly before another pesticide is added (See General Cautions).
- 4) Resuscitate spray mixtures at frequent intervals after mixing to maintain a uniform suspension.
- 5) Mix only as much spray as can be applied on day of mixing.

STORAGE OF UNUSED MATERIAL AND CONTAINER DISPOSAL:

- 1) Store unused Prokil Sevimol in original container only, in cool, dry area, out of reach of children and animals.
- 2) Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

GENERAL CAUTIONS

- 1) Physical Compatibility of Prokil Sevimol with other pesticides is not fully known. Incompatible mixtures curdle or precipitate and may injure crops or reduce control. Compatibility can be affected by water in your local area, surfactants and other formulation ingredients. Do not apply if physically incompatible.
- 2) Do not mix Prokil Sevimol with nutrients or adjuvants unless pesticides manufacturer's experience indicates that the mixture is safe and effective. If you are unsure, prepare a small amount of the mixture and test on a representative portion of the crop.
- 3) Do not use under highly alkaline conditions. Not to be used with alkaline materials such as Bordeaux lime-sulfur and lime-line sprays.
- 4) The worst possible injury on tender foliage do not apply when foliage is wet or when rain or excessive humidity is expected during the next 24 hours.

NOTICE ON CONDITIONS OF SALE

Directions and formulations for use of this product are based upon tests conducted to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of which or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

7-74-191

EPA Est. 10163-CA-1
Calif. Reg. No. 10163-50654 AA
EPA Reg. 10163-60