

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION

10165-62 PM 12, 176

PROKIL PARATHION 25WP

Active Ingredients:	By Wt.
Parathion: O,O-diethyl O-p-nitrophenyl phosphorothioate.....	23.86%
Related Active Isomers of O,O-diethyl O-p-nitrophenyl phosphorothioate.....	1.14%
Inert Ingredients:.....	75.00%
Total	<u>100.00%</u>

STOP - READ LABEL BEFORE USING!

KEEP OUT OF REACH OF CHILDREN

POISON

See Side Panels for Additional Precautions



DANGER

PELIGRO

NOT FOR HOME USE

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

CALL a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Parathion and describe his condition.

IF BREATHING HAS STOPPED--Start artificial respiration immediately and maintain until doctor sees victim.

IF SWALLOWED--Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IN CASE OF CONTACT--Immediately flush eyes or skin with plenty of water while removing contaminated clothing and shoes. See doctor immediately

ACCEPTED

FEB 24 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 10165-62

Net Weight _____ Pounds

Gowan Company

AGRICULTURAL CHEMICALS

EPA Reg. No. 10165-62 AA
EPA Est. No. 10165-AZ-1

10165-62
P.O. Box 3075
Yuma, AZ 85304
(602) 781-5511



DANGER -- POISON -- PRECAUTIONS

POISONOUS IF SWALLOWED

This product can kill you if swallowed even in small amounts: spray mist or dust may be fatal if swallowed.

POISONOUS BY SKIN CONTACT

Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes.)

POISONOUS IF BREATHED

Breathing vapors, spray mist or dust may be fatal.

NOTE TO PHYSICIAN

Antidote-- administer atropine sulfate in large doses, TWO to FOUR mg. intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

2/6

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER--POISON

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING:

(1) Wear water-proof pants, coat, hat, rubber boots or rubber over-shoes. (2) Wear safety goggles. (3) Wear mask or respirator approved by the Mining Enforcement and Safety Administration for parathion protection. (4) Wear heavy duty, natural rubber gloves.

WORK SAFETY RULES

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Keep all unprotected persons and children away from treated area or where there is a danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away (see Statement of Practical Treatment section).

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash face, hands and arms with soap and water before smoking, eating or drinking. AFTER WORK, take off all work clothes and shoes. Shower, using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.



POISON SIGNS (Symptoms)

Parathion is a very dangerous poison. It rapidly enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: headache, nausea, vomiting, cramps, weakness, blurred vision, pin-point pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your cooperative Agricultural Extension Service.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas for 48 hours unless appropriate protective clothing is worn. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and oral warnings must include the following information: DANGER-- Area treated with Prokil Parathion 25 WP on (date of application). Do not enter without appropriate protective clothing for 48 hours. In case of accidental exposure, follow instructions under Statement of Practical Treatment section.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE water, food or feed by storage or disposal.
PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CROP USE DIRECTIONS

Use only according to instructions listed on the attached booklet/folder.

(Don't lift wags)

PROKIL

Parathion 25 WP

Directions For Use

GOMAN COMPANY
P.O. Box 5696
Yuma, AZ 85364

EPA Reg. No. 10163-62
EPA Est. No. 10163-AZ-1

OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops. In order that pesticide residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to be applied to areas occupied by unprotected humans or beneficial animals or onto adjacent food fiber or pasture crops. The grower is responsible for residues on his crop as well as for damages caused by drift to other property.

Note to Users:

Do not permit workers to enter treated area to engage in any activity requiring substantial contact with treated foliage for the following periods: CITRUS, up to 16 lbs. per acre per year—21 days; greater than 16 lbs. but not more than 30 lbs. per acre per year—30 days; more than 30 lbs. per acre per application or more than 40 lbs. per acre per year—45 days. Grapes, Nectarines Peaches—61 days.

DIRECTIONS FOR USE:

Applications can be made by aircraft or ground power equipment by trained personnel only using approved protective equipment. Measure the specific amount of this product then wash through filter screen in neatly filled spray tank or premix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spray operations. Do not allow mixture to stand. Do not use in low volume gear-type spray equipment. **DO NOT APPLY with hand equipment.** Do not combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants.

RECOMMENDATIONS

Unless otherwise specified, rates are given in terms of Parathion 25 WP per 100 gallons of water for fruit and nut trees, for vegetables and field crops. Parathion 25 WP per acre in sufficient water to provide adequate coverage. Strawberries: Dilution Air, 5-15 gals per acre Ground, conventional 20-200 gals per acre. Small Fruits: Air, 7-15 gals per acre Ground 20-200 gals per acre; Tree Crops (mature) Air, 10-20 gals per acre Ground 40-200 gals per acre. Citrus (mature) Air 10-20 gals per acre Ground, 100-1500 gals per acre. Except as specified begin application at first sign of infestation and repeat as needed to maintain control but otherwise use must given for specific crops.

FRUIT AND NUT CROPS

APPLE — For control of codling moth plum curculio (apply at petal fall and every 10-14 days thereafter until control is achieved), orange (larvae) San Jose Forbs, and scurfy scales, red-banded leafroller (second and third brood), woolly apple aphid and mealybug, use 1 lb. Do not apply within 14 days of harvest. Apple pests given below when present are also controlled. Apple aphid, green apple aphid, red-banded leafroller (first brood) For control of bud moth, red bug, fruit tree leafroller, European red mite, two-spotted mite, Pacific mite, Schoenh mite, Willamette mite, and clover mite, use 1/4 lb.

For certain mites, such as two-spotted and Willamette, repeat applications at 7-10 day intervals during the summer months. The directions for control of mites on apples are applicable to other deciduous fruits such as peaches, pears, plums, and plums (240 lb). **CAUTION:** Injury to fruit and foliage may result from use of this material on McIntosh and certain varieties. Consult agricultural authorities.

APRICOTS — For control of codling moth, Oriental fruit moth, apply at petal fall and every 10-14 days thereafter until control is achieved. For control of scale insects, use 1 to 2 1/4 lbs. Use 1/2 lb. for post-bloom borer control. Use 1/2 lb. for bud moth, fruit tree leafroller, spider mite, and aphids as for apple. Do not apply within 14 days of harvest (144 lb).

CHERRY — Bud moth, black cherry aphid, caterpillar, fruit worm, San Jose scale, shot hole borer, western cherry fruit fly, tartar and Oriental fruit moth (apply at petal fall and every 10-14 days thereafter until control is achieved)—1 lb. For San Jose scale apply at time crawlers emerge. For fruit fly control, begin applications at fly emergence and repeat at 7 day intervals. For mites and aphids—1/2 to 3/4 lb. Do not apply within 14 days of harvest (84 lb).

CITRUS (California) — For control of California red scale and yellow scale - use 2 1/2 lbs. or 1/2 lb. with 1 1/2 to 1 3/4 gallons of a light or light medium oil emulsion per 100 gallons in a full coverage spray any time of year. For control of Purple scale use 1 1/2 lbs. or 1/2 lb. with 1 1/2 to 1 3/4 gallons of a light or light medium oil emulsion per 100 gallons in a full coverage spray any time of year. For control of Citricola and Cottony-cushion scale use 6 lbs. per 100 gallons and 100 gallons per acre or 1/2 lb. per 100 gallons in a full coverage spray during August to November, increase dose to 1/2 lb. per 100 gallons during February to March and 1 lb. during May and June applications. For control of black scale, use 1 1/2 lbs. per 100 gallons of water in a full coverage spray during August to November. For control of citrus thrips use 6 lbs. per 100 gallons and 100 gallons per acre or 1 lb. per 100 gallons in full coverage spray at time of petal fall. Repeat in late summer and early fall to protect new growth and prevent scarring of fruit. Do not apply within 14 days of harvest. Do not apply dosages of more than 16 lbs per acre application within 30 days of harvest. Avoid use of citrus peel for food purposes (400 lb).

GRAPES (California) — Grape mealybug—Apply 2 lbs. per 100 gal. Use 300 gals of spray per acre. Apply in late June or early July. Do not apply within 14 days of harvest. Do not apply more than 6 lbs. per acre application after fruit is the size of buckshot (60 lb).

NECTARINES, PEACHES — Plum curculio and Oriental fruit moth (apply at petal fall and every 10-14 days thereafter until control is achieved), collecting insects, Fruit tree leafroller and cottony peach scale—Use 1 lb. For San Jose scale, use 1 1/4 lb. For Peach twig borer, use 2 lb. Green peach aphid and spider mites, use 1/2-3/4 lb. Areas other than Calif. Fu. coverage spray for control of scale insects. Do not apply within 14 days of harvest. Do not apply more than 5 lbs. actual Parathion per acre per year. (20 lb.) California: Do not apply more than 25 lbs. actual Parathion per acre for any application. Do not apply within 21 days of harvest. Do not apply more than once after bloom. Do not apply more than 5 lbs. actual Parathion per acre between Jan 1 and harvest (20 lb.).

BEST AVAILABLE COPY

OLIVE — and black regular applicat
PEAR — use 1 lb. paylla post-blo der app bug, wa Under on Bos apply v
PRUNE — curculio and ev trial is mealy hids an apply v
STRAW — spider s acre in APPL
WALNUT — spider s 7-10 day not app
BEANS: star). L
- Apply to cover harvest
BEETS: (1-star) in sufficient within 15 are to be
BROCCO — worms (ture)-ACI water to days of 1
CABBAGE — 3rd inst Aphids water h days of CARRO acre in APPL use 1/2 COTTO Spider sufficient pect or Workers hours si CUCUM lee, Lea acre in

APPLY within 15 days of harvest. DO NOT APPLY before vining.

LETTUCE: (Head) - Aphids, Loopers (immature), Armyworms (up to 3rd instar), Leafminers - Apply 2 lbs. per acre in sufficient water to cover. DO NOT APPLY within 7 days of harvest.

SQUASH (winter & summer): Fleabeetles, Aphids, Leafminers - Apply 2 lbs. per acre in sufficient water to cover. DO NOT APPLY within 15 days of harvest.

SUGAR BEETS: Aphids, Armyworms, Blister Beetles, Flea Beetles, Leafhoppers, Leafminers, Webworms - Use 2 - 3 2/10 lbs per acre DO NOT APPLY within 15 days of harvest

TOMATOES: Leafminers, Armyworms (up to 3rd instar), Leafhoppers, Red spider mites - Apply 2-4 lbs. per acre in sufficient water to cover DO NOT APPLY within 10 days of harvest

VETCH, ALFALFA, CLOVER: Aphids, Armyworms, Grasshoppers - Apply 1 1/2 lbs per acre in sufficient water to cover. DO NOT APPLY within 15 days of harvest

WHEATS, OATS, BARLEY: Aphids (Greenbug), Armyworms, Grasshoppers - Apply 3 lbs per acre in sufficient water to cover. DO NOT APPLY within 15 days of harvest

CONDITIONS OF SALE

Our recommendations for use of this pro-

duct are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not in accordance with directions or established sale practice. The buyer must assume all responsibility, including injury or damage resulting from its misuse or such or its combination with other materials.

BEST AVAILABLE COPY

11-11-11

11-11-11

2 for 9