

PROKIL

10163-66

REGISTERED
TRADE MARK
M. S. STANLEY
MONTANA INTELLECT
PROPERTY, B.T.

SUGGESTED USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE VARY GREATLY BETWEEN DISTRICTS, AS A CONSEQUENCE OF LOCAL WEATHER AND OTHER CONDITIONS. ALWAYS FOLLOW THE RECOMMENDATIONS OF STATE AGRICULTURAL EXPERIMENT STATION FOR TIMING OF TREATMENTS.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

Apply thoroughly. Do not apply on food crops within 3 days of harvest unless otherwise specified. Remove residues remaining at harvest by washing, brushing or other effective means.

CUCUMBERS, SQUASH, MELONS: Aphids, Spider Mites, Pickleworm, Cucumber Beetles, Flea Beetles—25 to 37 lbs. per acre, repeat as necessary. **Squash:** Squash Vine Borers—30 to 37 lbs. per acre. **PUMPKINS:** Squash Vine Borers—25 to 33 lbs per acre. Apply twice a week to stems and vines at base of plants. Do not apply to Pumpkins within 3 days of harvest, and to Cucumbers, Squash and Melons within 1 day of harvest. Do not apply unless plants are dry.

LETTUCE: Aphids, Mites, Cabbage Looper, Leafhoppers, Climbing Cutworms, Alfalfa Looper, Corn Earworms—25 to 40 lbs. per acre. Do not apply after the seedling stage.

Malathion Cryolite 6-50 Dust

Active Ingredients	By Wt.
Malathion*	6%
Sodium Fluoaluminate	48
Inert Ingredients	46
Total	100
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.	
Fluorine expressed as elemental	25%

READ THE LABEL
KEEP OUT OF REACH OF CHILDREN

CAUTION

Avoid breathing of dust.

Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered unfit for sale, use or consumption.

SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS!

NET WEIGHT 50 LBS.

CABBAGE: Aphids, Cabbage Looper, Leafhopper, Climbing Cutworms and Alfalfa Looper—25 to 40 lbs. per acre. Do not apply after head starts to form.

TOMATOES: Aphids, Spider Mites, Tomato Russet Mites, Trioxiphila, Armyworms, Flea Beetles—20 to 45 lbs. per acre. Do not apply within 1 day of harvest for rates below 33 lbs. per acre. Do not apply within 5 days of harvest for rates in excess of 33 lbs. per acre.

CAUTION

This product will kill fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such or in combination with other materials.

EPA Est 10163-CA-1

EPA Reg No 10163-66

CALIFORNIA (Plant)
P. O. Box 458
340 East Main Street
Calipatria, Calif. 92233
Telephone: 348-2262

The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA
P. O. Box 5696
2760 W. Cty. 15 St.
Yuma, Arizona 85364
Telephone: 344-0040