

# PROKIL DIPEL® 150 DUST

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and oral warnings must include the following information: **CAUTION!!** Area treated with DIPEL on (date of application). Do not enter without appropriate protective clothing until dusts have settled. In case of accidental exposure, wash with soap and water.

## STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## DIRECTIONS FOR USE

DIPEL® 150 DUST is recommended for use by ground or by air. Apply when pests are first noticed and repeat as necessary. Thorough plant coverage is necessary for control.

Bacillus thuringiensis attacks the larva's gut and so must be eaten by the insect to be effective. After eating Bacillus thuringiensis, susceptible larvae generally stop feeding and die within 2 to 3 days. Do not allow dilute sprays to stand in the tank for more than 12 hours.

DIPEL® is the registered trademark of the Agricultural and Veterinary Products Division, Abbott Laboratories

ACTIVE INGREDIENT:	BY WT.
Bacillus thuringiensis, Berliner, Potency of 320 International Units per mg. (at least 0.5 billion viable spores per g.) Equivalent to potency of 0.15 billion International Units per pound	05%
INERT INGREDIENTS	99.95%
TOTAL	100.00%

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

Reasonable care should be exercised to avoid inhalation or contact with open wounds.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not enter treated areas without protective clothing until dusts have settled. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

NET WEIGHT 50 LBS.

**GOWAN COMPANY**  
AGRICULTURAL CHEMICALS

## APPLICATION RATE

**BROCCOLI, CABBAGE, CAULIFLOWER, COLARDS, KALE, LETTUCE, MUSTARD GREENS, TURNIP GREENS:** For the control of Cabbage Looper and Imported Cabbageworm—apply 25 to 40 lbs. per acre.

**CELERY:** For the control of Cabbage looper and celery looper—apply 25 to 40 lbs. per acre.

**GRAPES:** For the control of Grape Leaf Folder—apply 25 to 40 lbs. per acre.


**MELONS:** For the control of Cabbage looper apply 25 to 40 lbs. per acre.

**POTATOES:** For the control of Cabbage looper—apply 25 to 40 lbs. per acre.

**TOMATOES:** For the control of Cabbage looper and Tomato Hornworms—apply 25 to 40 lbs. per acre.

## NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such or in combination with other materials.



EPA REG. NO. 10163-69  
EPA EST. NO. 10163-AZ-1  
rev 10-3-84

12300 E. Co. 8th St.  
P.O. Box 5696  
Yuma, AZ 85364  
(602) 783-8844

INSECTICIDE OR FUNGICIDE, AGRICULTURAL N.O.I.B.N., DRY.