



115

PROKIN FIFAM 20 W

Fungicide

| | |
|---|----------|
| Active Ingredient: | % by Wt. |
| Firam (Cine dimethyldithiocarbamate)..... | 76.0% |
| Inert Ingredients:..... | 24.0% |
| Total | 100.0% |
| Fine expressed as metallic..... | 10.25% |

ACCEPTED WITH COMMENTS in EPA Letter Dated

JUN 8 1984

| | |
|--|--|
| KEEP OUT OF REACH OF CHILDREN | Under the Federal Insecticide, Fungicide, and Herbicide Act as amended, for the pesticide registration - EPA Reg. No. 100-10163-74 |
| CAUTION | |
| STATEMENT OF PRACTICAL TREATMENT | |
| IF IN EYES, flush eyes with plenty of clear water for at least 15 minutes and get medical attention. IF ON SKIN, wash skin with soap and water. | |

PRECAUTIONARY STATEMENTS
REGARDING HUMANS AND DOMESTIC ANIMALS
CAUTION

Avoid contact with eyes, nose, throat and skin. Avoid breathing dust or spray. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Protective clothing (long pants, long sleeve shirt, gloves, hat and boots) must be worn during mixing and application.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply in weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of this product. Apply this product only as specified on this label.

PHYSICAL AND CHEMICAL HAZARDS

Flammable liquid.

Net weight: 1.0 lb (454 g)

Net content: 1.0 lb (454 g)

Net weight: 1.0 lb (454 g)

115 - E. Co. 8th St.
P. O. Box 5696
Tulsa, OK 74204
(602) 752-8821

BEST DOCUMENT AVAILABLE

... ..

... ..

Use the lower rate

This product may be applied by

... ..

... ..

... ..

... ..

10163-74

APPLES: Brown Rot; Shot-hole Fungus--Use 1 1/2 to 2 lbs. per 100 gallons of water in prebloom, blossom and petal fall through early cover sprays. Under severe disease conditions, or when weather is extremely wet, make additional applications as often as every 5 days until full bloom. DO NOT apply after early cover spray.

BEANS: Rust; Anthracnose--Use 1 1/2 to 2 lbs. per 100 gallons spray per acre when plants are small. Increase dosage to 3 lbs. on large plants. Make applications at 7 day intervals, starting when true leaves appear. DO NOT apply to Snap Beans later than 4 days before picking.

BEETS: Leaf Spot--Use 1 to 2 lbs. per 100 gallons spray per acre. Apply when disease first appears and repeat at 7 to 10 day intervals. DO NOT apply within 7 days of harvest if tops are to be used as food or feed.

BLACKBERRIES (Pacific Northwest): Evergreen Blackberry Canker--Use 1 1/2 to 2 lbs. per 100 gallons of water. Apply once between mid-June and early July. DO NOT apply later than early July.

BLUEBERRIES: Mummy Berry--Use 3 lbs. per 100 gallons at loose bud scale stage and 7 days later. DO NOT apply later than 3 weeks after full bloom.

CABBAGE; CAULIFLOWER: Alternaria Leaf Spot; Downy Mildew--Use 1 1/2 to 2 lbs. per 100 gallons spray per acre. Apply when disease first appears and repeat at 7 to 10 day intervals. DO NOT apply within 7 days of harvest.

CHERRY: Early Blight (Cercospora); Late Blight (Septoria): Use 1 1/2 to 2 lbs. per 100 gallons. Begin in the plant bed and continue in the field at 5-10 day intervals, or at 3-5 day intervals under severe disease conditions. Remove residues by stripping, burning and washing.

CHEERLEAF: Brown Rot; Blossom Blight--Use 1 1/2 lbs. per 100 gallons of water in popcorn, full blossom and petal fall stages. Leaf Spot--Use 1 1/2 lbs. per 100 gallons of water in petal fall and blossom fall stages and approximately 2 weeks after chuck fall. DO NOT apply within 7 days of harvest.

CITRUS: Fruit Rot--Use 1 1/2 lbs. per 100 gallons, applying three times at two week intervals, beginning at mid-bloom. DO NOT apply later than 2 weeks after mid-bloom.

CORN: Leaf Blight, Anthracnose, Rust, Downy Mildew, Fusarium, and other diseases. Use 1 1/2 to 2 lbs. per 100 gallons of spray per acre. Make applications at 7-10 day intervals, beginning when the disease threatens. Continue at 7-10 day intervals as long as the disease threatens. Under severe disease conditions, shorten interval to 5-7 days. After the plants are ripe, the rate can be increased to 3 lbs. per 100 gallons of water per acre.

10163-14

PEACHES; APRICOTS; NECTARINES; Leaf Spot--Use 1 lb. per 100 gallons of water per acre. Begin in plant bed when plants emerge. Repeat in plant bed at 7-10 day intervals or as necessary.

GRAPES: Black Rot; Downy Mildew--Use 1/2 lb. per 100 gallons. Apply before fruit swell. Repeat after blossoming, but before fruit forms. Do NOT apply after fruit begins to form.

NECTARINES; PEACHES: Brown Leaf Spot; Shot-hole Fungus--Use 1 1/2 to 3 lbs. per 100 gallons at dormant and popcorn stages. For West Coast use a Peacher for Leaf Spot, apply just before Fall rains or in November and repeat in three months while trees are still dormant and again in popcorn stage. Brown Rot; Blossom blight--Use 1 1/2 to 2 lbs. per 100 gallons at pink bud and 25% to 75% of full bloom. Fruit Brown Rot--Use 1 1/2 to 2 lbs. per 100 gallons at petal fall, two weeks later, and one or more additional applications at 7-14 day intervals beginning about one month before harvest. If applied closer than 2 weeks before harvest, remove residues by washing or wiping. If lead arsenate is added to the spray, include the correct amount of zinc sulfate and lime to reduce the danger of arsenical injury. Do NOT apply more than 1 lb. per acre after petal fall.

PEACHES: Scab--Use 1 lb. per 100 gallons, applying four times at 11-26 day intervals when tips of small nuts become brown. Under severe disease conditions, a fifth application in August may be required. Do NOT apply after harvest.

STRAWBERRIES: Leaf Spot--Use 1 lb. to 1 1/2 lbs. per 100 gallons. Apply when plants first show signs and again when earliest blossoms open. Gray Mold (Botrytis)--Use 1 lb. to 2 lbs. per 100 gallons. Make 2-3 applications at 7-10 day intervals, starting at early bloom stages. If applied within 7 days of picking, remove residues by washing.

STRAWBERRIES: Anthracnose; Early Blight; Septoria leaf Spot--Use 1 1/2 to 2 lbs. per 100 gallons of water per acre at 7-10 day intervals, starting one week after plants are set. Under severe anthracnose conditions, apply 1 lb. in 100-150 gallons per acre at 7 day intervals for control thereafter.

STRAWBERRIES: Botrytis--Use 1 lb. to 2 lbs. per 100 gallons. Apply when plants first show signs and repeat at 7-10 day intervals thereafter. Do NOT apply after harvest.

STRAWBERRIES: Downy Mildew--Use 1/2 lb. per 100 gallons. Apply before plants emerge and repeat at 7-10 day intervals thereafter. Do NOT apply after harvest.

STRAWBERRIES: Leaf Spot--Use 1 lb. per 100 gallons. Apply when plants first show signs and repeat at 7-10 day intervals thereafter. Do NOT apply after harvest.

10163-74

NOTICE OF CONDITIONS OF SALE

Directions for use of this product are based upon tests
which are reliable. The use of this product being beyond the
control of the manufacturer, no guarantee, expressed or implied,
is made as to the effects of such or the results to be obtained
therefrom in accordance with directions or established safe
practices. The buyer must assume all responsibility, including
damages, resulting from its misuse as such, or in com-
bination with other material.

BEST DOCUMENT AVAILABLE

10F13

| | | |
|---|---------------------------|--------------------------------|
| U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDE PROGRAMS REGISTRATION DIVISION, III 521 WASHINGTON, D.C. 20460 | EPA REGISTRATION NO. | DATE OF ISSUANCE JUN 8 1984 |
| | TERM OF ISSUANCE | |
| NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> RENEWAL <input type="checkbox"/> Reregistration <i>(Under the Federal Insecticide, Fungicide and Rodenticide Act, 7 U.S.C. 136-136f)</i> | NAME OF PESTICIDE PRODUCT | |
| NAME AND ADDRESS OF REGISTRANT (include ZIP code) | | |
| <p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> | | |
| <input type="checkbox"/> ATTACHMENT AVAILABLE | | |
| SIGNATURE OF APPROVING OFFICIAL | | DATE |

LAST COPY AVAILABLE