

Gowan Company

ACCEPTED

SEP 29 1992

 Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 10163-80

BEST AVAILABLE

Information Bulletin

RESTRICTED USE PESTICIDE

DUE TO ACUTE TOXICITY

For retail sale to and use by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING CONTAINING INFORMATION WHICH MAY NOT APPEAR ON PACKAGE LABELS

FOR DISTRIBUTION AND USE ONLY WITHIN LOUISIANA

GOWAN AZINPHOS-M 2 EC

EMULSIFIABLE INSECTICIDE

EPA Reg. No. 10163-80

All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

This supplemental labeling must be in the possession of the user at the time of pesticide application when used on sugarcane in the state of Louisiana.

This supplemental label replaces the directions for use on the container and booklet of this product on sugarcane only in the state of Louisiana. These directions for use must be followed when applying this product to sugarcane in Louisiana.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SUGARCANE (Conventional Spray - Louisiana only): For control of Sugarcane borer apply 3 pints per acre by air application using a minimum of 2 gallons of water per acre. For best results, applications should be made at approximately cane-top level and not more than 10 feet above cane-top level. Do not apply more than 3 times per season nor within 30 days of harvest. Bagasse from sugarcane treated with Gowan Azinphos-M 2 EC may be used for feed to cattle, goats and sheep. Consult your State Agricultural Extension Service or Experiment Station for specific use information.

SUGARCANE (Ultra-Low-Volume-Spray* - Louisiana only): For control of Sugarcane borer apply 3 pints per acre undiluted by air application; 1:1 (water to product). Aircraft must be equipped with low volume device that will produce droplets within the range of 30 to 100 microns. Applications should be made at approximately cane-top level and not more than 10 feet above cane-top level. Do not apply more than 3 times per season nor within 30 days of harvest. Bagasse from sugarcane treated with Gowan Azinphos-M 2 EC may be used for feed to cattle, goats and sheep. Consult your State Agricultural Extension Service or Experiment Station for specific use information.

* **Caution:** This formulation, when used undiluted, may cause spotting of automobile finishes if prolonged exposure is permitted. Do not spray directly over automobiles. If accidental exposure does occur, automobiles should be washed immediately.

RESTRICTIONS

- Do not apply in rain.
- Do not make applications during temperature inversions. A temperature inversion is a stable atmospheric condition characterized by an increase in air temperature with increased height above the ground until at some height a "ceiling" or barrier of colder air is met.
- Make applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph).
- In Louisiana do not apply when the wind velocity exceeds 10 mph.
- For air application, the spray boom must be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vertices. Boom length must not exceed 75% of wing span or rotor diameter.
- Do not apply within 75 feet of lakes, reservoirs, rivers, permanent streams, marshes or ponds, canals, estuaries and commercial fish farm ponds. Do not apply if soil is saturated with water.
- Do not apply under conditions that favor runoff or drift from treated areas.
- Allow at least 21 days between applications and at least 30 days between the last application and harvest.

BEST AVAILABLE COPY