

GOWAN PROMETRYNE 4L

HERBICIDE

ACTIVE INGREDIENT	% BY WT
Prometryne (2,4-bis(isopropylamino)-6-methylthio-s-triazine)	44.4%
Inert Ingredients	55.6%
Total	100.0%

Contains 4 lbs. Prometryne per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If SWALLOWED, do not induce vomiting. Drink 1 or 2 glasses of water or fruit juice with 100 mg. baking soda with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.


- IF INHALED, remove victim to fresh air. Apply artificial respiration if indicated.
 - IF IN EYES, flush eyes with plenty of clear water for at least 15 minutes. Get medical attention if irritation persists.
 - IF ON SKIN, wash with soap and water.
- NOTE TO PHYSICIAN: If ingested, there is no specific antidote. Induce emesis or lavage stomach. Give saline laxative and supportive therapy.

NET CONTENTS _____ GALLONS

Gowan Company

EPA Reg. No. 10163-94
EPA Est. No. 11603 IS 1

P.O. Box 5569
Yuma, Arizona 85366-5569
(602) 783-8844



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, marshes, and potholes). Do not contain more water when disposing of equipment washwaters. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENTS

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and oral warnings must include the following information: CAUTION—Area treated with Prometryne on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, follow instructions under Statement of Practical Treatment section.

CROP USE DIRECTIONS

See directions for use inside booklet

2/89



GOWAN PROMETRYNE 4L

DIRECTIONS FOR USE

Gowan Company

P.O. Box 5696
Yuma, AZ 85364

EPA Reg. No. 10163-94
EPA Est. No. 11602-1S-1

DIRECTIONS FOR USE

This is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Gowan Prometryne 4L is a selective herbicide that may be applied either before or after weeds emerge for control of most annual broadleaf weeds and grasses, including groundcherry, lamsquarters, annual morningglory, mallow, mustard, black night shade, pigweed, karnalsweed, purslane, Florida pusley, ragweed, smartweed, tearweed (prickly sida), barnyardgrass (watergrass), crabgrass, foxtail, goosegrass, junginice, *Fanicum* spp., signalgrass (and other *Bracharia* spp.) and wild oats. Prometryne also controls shallow-germinating seedlings of cocklebur, coffee-weed, and sandspur. It does not control johnsongrass, bermudagrass, other established perennial or sedge/poro, or selective rate.

When applied before weeds emerge, Prometryne enters weeds through their roots. Thus its effectiveness depends on moisture to move it into the soil. Under very dry soil conditions after application, a shallow cultivation or rotary hoeing will generally result in better weed control.

When applied to emerged weeds, Prometryne provides foliar knockdown and/or residual control of later germinating weeds, depending on the rate applied.

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected, we recommend the use of this product in combination with other registered herbicides which are not triazines. Consult with your State Agricultural Extension Service for specific recommendations.

FAILURE TO FOLLOW PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PERFORMANCE, AND/OR ILLEGAL RESIDUES

Equipment and Mixing

Thoroughly clean sprayer prior to use. Do not use a sprayer contaminated with 2,4-D or other materials, as crop damage or sprayer clogging may result.

Use conventional spray equipment with hydraulic or mechanical agitation except in California and Arizona where only mechanical agitators are recommended.

Screens in nozzles and in suction and in line strainers should be no finer than 50 mesh. Use a pump with capacity to maintain 35-40 psi at the nozzles. If hydraulic agitation is used, the pump should also provide sufficient agitation in the tank to keep the mixture in suspension.

Nozzles: For preplant incorporated or preemergence application, use flat fan nozzle tips. For postemergence band application, use off-center nozzle tips. For postemergence broadcast application, use flat fan or off-center nozzle tips. Use flood nozzle tips only in Arizona and California for lay-by treatment in cotton at least 18 inches tall.

Mixing: (1) Fill spray tank 1/2 to 3/4 full with clean water. (2) Start agitation. (3) Pour the product directly from the container into the spray tank partially filled with water, and then add the rest of the water. (4) Provide agitation during mixing and application to maintain a uniform suspension.

Cleaning: Wash sprayer thoroughly with clean water immediately after use. Do not use the same sprayer on sensitive crops, as even small residues of Prometryne in the tank may cause injury to these crops.

Seedbed Preparation

To insure proper placement of Gowan Prometryne 4L, seedbeds must be well prepared and as free as possible from trash and clods. A firm seedbed is best for obtaining effective weed control. Uniformity in height and width of seedbed is essential for proper postemergence applications of Prometryne. Beds should be low and flat. Take care to avoid planter marks. Wear planter packing wheels or rollers are recommended. Wheel furrows should be in firm, dry dirt. Mount the sprayer so that it will be the same row as the planter.

Band Treatment

For broadcast application, use the amount indicated in the tables needed by the soil.

For band application, use the amount indicated in the tables needed by the soil, multiplied by the number of bands per acre of field.

COTTON

Do not allow livestock to feed or graze on treated cotton crops.

Gowan Prometryne 4L may be applied preplant incorporated or preemergence and/or postemergence as recommended in the following tables. The post-emergence applications may follow preplant incorporated or preemergence treatment of Gowan Prometryne 4L.

Apply Gowan Prometryne 4L in a minimum of 20 gals. of water per acre as broadcast or band application at the rates indicated in the following tables.

Do not use on glandless cotton varieties or crop injury will occur.

Note: Do not feed treated forage to livestock, or graze treated areas.

PREPLANT INCORPORATION (Arizona, California, and New Mexico)

Apply Gowan Prometryne 4L at the appropriate rate as a broadcast or band treatment. If broadcast, treat the flat soil surface prior to tilling. If banded, apply over partially finished or finished beds. Incorporate up to 4 inches deep immediately after application with FTO or similar equipment, and a disk rolling, cultivating, or bed conditioner.

Preplant Incorporation

Region	Soil Texture	Broadcast rate per acre
Arizona, California, and New Mexico	sand, loamy sand	Do not use
	sandy loam (AZ & CA only)	24-32 pts
	sandy loam, loams (NM only)	32 pts
	silt loam, clay	48 pts

Precautions: Do not use Gowan Prometryne 4L in our areas of newly leveled fields, in areas of excess soil, or in areas where flooding over the beds is likely to occur or crop injury may result. Do not plant cotton in tractor wheel depressions or crop injury may result. On mulch planted cotton, water back only after cotton seedlings are well established. In New Mexico, apply either preplant incorporated or preemergence (not both)—See Preemergence section.

Gowan Prometryne 4L may be applied aerially and preplant incorporated for weed control in furrow-irrigated cotton (except in California).

BEST AVAILABLE COPY

... applicator only where broadcast applications are specified. Apply at a rate of 5 gallons per acre of total spray. Avoid application under conditions where uniform coverage cannot be obtained—where excessive wind or uneven terrain may occur. To assure that spray will be controlled within the target area, when directed according to label directions, make applications at a minimum height of 10 feet using low drift nozzles at a maximum pressure of 40 psi, and avoid applications during periods when wind speed does not exceed 10 mph. Applications may adversely affect adjacent sensitive nontarget plants. Apply Gowan Prometryne 4L by aircraft at a minimum upward distance of 400 feet.

... of humans or animals. Flagmen and loaders should avoid contact with spray mist and prolonged contact with skin and should wash thoroughly before eating and at the end of each day's operation.

PREEMERGENCE

Apply Gowan Prometryne 4L shortly after planting of the appropriate row. Gowan Prometryne 4L may be used on cotton planted flat or in furrows. To ensure penetration of Prometryne in the seed furrow, do not use broadcast applications on cotton planted in furrows deeper than 2 inches. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow. If banded, do not cover treated bands with soil while cultivating untreated row middles. Do not use on hard or loamy sand, on shallow soils with caliche subsols, or in areas with calciche outcroppings.

Cotton may be replanted in soil previously treated with Gowan Prometryne 4L. Do not apply a second preemergence application of Prometryne or injury may occur.

Preemergence

Region	Soil Texture	Broadcast rate per acre
Arkansas	sandy loam	3.2 pt.
	sandy clay	4.6 pt.
California	loam	4.6 pt.
	clay	4.6 pt.
Florida	loam	4.6 pt.
Georgia	sandy loam	3.2 pt.
	sandy clay	4.6 pt.
Illinois	loam	4.6 pt.
Indiana	loam	4.6 pt.
Mississippi	loam	4.6 pt.
Missouri	loam	4.6 pt.
North Carolina	loam	4.6 pt.
North Dakota	loam	4.6 pt.
Oklahoma	loam	4.6 pt.
South Carolina	loam	4.6 pt.
South Dakota	loam	4.6 pt.
Texas	loam	4.6 pt.
	clay	4.6 pt.
Virginia	loam	4.6 pt.
Washington	loam	4.6 pt.
West Virginia	loam	4.6 pt.
Wisconsin	loam	4.6 pt.
Wyoming	loam	4.6 pt.

*Rio Grande Valley of Texas—Use with caution on soils where water does not drain well. Use less water in shallow cultivation with no ridge weed control.

**New Mexico—Apply either preplant incorporated or preemergence (not both)—See: Preplant Incorporation section.

Winter Weed Control in Texas

For control of winter weeds only, such as henbit (purpletop) and seedling dock, on fall bedded cotton land in the Texas Gulf Coast and Blacklands of Texas, apply 1.2 to 1.6 pt. of Prometryne per acre in the fall or winter to land that will be planted to cotton the following spring. For best results, apply before weeds emerge. Gowan Prometryne 4L will give effective control of emerged henbit if applied before it reaches 4.6 inches tall. For postemergence henbit control, add a suitable surfactant such as X-77® at 0.5% of spray volume or an emul-

sifiable oil at 1.0% of spray volume to the spray tank.

POSTEMERGENCE DIRECTED

Be especially careful in applying Gowan Prometryne 4L postemergence to prevent contact of the spray with cotton leaves, or injury may occur. Use precision application equipment so the spray is accurately directed to the base of the cotton plant and soil thoroughly covers the soil and weeds beneath the cotton plant. Apply during calm periods to prevent drift. Use leaf litter or shields of lead contact cannot be avoided merely by directing the spray. Apply only when drift potential has exceeded the minimum recommended height. Apply to level, well-prepared soil to protect or treat velvetleaf-free beds made with bed shapers. Do not apply to furrow planted cotton until furrows are leveled (plowed in). Do not treat cotton with Prometryne 4L at night, with cold water damage, or in late application.

When applying to emerged weeds, add 2 pt. of surfactant per 100 gal. of spray mixture. Use a surfactant that is compatible with Prometryne 4L when applied in cotton and is approved by EPA for use on food and feed crops. Examples include X-77, Tronic, and Triton.

CHEMICAL HOE (Emerged Weeds Only)

Apply Gowan Prometryne 4L at the appropriate rate two or three times if necessary. In cotton 3 to 6 inches tall, be extremely careful to avoid spray contact with cotton leaves by applying Prometryne 4L with a precision applicator equipped with fenders or shields such as Bell Row Shield, Dickey Fenders, or W&A Fenders. In cotton less than 10 inches tall, apply only if cotton is bed or flat planted.

Chemical Hoe

Height of cotton and area of use	Height of weeds	Broadcast rate per acre
3 to 6 inches	2 to 4 inches	2.4 pt.
Arkansas	2 to 4 inches	3.2 pt.
Mississippi	2 to 4 inches	4.6 pt.
Texas	2 to 4 inches	4.6 pt.
California	2 to 4 inches	4.6 pt.
Florida	2 to 4 inches	4.6 pt.
Georgia	2 to 4 inches	4.6 pt.
Illinois	2 to 4 inches	4.6 pt.
Indiana	2 to 4 inches	4.6 pt.
Mississippi	2 to 4 inches	4.6 pt.
Missouri	2 to 4 inches	4.6 pt.
North Carolina	2 to 4 inches	4.6 pt.
North Dakota	2 to 4 inches	4.6 pt.
Oklahoma	2 to 4 inches	4.6 pt.
South Carolina	2 to 4 inches	4.6 pt.
South Dakota	2 to 4 inches	4.6 pt.
Texas	2 to 4 inches	4.6 pt.
Virginia	2 to 4 inches	4.6 pt.
Washington	2 to 4 inches	4.6 pt.
West Virginia	2 to 4 inches	4.6 pt.
Wisconsin	2 to 4 inches	4.6 pt.
Wyoming	2 to 4 inches	4.6 pt.

LAY BY Emerged Weeds and Germinating Weeds

Apply Gowan Prometryne 4L at the lay by rate once per season when cotton is in the 12 to 18 inch stage, if nozzles are used in Arizona and California. Apply before weeds are 2 inches tall.

Lay-by

Region	Soil texture	Broadcast rate per acre
mid South and Southeast	sandy	2.4 pt.
	loam	2.6 pt.
	clay	3.2 pt.
Blacklands of Texas and Oklahoma	loam	1.6 pt.
	clay	2.6 pt.
High Plains of Texas and New Mexico	clay	1.6 pt.
	loam and clay	2.4 pt.
Southwest Texas	loam	2.4 pt.
	clay	3.2 pt.
Rio Grande Valley of Texas		Do not use
Arizona and California (Do not use in the Coachella Valley)	sand and loamy sand	Do not use
	sandy loam	2.4-3.2 pts
	loam	3.2 pts

Rotational Crops

The following vegetable and cover crops may be planted in the fall when Prometryne 4L was applied on cotton by no more than one of these methods that year: preplant incorporated, preemergence, or only one chemical hoe treatment.

BEST AVAILABLE COPY

vegetables.
Cabbage, cauliflower, pea, and field sweet corn.
Cover Crops: Do not grow down and not used for food or feed.
Cotton: Do not use with barley, water eye, winter wheat.
Spring-seeded crop in California and Arizona and spring-seeded vegetables in the Kern County valley of Texas should not be planted until after April 1.

Gowan Prometryne 4L Combinations for Cotton

A tank mix of Gowan Prometryne 4L and Trifluralin or a tank mix of Gowan Prometryne 4L and Frowl may be applied in a preplant or preemergence application for weed control in cotton.

This tank mix may be applied in a preplant or preemergence application (except in California) to control weeds in cotton where broadcast applications are specified. Apply a minimum of 5 gal. per acre of total spray. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. To assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft using low drift nozzles at a maximum pressure of 40 psi and a total application of 10 periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive non-target plants, apply Gowan Prometryne 4L by aircraft at a minimum upward drift height of 100 feet above the target area. Avoid applications to the ground or plants with a wind speed of 10 mph or greater. Do not apply to plants or ground with a wind speed of 10 mph or greater. Do not apply to plants or ground with a wind speed of 10 mph or greater.

Frowl Tank Mixture: Frowl may be tank mixed with Gowan Prometryne 4L in a preplant or preemergence application for weed control in cotton.

This tank mix may be applied in a preplant or preemergence application (except in California) to control weeds in cotton where broadcast applications are specified. Apply a minimum of 5 gal. per acre of total spray. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. To assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft using low drift nozzles at a maximum pressure of 40 psi and a total application of 10 periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive non-target plants, apply Gowan Prometryne 4L by aircraft at a minimum upward drift height of 100 feet above the target area. Avoid applications to the ground or plants with a wind speed of 10 mph or greater. Do not apply to plants or ground with a wind speed of 10 mph or greater.

Fill tank 1/2 to 3/4 full with clear water and agitate. Premix one part Gowan Prometryne 4L with one part water and add this diluted mixture into the partially filled tank while agitating. After complete mixing, add Frowl to the tank. Fill the remainder of the tank with water while agitating.

Apply the tank mixture with ground equipment at least 10 gal. of water per acre or by air in at least 5 gal. of water per acre according to tank mix directions above in the appropriate label below.

Tank Mixture with Frowl

Soil Texture	Broadcast Rate Per Acre	
	Frowl	Gowan Prometryne 4L
Sandy loam	1.1 5 pts	24.3 2 pts
Loam	1.5 2 pts	3.2 pts
Silty loam, sandy clay loam	1.5 2 pts	3.2 4 8 pts
clay loam, silty clay loam, clay	1.5 3 pts	3.2 4 8 pts

Use the high rate for each soil texture above if heavy weed populations are not present. Use the 3 pt. rate of Frowl for heavy clay soils.

- Precautions:** To avoid crop injury, 1) Do not use in cut areas of newly leveled fields, in areas of excess salt, or in areas where flooding over the bed is likely to occur. 2) Do not plant cotton in tractor wheel depressions. 3) Do not use this tank mixture when cotton is irrigated up. 4) On mulch planted cotton, water back only after cotton seedlings are well established.

Note: Do not feed treated forage to livestock, or graze treated areas, or illegal residues may result.

Rotational Crops: Cotton, sorghum, and other may be replanted. Do not rework the soil before the Frowl label directions in sections of this label for rotational crop restrictions.

Trifluralin Tank Mixture: Trifluralin may be tank mixed with Gowan Prometryne 4L in a preplant or preemergence application for weed control in cotton.

This tank mix may be applied in a preplant or preemergence application (except in California) to control weeds in cotton where broadcast applications are specified. Apply a minimum of 5 gal. per acre of total spray. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. To assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft using low drift nozzles at a maximum pressure of 40 psi and a total application of 10 periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive non-target plants, apply Gowan Prometryne 4L by aircraft at a minimum upward drift height of 100 feet above the target area. Avoid applications to the ground or plants with a wind speed of 10 mph or greater. Do not apply to plants or ground with a wind speed of 10 mph or greater.

Note: Under conditions of very soft water, conduct a compatibility test with Prometryne 4L and Trifluralin to assure compatibility before mixing in spray tank. Follow procedures on the Trifluralin label for soil preparation and incorporation. Apply the tank mix combination to the field before dissolving. Fill Gowan Prometryne 4L directly into spray tank one half to three fourths full of water, allow it to disperse with agitation, add Trifluralin, and then add the rest of the water. Continuous agitation in the spray tank is required to keep the material in suspension. Apply the tank mixture by ground in at least 20 gal. of water per acre or by air in at least 5 gal. of water per acre according to tank mix directions above of the appropriate rate from table below.

Tank Mix with Trifluralin

Soil Texture	Broadcast Rate Per Acre		
	4lb gal Trifluralin	5lb gal Trifluralin	Gowan Prometryne 4L
Sandy loam	DO NOT USE	DO NOT USE	DO NOT USE
Loam	1.1 5 pts	1.1 5 pts	24.3 2 pts
Silty loam	1.5 2 pts	1.5 2 pts	3.2 pts
Sandy clay loam	1.5 2 pts	1.5 2 pts	3.2 4 8 pts
clay loam, silty clay loam, clay	1.5 3 pts	1.5 3 pts	3.2 4 8 pts

Note: Under conditions of very soft water, conduct a compatibility test with Prometryne 4L and Trifluralin to assure compatibility before mixing in spray tank. Follow procedures on the Trifluralin label for soil preparation and incorporation. Apply the tank mix combination to the field before dissolving. Fill Gowan Prometryne 4L directly into spray tank one half to three fourths full of water, allow it to disperse with agitation, add Trifluralin, and then add the rest of the water. Continuous agitation in the spray tank is required to keep the material in suspension. Apply the tank mixture by ground in at least 20 gal. of water per acre or by air in at least 5 gal. of water per acre according to tank mix directions above of the appropriate rate from table below.

Trifluralin Split Application: Trifluralin may be applied in a preplant or preemergence application for weed control in cotton. Apply a minimum of 5 gal. per acre of total spray. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. To assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft using low drift nozzles at a maximum pressure of 40 psi and a total application of 10 periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive non-target plants, apply Gowan Prometryne 4L by aircraft at a minimum upward drift height of 100 feet above the target area. Avoid applications to the ground or plants with a wind speed of 10 mph or greater. Do not apply to plants or ground with a wind speed of 10 mph or greater.

Note: Under conditions of very soft water, conduct a compatibility test with Prometryne 4L and Trifluralin to assure compatibility before mixing in spray tank. Follow procedures on the Trifluralin label for soil preparation and incorporation. Apply the tank mix combination to the field before dissolving. Fill Gowan Prometryne 4L directly into spray tank one half to three fourths full of water, allow it to disperse with agitation, add Trifluralin, and then add the rest of the water. Continuous agitation in the spray tank is required to keep the material in suspension. Apply the tank mixture by ground in at least 20 gal. of water per acre or by air in at least 5 gal. of water per acre according to tank mix directions above of the appropriate rate from table below.

Rotational Crops: Cotton, sorghum, and other may be replanted in the fall after a preplant application of Trifluralin. For winter barley, winter barley, and winter wheat, preplant applications of Trifluralin would be preemergence for winter wheat. Refer to the Trifluralin label for directions and precautions.

Trifluralin Split Application: Trifluralin may be applied in a preplant or preemergence application for weed control in cotton. Apply a minimum of 5 gal. per acre of total spray. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. To assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft using low drift nozzles at a maximum pressure of 40 psi and a total application of 10 periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive non-target plants, apply Gowan Prometryne 4L by aircraft at a minimum upward drift height of 100 feet above the target area. Avoid applications to the ground or plants with a wind speed of 10 mph or greater. Do not apply to plants or ground with a wind speed of 10 mph or greater.

DSMA or MSMA: For faster knockdown of the weeds controlled by Prometryne 4L alone, apply 1.1-3 pints of Prometryne 4L plus 5 lbs. active ingredient of DSMA or 2 lbs. active ingredient of MSMA per acre, following the same directions, precautions, and limitations as given on this label for Prometryne 4L applied alone post-emergence directed (chemical hoe). If needed, make a second application 1-3 weeks after the first application. Do not apply after first bloom.

Several formulations of DSMA and MSMA are available under various trade names from several manufacturers. Observe the directions, limitations, and precautions on the label of the product used.

BEST AVAILABLE COPY

CELERY—EXCEPT IN CALIFORNIA

Seedbeds Florida

Broadcast 2-4 pts. in a minimum of 20 gals. of water per acre after celery has 2-5 true leaves. Application may be made over the celery. Apply only after seedbed covers have been removed from seedbeds for at least one week. Apply only once per year to seedbeds.

Transplants

Apply one application at the appropriate rate from the Table in a minimum of 20 gals. of water per acre during the 2-6 week period after transplanting. Within the rate range given, use the lower rate on relatively coarse textured soils and soils low in organic matter; use the higher rate on relatively fine textured soils and soils high in organic matter. Application may be made over the celery. Apply before weeds are 2 inches tall.

Transplanted Celery

State	Soil	Broadcast rate per acre
Florida	sandy or muck	1.6-3.2 pts
Hawaii	coarse textured	3.2-4.8 pts
	fine-textured	4.8-6.4 pts
Michigan, Ohio	fine-textured or muck	2-4 pts
Wisconsin	fine textured	3.2-4 pts

Rotational Crops

This low-insecticide may be seeded in rotation, applying no more than 4 gals. per acre from 1/4 to 1/2 acre of celery, cabbage, celery, corn, or onion, and not more than ¹ gallon.

STORAGE AND DISPOSAL

Do not allow to enter water, land, or feed by storage or disposal.

PESTICIDE DISPOSAL Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

8/88

BEST AVAILABLE COPY