



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input type="checkbox"/>	Other

OPP Identifier Number
220227

Application for Pesticide - Section I

1. Company/Product Number Gowan Company / 10163-100	2. EPA Product Manager George La Rocca	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Gowan Company / Gowan Diazinon 4E	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) Gowan Company P.O. Box 5569 Yuma, AZ 85366-5569 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION MAR 24 1996
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Addition of advisory statements for Phytotoxicity, Resistance Management and ~~Ornamental~~ Ornamental Use per PR Notice 95-2. Advisory statements have been highlighted.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. _____ No. per container _____	If "Yes" Package wgt. _____ No. per container _____		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Cindy Baker	Title Registration Specialist	Telephone No. (Include Area Code) (520) 783-8844
2. Signature Cindy Baker		6. Date Application Received (Stamped)
3. Title Registration Specialist		
4. Typed Name Cindy Baker	5. Date 2/27/96	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		

2/9

RESTRICTED USE PESTICIDE
DUE TO AVIAN AND AQUATIC TOXICITY
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATORS CERTIFICATION.

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.

GOWAN DIAZINON 4E

Insecticide

For control of certain insects on fruits, nuts, vegetables, field crops, lawns and ornamentals

ACTIVE INGREDIENT: Diazinon	% By Wt.
0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	48.00%
INERT INGREDIENTS*	52.00%
	TOTAL 100.00%

Contains 4 lbs. diazinon per gallon
*Contains xylene range aromatic solvent

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT Organophosphate Insecticide

IF SWALLOWED: Call physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. Do not give anything by mouth if person is unconscious or convulsing.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

IF IN EYES: Rinse eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains a xylene range aromatic solvent which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

FOR 24-HOUR EMERGENCY ASSISTANCE CALL CHEMTREC (800) 424-9300
For other product information, contact Gowan Company or see Material Safety Data Sheet.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Avoid contamination of food and feed. Food utensils such as tablespoons and measuring cups should not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets or livestock. Do not contaminate ornamental fish ponds.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as Barrier Laminate, butyl or nitrile rubber ≥ 14 mils, or Viton ≥ 14 mils
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

EPA Reg. No. 10163-100
EPA Est. No. 67545-AZ-1



Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

GOWAN DIAZINON 4E



Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

EPA Reg. No. 10163-100
EPA Est. No. 67545-AZ-1

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of Atlantic coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping of sprays. If water-fowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill, or store near heat or open flame.

PHYTOTOXICITY STATEMENT

As is common with most emulsifiable concentrate formulations adverse effects, such as spotting or discoloration of the fruit or foliage can occur. Some conditions known to contribute to phytotoxicity include, but are not limited to: high temperatures, poor spray drying conditions, excessive spray runoff, certain spray mixtures, stage of crop development or tank mixes with other pesticides.

RESISTANCE MANAGEMENT STATEMENT

Based on historical use patterns in some areas, certain pest species listed on this label may have developed resistance to Gowan Diazinon 4E. Consult your local agriculture advisor, State Cooperative Extension Service, or regional Gowan Company representative for recommendations.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI). Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves, such as Barrier Laminate, butyl or nitrile rubber \geq 14 mils, or Viton \geq 14 mils
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants in farms, forests, nurseries, or greenhouses.

- Do not permit children or pets to go onto treated areas until sprays have dried.

WORK SAFETY RULES

Repeated exposures to cholinesterase inhibitors such as are contained in this product may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. When handling Gowan Diazinon 4E, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See Practical Treatment section of this label.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

GENERAL INFORMATION

Gowan Diazinon 4E is an emulsifiable solution which, when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops and other plants. Do not use on food crops grown in greenhouses. Do not use in barns or dwellings. Do not use on livestock.

In the directions for use which follow, best control is obtained when application is made at first sign of pests. Application should be repeated only as directed to maintain control.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

Note: The California Department of Agriculture has set a 5-day reentry period for diazinon on grapes, peaches, and nectarines. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

FRUIT AND NUT CROPS- REI 24 hours

Gowan Diazinon 4E may be applied using ground or aerial application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for the various methods of application. The rate of Gowan Diazinon 4E for aerial spray or concentrated spray applied to fruit trees by ground equipment is based on a dilute full cover spray applied with conventional ground equipment at a rate of 300-400 gals. of water per acre. The specific amount of product to be used appears under the separate directions for use for each crop. If the Rate calls for 1 pt. of Gowan Diazinon 4E in 100 gals. of water, this amounts to 3-4 pts. of product per acre when applied at the usual dilute ground spray volumes of 300-400 gals. of water per acre. In no case should the amount of product used per acre exceed the maximum amount per acre per application that is specified in the Directions for individual crops.

Crop	Aerial Application Min. GPA	Ground Application Min. GPA	
		Dilute	Concentrate
Almonds*	20	100	20
Apricots	20	100	20
Blackberries**	20	100	20

Crop	Aerial Application Min. GPA	Ground Application Min. GPA	
		Dilute	Concentrate
Boysenberries**	20	100	20
Cherries	20	100	20
Cranberries	20	15	-
Dewberries**	20	100	20
Grapes	20	100	20
Loganberries**	20	100	20
Nectarines	20	100	20
Peaches	20	100	20
Plums	20	100	20
Prunes	20	100	20
Raspberries**	20	100	20
Strawberries	20	100	20
Walnuts	20	100	20

*CA only.

**CA, OR, or WA only

Observe stated time intervals between last application and harvest, as well as intervals between applications and total amount of product that can be applied per season.

Crop	PHI*
Almonds	**
Apricots	21
Blackberries	7
Boysenberries	7
Cherries	21
Cranberries	7
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums, Prunes	21
Raspberries	7
Strawberries	5
Walnuts	45

*PHI = Pre-Harvest Interval or the number of days between the last application and harvest; may vary based on use pattern selected. See specific directions for use on grapes.

** = Dormant spray only.

Note: To protect bees, do not apply this product to fruit trees when trees or substantial numbers of weeds in the orchard are in bloom.

ALMONDS (California only): San Jose Scale, Parlatoria Scale: Use 1-1 1/2 pts. plus 2-3 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water. Apply as dormant spray. Do not apply more than 6 pts. of diazinon or 9 gals. of oil per acre. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

APRICOTS: Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers: Use 1 pt. in 100 gals. of water. Apply as infestations occur. Repeat application after 14 days if necessary. Olive Scale Crawlers: Use 1/2 pt. plus 1 1/2 gals. of light medium horticultural oil per 100 gals. of water. Apply when scale crawlers are present. Apricot mealybugs: Use 1 pt. per 100 gals. of water. Apply as a cover spray from petal fall to June. Allow 14 days between applications. Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CANEBERRIES (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries) (California, Oregon and Washington only): Leafhoppers, Thrips, Aphids, Two-spotted Spider Mites, Raspberry Fruitworms, Raspberry Sawflies: Use 1 qt. per acre. Apply as insects occur. Repeat application after 14 days if necessary. Do not make more

than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Apply in a minimum of 100 gals. of water per acre (200 gals. per acre maximum). For Raspberry Fruitworms make one application when blossom buds separate. Make a second application immediately before blossoms open. Dryberry Mites: Use 1-2 qts. per acre. Apply as a dormant spray in the early Spring when canes are still on the ground and buds are closed. Make a second application when buds are well developed but still closed. Apply in a minimum of 100 gals. of water per acre. Do not apply within 7 days of harvest or illegal residues may occur. Raspberry Crown Borers (Raspberry Root Borers): Use 2 qts. per acre. Apply a single application in a minimum of 100 gals. of water per acre as a drench to the crown and lower canes in the Spring before buds break. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CHERRIES: Eyespotted Bud Moths, Fruit Tree Leafrollers: Use 1 pt. in 100 gals. of water. Apply in cover spray as infestations occur beginning at petal fall. Repeat application after 7 days if necessary. Leafhopper: Use 1 pt. in 100 gals. of water. Make 3-4 applications beginning at petal fall and continuing through the growing season as infestations occur. Allow 7 days between applications. Cherry Fruitflies: Use 1/2-1 pt. in 100 gals. of water. Make 3-4 applications beginning when adult flies begin to emerge and repeating at 10 day intervals. Black Cherry Aphids: Use 1 pt. in 100 gals. of water. Apply as cover spray when aphids appear. Repeat application after 10 days if necessary. San Jose Scale Crawlers: Use 1 pt. in 100 gals. of water. Apply during the pre-pink stage or when crawlers first appear. Cherry Rust Mites: Use 1/2-1 pt. in 100 gals. of water. Apply to trees as a single supplemental spray after harvest. Notes: (1) Do not make more than 3 applications per season. (2) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (3) Allow a minimum of 21 days between the last application and harvest. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CRANBERRIES: Blackheaded Fireworms (*Rhopobota neavans*): Use 2 qts. in a minimum of 15 gals. of water per acre for ground application and up to 400 gals. of water per acre for chemigation. Apply when pest appears. Repeat application after 14 days if necessary. A maximum of 6 applications per season is permitted. Do not apply within 7 days of harvest or illegal residues may occur. Cranberry Fruitworms, Cranberry Tipworm: Use 3 qts. in a minimum of 15 gals. of water per acre for ground application and up to 400 gals. of water per acre for chemigation. Apply when pest appears. Repeat application after 14 days if necessary. A maximum of 4 application per season is permitted. Do not apply within 7 days of harvest or illegal residues may occur. Note: This product may be applied to cranberries through sprinkler irrigation. Refer to the Chemigation section for specific chemigation directions for use. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

GRAPES: Pacific Spider Mites, Leafhoppers, Grape Berry Moths, Grape Leafhoppers, Omnivorous Leafrollers, Aphids, Spider Mites: Use 1-2 pts. per acre in a minimum of 100 gals. of water. Apply as pest occurs as a thorough cover spray. Repeat application after 7 days if necessary. *Drosophila* spp.: Use 6 fl. oz. per acre in a minimum of 100 gals. of water (300 gals. of water maximum). Apply as pest occurs as a thorough cover spray. Repeat application after 7 days, if necessary. Notes: (1) Do not apply more than 2 pts. of product per acre per application and no more than 10 pts. per acre per season. (2) To avoid illegal residues, allow 28 days between the last application and harvest. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

NECTARINES: Oriental Fruit Moths: Use 1 pt. in 100 gals. of water. Apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight. Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers, Peach Twig Borers: Use 1 pt. in 100 gals. of water. Apply as infestations occur. Repeat application after 14 days if necessary. Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

PEACHES: Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers: Use 1 pt. in 100 gals. of water. Apply as infestations occur. Repeat application after 14 days if necessary. Oriental Fruit Moths: Use 1 pt. in 100 gals. of water. Apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight. Peach Twig Borers: Use 1 pt. in 100 gals. of water. Apply as insects occur in May and June. Allow 7 days between applications. Leafhoppers: Use 1 pt. in 100 gals. of water. Make 3-4 applications beginning at petal fall and continuing through the growing season. Allow at least 7 days between applications. White Peach Scale: Use 1 1/2-2 pts. in 100 gals. of water. Apply post-harvest timed to coincide with peak crawler and immature scale

activity. Under conditions of heavy infestation 2 applications (7 days apart) may be needed. Do not apply before fruit is harvested. Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

PLUMS: Brown Mites (Clover Mites), European Red Mites: Use 1 pt. in 100 gals. of water. Apply as needed every 7 days. Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids: Use 1/2-1 pt. in 100 gals. of water. Apply as needed every 7 days. Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

PRUNES: Brown Mites (Clover Mites), European Red Mites: Use 1 pt. in 100 gals. of water. Apply as needed every 7 days. Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids: Use 1/2-1 pt. in 100 gals. of water. Apply as needed every 7 days. Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

STRAWBERRIES: Aphids, Two-spotted Spider Mites: Use 1 pt. in 100 gals. of water per acre. Apply as pests occur. Repeat applications if necessary every 7 days, but do not make more than 3 applications per season. Do not apply within 5 days of harvest or illegal residues may occur. Cyclamen Mites: Use 1 qt. in 100 gals. of water per acre. Direct spray to the plant crown. Plants should be sufficiently agitated to assure thorough coverage of the foliage. Repeat application after 7 days if necessary. Do not make more than 3 foliar applications per season. Do not apply within 5 days of harvest or illegal residues may occur. Application may be made to plants after harvest. Mole Crickets: Use 1 qt. per acre. Broadcast in sufficient water to obtain even coverage when insects are present. Apply 1-2 days before transplanting and immediately incorporate into the top 1-2 inches of soil. Strawberry Leafrollers: Use 3/4-1 pt. in 100 gals. of water. Apply 200 gals. of spray per acre when blossoms show color. Do not apply within 5 days of harvest or illegal residues may occur. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

WALNUTS (California only): Aphids, Spider Mites: Use 1 qt. per acre. Codling Moths, Scale Crawlers, Walnut Caterpillars: Use 3 qts. per acre. Apply as a foliar spray. Repeat application in 14 days if necessary. Do not make more than 3 applications per season. Do not apply within 45 days of harvest or after husks open or illegal residues may occur. Make ground application in a minimum of 100 gals. of water per acre. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

VEGETABLE CROPS- REI 24 hours

For foliar or soil applications, Gowan Diazinon 4E should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. Gowan Diazinon 4E may be applied using ground or aerial application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for the various methods of application. The specific amount of product to be used appears under the separate directions for use for each crop.

Crop	Ground Application Min. GPA	Min. Gals./ 1000 Sq. Ft. (Ground)	Aerial Application Min. GPA
Beans (Lima, Pole, and Snap) (succulent only)	10	2	5
Beets, Red (Table)	10	2	5
Broccoli	10	2	5
Brussels Sprouts	10	2	5
Cabbage	10	2	5
Cantaloupes	5	1	5
Carrots	10	2	5
Casabas	5	1	5
Cauliflower	10	2	5
Chinese Broccoli	10	2	5

Crop	Ground Application Min. GPA	Min. Gals./ 1000 Sq. Ft. (Ground)	Aerial Application Min. GPA
Chinese Cabbage	10	2	5
Chinese Mustard	10	2	5
Chinese Radish (California and Florida only)	10	2	5
Collards	10	2	5
Crenshaws	5	1	5
Cucumbers	5	1	5
Endive (Escarole)	10	2	5
Ginseng	10	2	5
Honeydew Melons	5	1	5
Kale	10	2	5
Lettuce, Head	10	2	5
Lettuce, Leaf	10	2	5
Muskmelons	5	1	5
Mustard	10	2	5
Onions (Bulb and Green)	10	2	5
Parsley	10	2	5
Parsnips	10	2	5
Peas (succulent only)	10	2	5
Peppers	10	2	5
Persian Melons	5	1	5
*Potatoes (Irish)	10	2	5
Radishes	10	2	5
Spinach	10	2	5
Squash, Summer	5	1	5
Squash, Winter	5	1	5
Sweet Corn	20	4	5
Sweet Potatoes	10	2	5
Swiss Chard	10	2	5
Tomatoes	10	2	5
Turnips	10	2	5
Watermelons	5	1	5

*Do not use on commercially grown potatoes which will be hand-harvested. Observe stated time intervals between last application and harvest as well as intervals between and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (PHI*) for Foliar Applications

Crop	PHI* (Days)
Beans (Lima, Pole, and Snap)	7
Beets, Red (Table)	14
Broccoli	7
Brussels Sprouts	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casabas	3
Cauliflower	7
Chinese Broccoli	10
Chinese Cabbage	10
Chinese Mustard	10
Chinese Radish	10
Collards	10
Crenshaws	3

Pre-Harvest Intervals (PHI)* for Foliar Applications

Crop	PHI* (Days)
Cucumbers	7
Endive (Escarole)	14
Ginseng	30
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Mustard	10
Onions (Bulb and Green)	14
Parsnips	14
Peas	7
Peppers	5
Persian Melons	3
Potatoes (Irish)	35
Radishes	14
Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Swiss Chard	14
Tomatoes	1
Turnips, Roots and Tops	14
Watermelons	3

*PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

Notes: (1) Soil Incorporation: Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see Directions for individual crops) using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches, for subterranean cutworms incorporate 3-6 inches. (2) To Protect Bees: Foliar applications of this product should not be made to peas if these crops or weeds in the treatment area are in bloom, or to corn during the pollen shed period.

BEANS (Lima, Pole, and Snap (succulent only): Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Black Bean Aphids, Mexican Bean Beetles, Dipterous Leafminers: Use 1-1½ pts. per acre. Cucumber Beetles: Use ¾ pt. per acre. Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay. Make cucumber beetle application in 100 gals. of water per acre. **Notes:** (1) Diazinon will not control organophosphate-resistant leafminers. (2) To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

BEANS, LIMA (succulent only): Leafhoppers, Spider Mites: Use 1 pt. per acre. Apply as pests occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay. **Note:** To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

BEETS, RED (Table): Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or field 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Dipterous Leafminers: Use ½-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified

dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

BROCCOLI: Root Maggots: Use 2-3 qts. per acre. Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil. Root Maggots: Use 4-8 oz. in 50 gals. of transplant water as a drench application when 200-300 gals. of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. **Note:** Transplant water treatments may result in stand reduction due to plant stress at time of transplanting. Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers: Use ½-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

BRUSSELS SPROUTS: Root Maggots: Use 2-3 qts. per acre. Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil. Root Maggots: Use 4-8 oz. in 50 gals. of transplant water as a drench application when 200-300 gals. of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted spray equipped with drop nozzles to direct spray to the base of the plant. **Note:** Transplant water treatments may result in stand reduction due to plant stress at the time of transplanting. Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Diamondback Moths, Imported Cabbageworms: Use ½-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CABBAGE: Root Maggots: Use 2-3 qts. per acre. Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil. Root Maggots: Use 4-8 oz. in 50 gals. of transplant water as a drench application when 200-300 gals. of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted spray equipped with drop nozzles to direct spray to the base of the plant. **Note:** Transplant water treatments may result in stand reduction due to plant stress at the time of transplanting. Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers: Use ½-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 21 days of harvest, or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CARROTS: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids: Use 1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest, or illegal residues may occur. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CAULIFLOWER: Root Maggots: Use 2-3 qts. per acre. Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil. Root Maggots: Use 4-8 oz. in 50 gals. of transplant water as a drench application when 200-300 gals. of water are used per acre. Apply ½-1 cup (4-8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. **Note:** Transplant water treatments may result in stand reduction due to plant stress at time of transplanting. Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and

immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre: Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat applications, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 7 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CHINESE BROCCOLI: Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CHINESE CABBAGE (Bok Choy and Napa): Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CHINESE MUSTARD (Gal Choy): Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CHINESE RADISH (Japanese Radish or Daikon) (Florida and California only): Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers: Use 1 pt. per acre. Mix in a minimum of 100 gals. of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

COLLARDS: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Diamondback Moths, Imported Cabbageworms, Harlequin bugs, Corn Earworms: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

CUCUMBERS: Cutworms: Use 2-4 qts. per acre: Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms: Use 1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

ENDIVE (Escarole): Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Dipterous Leafminers: Use 1/2-1 pt. per acre: Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not

7/9
apply within 14 days of harvest. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

GINSENG: Leafhoppers, Aphids, Lygus Bugs, Flea Beetles, Jumping Plant Lice: Use 3/4-1 pt. per acre. Spray when insects first appear. For heavy insect infestations, use 1 pt. per acre. Apply by ground equipment using 10 to 100 gals. of water per acre. Do not apply more than one application per growing season. Do not apply during flowering of three or four year old crops. Do not apply within 30 days of harvest. Do not graze treated areas or feed treated forage to livestock. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

KALE: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Diamondback Moths, Imported Cabbageworms, Harlequin bugs, Corn Earworms: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

LETTUCE (head and leaf): Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Dipterous Leafminers: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

MELONS (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these, and Watermelons): Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Dipterous Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites: Use 1/2-1 1/2 pts. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

MUSTARD: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Dipterous Leafminers, Diamondback Moths, Imported Cabbageworms: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

ONIONS (Bulb and Green): Onion Maggots: Use 2-4 qts. per acre. Broadcast just before planting and mix into the top 3-4 inches of soil. **Note:** Diazinon will not control organophosphate-resistant onion maggots. Onion Thrips: Use 1 pt. per acre. Apply as thrips appear. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

PARSLEY: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before

planting and immediately incorporate into the top 4-8 inches of soil. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

PARSNIPS: Aphids, Flea Beetles, Dipterous Leafminers: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

PEAS (succulent only): Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Dipterous Leafminers: Use 3/4-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. To avoid illegal residues, allow a minimum of 7 days between the last application and harvest. Vines and hay may be fed to dairy and beef cattle and sheep. To avoid illegal residues, wait 7 days following treatment if forage is to be cut for hay. **Notes:** (1) Diazinon will not control organophosphate-resistant leafminers. (2) To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

PEPPERS: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Serpentine Leafminers: Use 1/2-1 pt. per acre. Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 5 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

POTATOES: Cutworms (Surface and Subterranean): Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Southern Potato Wireworms (Southeastern U.S.): Use 2-4 qts. per acre. For potatoes grown on mineral soils in the Southeast, apply before planting and immediately incorporate into the top 3-6 inches of soil depending on location of wireworms. Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leafminers: Use 1/2-3/4 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 35 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. Banded Cucumber Beetles, Leafhoppers, Southern Armyworms: Use 3/4-1 pt. per acre. Apply as insects occur. Repeat application as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 35 days of harvest or illegal residues may occur. **Note:** Do not use on commercially grown potatoes which will be hand-harvested. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

RADISHES: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre: Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Flea Beetles, Dipterous Leafminers: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

SPINACH: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Dipterous Leafminers: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

SQUASH, SUMMER: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites: Use 1/2-1 1/2 pts. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

SQUASH, WINTER: Cutworms: Use 2-4 qts. per acre: Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites: Use 1/2-1 1/2 pts. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

SWEET CORN: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Seed Corn Maggots: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast before planting and immediately incorporate into the top 4-8 inches of soil. Corn Earworms: Use 1-1 1/4 qts. per acre. Apply when silks first appear. Repeat application every 7 days if necessary. Do not make more than 5 applications per season. Corn Rootworm (Adults): Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Corn Leaf Aphids: Use 1-2 pts. per acre. Apply in sufficient water to provide complete coverage to the upper portions of the plants. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Grasshoppers, Spider Mites, Flea Beetles: Use 1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Sap Beetles: Use 2-2 1/2 pts. per acre. Consult local agricultural authorities for proper timing of sprays. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. **Notes:** (1) To protect bees, do not apply foliarly to corn during the pollen shed period. (2) Allow 7 days after the last application before grazing or cutting for forage. Forage may be fed to beef and dairy cattle and sheep. (3) Corn may be picked 7 days after the last application. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

SWEET POTATOES: Wireworms, Flea Beetles: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

SWISS CHARD: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Imported Cabbageworms, Diamondback Moths, Harlequin bugs: Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

TOMATOES: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Dipterous Leafminers: Use 1/2 pt. per acre. Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within one day of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. Banded Cucumber Beetles, Fall Armyworms, Southern Armyworms, Beet Armyworms: Use 3/4-1 pt. per acre. Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within one day of harvest or illegal residues may occur. Vinegar Flies (*Drosophila* spp.): Use 1/2-1 1/2 pts. per acre. Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within one day of harvest or illegal residues may occur. For application by irrigation

systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

TURNIPS, ROOTS and TOPS: Cutworms: Use 2-4 qts. per acre. Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note. Mole Crickets: Use 1 qt. per acre. Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. Wireworms: Use 3-4 qts. per acre. Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. Aphids, Flea Beetles, Dipterous Leafminers. Use 1/2-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. **Note:** Diazinon will not control organophosphate-resistant leafminers. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

FIELD CROPS- REI 24 HOURS

Soil and foliar applications of Gowan Diazinon 4E should always be applied as a spray in sufficient water to ensure thorough coverage. For foliar applications, observe stated time intervals between last application and harvest, as well as intervals between applications and total number of applications that can be made per season.

Pre-Harvest Interval (PHI*) For Foliar Applications

Crop	PHI* (Days)
Hops	14
Sugar Beets	14

PHI* = Pre-Harvest Interval or the number of day between the last application and harvest.

HOPS: Aphids, Spider Mites: Use 1 qt. per acre. Apply as pests occur. Repeat application, as necessary, every 14 days. Do not make more than 4 applications per season. Do not apply within 14 days of harvest. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

SUGAR BEETS: Wireworms: Use 3-4 qts. per acre. Broadcast just before planting. Incorporate into soil 4-6 inches. Do not apply more than 4 qts. of this product to the soil, per season. Beet Leafhoppers, Dipterous Leafminers, Aphids: Use 3/4-1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. **Note:** Diazinon will not control organophosphate-resistant leafminers. Grasshoppers: Use 1 pt. per acre. Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. **Note:** Tops may be fed to beef and dairy animals. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the Supplemental labeling, "APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION."

GRASSLAND INSECTS

DITCH BANKS, ROADSIDES, WASTELAND, NONCROP AREAS, BARRIER STRIPS: GRASSHOPPERS: Spray 3/4-1 pt. per acre in water (minimum of 1 gal. of water per acre) for aerial applications or a minimum of 5 gals. of water per acre for ground applications or in oil (a minimum of 1 gal. of oil per acre) when insects first appear, preferably in the nymphal stage. Thorough coverage of the foliage is essential. Do not graze or use treated grass for forage or feed.

ORNAMENTAL STATEMENT

Before treating a large number of ornamental plants with Gowan Diazinon 4E alone or as a tank mixture with any other material, make a test application on a few plants and observe for 7 to 10 days prior to treating large areas to reduce the possibility of plant injury.

INSECT CONTROL ON ORNAMENTALS- REI 12 HOURS

To control certain insects on Arborvitae, Azalea, Birch, Boxwood, Camellia, Carnation, Chrysanthemum, Douglas Fir, Elm, Gladioli, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Rose, Spruce and Wiltow, apply the recommended rates indicated below. Try to spray underside of leaves and penetrate dense foliage.

Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Dipterous Leafminers, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Obscure Root Weevils, Omnivorous Leafhoppers, Privet Mites, Scale Crawlers (Cottony-cushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Two-spotted Spider Mites, Juniper Webworms, Whiteflies: Use 1/2 fl. oz. per 3 gals. of water, or use 1 pt. per 100 gals. of water. Apple-and-Thorn Skeletonizers, Cotoneaster Webworms, Fall Webworms, Hemlock Chermes, Oak Loopers, Obliquebanded Leafrollers, Pear Slugs, Tent Caterpillars: Use 1 1/2 fl. ozs. per 3 gals. of water, or 3 pts. per 100 gals. of water. Mimosa Webworms: Mix 1/2 fl. oz. in 3 gals. of water

(or 1 pt. in 100 gals. of water) and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

1 fl. oz. = 2 tablespoons;

16 fl. oz. = 1 pint

PRECAUTION: Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, and Gardenia, since injury to the plants may occur.

LAWN PEST CONTROL

(Excluding Golf Courses and Sod Farms)

PRECAUTIONS: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping sprays. If waterfowl, i.e., ducks or geese, can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least 1/4 inch water immediately after application, however, stop watering before puddling occurs.

The following rates for Lawn Pest Control are given as rate per 1,000 sq. ft.

Lawn Chinch Bugs: Use 2-3 fl. ozs. in 3 gals. of water. Use higher rate for longer residual control and in lawns of dense growth such as St. Augustine lawns. Ants, Armyworms, Bermudagrass Mites, Clover Mites, Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs), Springtails (Collembola): Use 2-3 fl. ozs. in 3 gals. of water. Spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk when wasps are not active. For frit flies, mow grass and water well before treatment, but delay application until grass is dry. For billbug control, apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect.

Brown Dog Ticks, Chiggers, Fleas: Use 3 fl. oz. in 3 gals. of water. **Note:** Do not apply to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house. Hyperodes Weevils: Use 3 fl. ozs. in 3 gals. of water. Treat problem areas in mid-April and again in mid-May. Millipedes: Use 3 fl. ozs. in 3 gals. of water. Rhodesgrass Scales: Use 3 fl. ozs. in 3 gals. of water. Apply when crawlers first emerge. White Grubs (such as Japanese Beetle Larvae, European Chafer, Southern Chafer): Use 3 fl. ozs. in 3 gals. of water. Apply any time between late July and early October.

NUISANCE PESTS IN OUTSIDE AREAS

Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (Collembola): Mix 3 fl. ozs. in 3 gals. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide. To prevent entry into houses, spray a five-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary. Do not apply to animals. Do not treat plants (other than tree trunks) at this nuisance pest rate.

For application through irrigation systems, see SUPPLEMENTAL LABELING APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If pesticide, spray mixture or rinsate cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER

Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or incinerate, or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

rev. 2/95