

) 10163-134

) 05/15/2000

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

Rebecca A. Lamas
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

MAY 15 2000

Subject: Gowan Carbaryl 80 WSB
EPA Registration No. 10163-134
Amendment dated February 18, 2000 in Response to EPA
Letter dated December 21, 1999 in Response to GDE.

Dear Ms. Lamas:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes indicated below before you release the product for shipment bearing the amended labeling:

1. This product is in water soluble packets and the label must be changed to indicate the proper formulation and include necessary safety measures. On page 2 under "Applications Recommendations" the statement must be changed to be similar as follows: "Gowan Carbaryl 80 WSB is a dry powder formulation of carbaryl insecticide and is packaged in water soluble paks (WSPs) for dispersion in water and application through hydraulic sprayers, mist blowers, low gallonage ground equipment and aircraft. Each pak contains 1.0 pounds of formulated product. Do not sell individual water soluble paks. Do not handle inner bag with wet hands or gloves. Do not allow paks to become wet prior to adding to the spray tank. Handle outer container (over wrap bag) carefully to avoid breakage of inner soluble paks. Always reseal outer (over wrap bag) in a manner that protects remaining paks from moisture. Do not remove the WSPs from the container except for immediate use. Use the entire contents of a WSP and do not break open to use partial contents of a WSP. Directions for water dilution are given under the CROP USE DIRECTIONS."

2. The directions on page 3 under the heading "Spray Preparation" must be changed to be applicable for water soluble paks. After the statement "Fill mix tank 1/2 to 3/4 full with water and start agitation", delete the next two label sentences and replace them with the following: "Slowly add the required number of unopened paks of Gowan Carbaryl 80 WSB into the mixing tank. Allow all the water soluble paks to dissolve and completely disperse. Continue agitation while

adding the remainder of the water." Then continue with the remainder of the present label directions under this heading.

3. On page 3 under the heading "Dilution Directions" the statement "Where small volumes of spray....per 100 gallons" must be deleted. The water soluble paks must not be opened for partial use as previously stated on the label. Additionally, it is specified this product is for agriculture use only and all use directions are in pounds per acre or pounds per 100 gallons of water.

4. Under the "TREES AND VINES" section make the following corrections:

The correct PHI for almonds is 14 days.
The PHI for apples under "Apple Thinning" should be 3.
After Cherries insert the PHI (3), and state "See PRUNES".
Place the PHI after LOGANBERRIES (7), NECTARINES (3), and PLUMS (3).
Correct the PHIs for the crops PRUNES (3), PLUMS (3), and CHERRIES (3).

5. Under the heading "FIELD AND ROW CROPS" make the following corrections:

Change "BROCCOL (3)I" to "BROCCOLI (3)".
Correct the PHI for PARSNIPS to (7), SWEET POTATOES to (7), and TOBACCO to (0).

6. Under the heading "Further directions for Chemigation systems" on page 8, the first four sentences should be changed to "Fill mix tank half to three-quarters of the desired amount of water. Begin agitation. Add the required amount of Gowan Carbaryl 80 WSB and once the water soluble paks are dissolved then add the remaining volume of water. Maintain agitation until application is complete."

Submit two copies of your final printed labeling before you release the product for shipment. A copy of the labeling stamped "Accepted with Comments" is enclosed for your records.

Sincerely,
George Tompkins
George Tompkins, Ph.D., Entomologist
Insecticide-Rodenticide Branch
Registration Division (7505C)

MAY 15 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
10163-134

GOWAN CARBARYL 80 WSB

For agricultural use only

ACTIVE INGREDIENT:		% BY WT.
Carbaryl (1-naphthyl N-methylcarbamate).....		80.0%
INERT INGREDIENTS.....		20.0%
	TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

PRECAUTION AL USARIO: Si usted no lee ingles, no use este producto hastat que la etiqueta le haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by placing finger on back of throat. Call a physician. Never give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove victim to fresh air. Contact a physician.

IF ON SKIN: Remove contaminated clothing and wash skin with soap and generous amounts of water. Contact a physician if irritation develops.

IF IN EYES: Immediately flush eyes with plenty of water. Contact a physician if irritation develops.

NOTE TO PHYSICIAN: Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible. Gastric lavage may be used if this product has been swallowed. Carbaryl poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

PRECAUTION: To prevent aspiration of petroleum distillates, lay patient on side with head lower than waist. Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. *Mild cases may be given 1 to 2 mg. Intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms appear. Severe cases should be given 2 to 4 mg. Intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.*

NARCOTICS AND OTHER SEDATIVES SHOULD NOT BE USED. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

FOR EMERGENCY MEDICAL RESPONSE AND HAZARD COMMUNICATION ONLY CALL HAZARD COMMUNICATIONS SERVICES AT (800) 228-5635 ext. 283

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS WARNING/AVISO

May be fatal if swallowed. Avoid breathing dust or spray mist. Do not take internally. Avoid contact with eyes, skin or clothing. Wash hands and face before eating. Wash thoroughly after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

CONTAINS 10 ONE-POUND WATER-SOLUBLE PACKETS
NET CONTENTS.....10.....POUNDS

GOWAN

EPA Reg. No. 10163-134
EPA Est. No.

Gowan Company
P O Box 5569
Yuma, AZ 85366

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands, except under the forest canopy and use on rice. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning of equipment or disposal of wastes.

BEE HAZARD

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds, if bees are visiting the treatment area. Contact your Agricultural Extension Service Office for further information on bee activity in your area.

AGRICULTURE USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agriculture workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Hat or suitable headgear

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warning are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and oral warnings include the following information: **WARNING---Area treated with carbaryl on (date of application). Do not enter without appropriate protective clothing until sprays have dried.** (Use re-entry interval required by state if more restrictive.) In cases of accidental exposure, follow instructions under STATEMENT OF PRACTICAL TREATMENT section.

CHEMIGATION STATEMENT

This product may be applied through center pivot irrigation systems to **alfalfa, corn and potatoes, only**. Follow directions in the CROP USE DIRECTIONS section. Do not apply through any chemigation system other than that specified on this label.

APPLICATIONS RECOMMENDATIONS

Gowan Carbaryl 80 WSB is a dry powder insecticide for dispersion in water and application through hydraulic sprayers, mist blowers, low gallonage ground equipment and aircraft. Directions for water dilution are given under the CROP USE DIRECTIONS.

DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM THE AREA TREATED.

Tolerances established by the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when carbaryl is used in accordance with label directions. Forage or feed crops may be grazed or harvested for use as feed for dairy or meat animals without resulting residues in meat or milk. **READ THIS LABEL. USE IN STRICT ACCORDANCE WITH DIRECTIONS AND CAUTIONS.**

DO NOT PLANT rotational food and feed crops not registered for carbaryl in carbaryl-treated soil. Do not use reclaimed irrigation water from crops treated with carbaryl on upland crops from which carbaryl residue tolerances are not established.

The directions on this label are based on tests and field experience relating to (a) effectiveness; (b) possible injury to plant growth; and (c) residues in food, feed and milk. Follow directions for most effective and safest use.

Consult State Agricultural Experiment Stations or State Agricultural Extensions Service for additional information as the timing will vary with local conditions.

Spray Preparation

ASSURE A UNIFORM SUSPENSION: AGITATE THE MIXTURE BEFORE APPLICATION AND CONTINUOUSLY DURING APPLICATION. DO NOT STORE THE SPRAY MIXTURE. Flush entire system with clear water before use. Do not flush with kerosene or diesel oil. For concentrate application, screens in the application system should be 50 mesh or larger and nozzles should be fan type, number 3 or larger. Cone nozzles may be used if applying at least 3 gallons of finished spray per acre. Be sure that mix tank and application system is clean and free of residues, scale or other contamination. Fill mix tank 1/2 to 3/4 full with water and start agitation. Add the wettable powder. When powder is uniformly suspended, add the remaining water. (Appropriate ranges of water rates are given in the DILUTION DIRECTIONS.) Flush tank and lines with clean water after spraying. Prepare only as much spray mixture as can be applied on the day of mixing. Maintain continuous agitation during mixing and application to assure a uniform suspension. Do not store spray mixture for prolonged periods.

DIRECTIONS FOR USE OF WATER-SOLUBLE BAGS

Packets containing Gowan Carbaryl 80 WSB are water-soluble. Do not allow packet to become wet prior to adding to the spray tank. Do not handle with wet hands. Reseal outer bag to protect remaining packets. To prepare the spray mixture, drop the required number of unopened packets into spray tank while filling with water to the desired level. Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the packets should be completely dissolved within 5 minutes of the time they were added to the water. Use the specified amount of Gowan Carbaryl 80 WSB in the amount of water necessary to give complete coverage of foliage. Determine the total amount of wettable powder to be added to the spray tank based on the rates given in the CROP USE DIRECTIONS. For each one (1) pound of product to be added to the spray tank, use one (1) pound packet. Do not attempt to break open the water-soluble packets. Where dosages are expressed as fractions of pounds, prepare tank load to the lower of the nearest whole pound.

Compatibility Cautions

- 1) Physical compatibility of carbaryl with other pesticides is not fully known. Incompatible mixtures curdle or precipitate and may injure crops or reduce control. Compatibility can be effected by water in your locality, various surfactants and other formulation ingredients. Do not apply if physically incompatible.
- 2) Do not mix this product with nutrients, adjuvants or other pesticides unless your experience indicates that the mixture is safe and effective. In particular, combinations with more than one other pesticide should be evaluated before application. If uncertain, test a small amount of the mixture on a representative portion of the crop.
- 3) Unstable under highly alkaline conditions. Not effective if used with alkaline materials such as Bordeaux, lime-sulfur and casein lime spreaders.

Plant Response Precautions

- 1) To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.
- 2) Do not use on Boston ivy, Virginia creeper and maidenhair fern as injury will result. During early season, it may also injure Virginia and sand pines.
- 3) Observe label instructions for apple thinning and on combinations with certain herbicides on rice and soybeans.

Insect Control

Apply when insects or their damage appear. To maintain control, repeat at 7 to 14 day intervals or as necessary unless different directions are given in the CROP USE DIRECTIONS. Where dosage range is indicated, use the lower rate on young plants and early instars and higher rate on mature plants, advanced instars and adults. This product does not control spider mites.

Dilution Directions

The rate required for thorough, uniform coverage varies with the plant growth at time of application. Except as specified for certain uses, the following rates are therefore intended to cover a broad range of conditions. Where small volumes of spray are required, one (1) Tablespoon of product per gallon of water corresponds to a label rate of one (1) pound per acre or one (1) pound per 100 gallons.

Dilute Application

Field and Row Crops: Apply specified rate in 20 to 75 gallons of water per acre.

Fruit and Nuts: For dilute sprays apply the specified dosage per 100 gallons of water. Typical spray gallonages range from 200 to 300 gallons per acre.

Concentrate Application

Field and Row Crops: Use at least 3 gallons of finished spray per acre.

Fruits and Nuts: For concentrate sprays increase the concentration of product in the spray mixture to apply an amount per acre equivalent to that in a dilute spray. Use 30 to 100 gallons of finished spray per acre.

Aerial Application

Field and Row Crops: Use at least one (1) gallon of finished spray per acre.

Fruits and Nuts: Use at least 10 to 25 gallons of finished spray per acre. Apply an amount of product per acre that is equivalent to that in a dilute spray.

CROP USE DIRECTIONS

Agricultural use rates are given in pounds per acre, except as specified for tobacco, dip treatments and apple thinning. See DILUTION DIRECTIONS for appropriate water rates per acre. Also observe spray preparation and compatibility precautions elsewhere on this label. Application can be made by ground or aircraft.

PREHARVEST INTERVALS

The required days to wait between the last application and harvest are given in () after each crop name.

TREES AND VINES

ALMONDS (28): Fruit tree Leafroller, Navel orangeworm, Peach twig borer, San Jose scale: Use 1 1/4 lbs. per 100 gals. Apply in popcorn and petal fall stages and thereafter as needed. Time application to coincide with moth flight peaks. Make late season application at hullsplit for navel orangeworm.

APPLE THINNING (1): Apply dilute spray 10-25 days after full bloom. Consider factors including variety, tree age, previous crop, pruning, bloom and degree of set and exercise caution to avoid excessive thinning and consequent yield reduction. Contact your County Extension Service or other experts for advice on the use of this product. For easily thinned varieties, including Cortland, Grimes, Jonathan, McIntosh, Orleans, Rome Beauty, Puritan, Red Delicious, Winesap and Yellow Newton: Use 1/3 - 2/3 lbs. per 100 gals. For difficult to thin varieties, including Baldwin, Ben Davis, Duchess, Early McIntosh, Golden Delicious, Lady Apple, Northern Spy, Rhode Island Greening, Steele Red, Turley, Wealthy, Yellow Transparent and York Imperial: Use 2/3 - 1 1/4 lbs. per 100 gals.

APPLES (3), PEARS (3): Apple aphid, Apple maggot, Apple mealybug, Apple rust mite, Apple sucker, Bagworms, California pearslug, Codling moth, Eastern tent caterpillar, European apple sawfly, Eyespotted bud moth, Fruitree leafroller, Green fruitworm, Japanese beetle, Lesser appleworm, Lygus bugs, Orange tortrix, Pear leaf blister mite, Pear psylla, Pear rust mite, Periodical cicada, Plum curculio, Redbanded leafroller, Scale insects, Tarnished plant bug, Tentiform leafminers, White apple leafhopper, Woolly apple aphid: Use 1-1 1/4 lbs. per 100 gals. For psylla control, apply when eggs hatch or young nymphs are present. For scale control, apply when crawlers are present. DO NOT apply until at least 30 days after bloom to avoid undesired thinning.

APRICOTS (3): See PEACHES.

BLACKBERRIES (7), RASPBERRIES (7), DEWBERRIES, INCLUDING BOYSENBERRIES AND LOGANBERRIES (7): European raspberry aphid, Japanese beetle, Leafhoppers, Leafrollers, Omnivorous leafroller, Raspberry sawfly, Rose Chafer, Snowy rose tree cricket: Use 1 1/4 - 2 1/2 lbs. per acre.

BLUEBERRIES (7): Blueberry maggot, Cherry fruitworm, Cranberry fruitworm, European fruit Lecanium, Japanese beetle: Use 1 7/8 lbs. per acre.

BOYSENBERRIES (7): See BLACKBERRIES.

CHERRIES: See PEACHES.

CHESTNUTS (14): Chestnut weevil: Use 2 1/2-3 3/4 lbs. per 100 gals. DO NOT USE IN CALIFORNIA. Last application should be made prior to shuck split.

CITRUS (5) (INCLUDING GRAPEFRUITS, LEMONS, LIMES, ORANGES, TANGELOS, TANGERINES, CITRUS CITRON, KUMQUATS AND HYBRIDS): Avocado leafroller, California orangedog, Citrus cutworm, Citrus root weevil, Fruitree leafroller, Fullers Rose Beetle, Orange tortrix, Western Tussock moth, West Indian sugarcane borer: Use 1-1 1/4 lbs. per 100 gals. May be mixed with petroleum oils commonly used on citrus. Use according to manufacturer's directions. Do not apply more than 25 lbs. per acre per application. Do not apply less than 10 gals. of dilute spray mixture per mature tree.

CRANBERRIES (7): Cutworms, Cranberry fireworms, Cranberry fruitworms, Japanese beetle, Leafhoppers, Rose Chafer: Use 1 7/8 - 3 3/5 lbs. per acre. MAY KILL FISH AND CRABS. OBSERVE PRECAUTIONS TO PROTECT AQUATIC ORGANISMS. DO NOT APPLY WHERE THESE ARE IMPORTANT RESOURCES.

DEWBERRIES (7): See BLACKBERRIES.

FILBERTS (14): Filbert aphid, Filbert leafroller, Filbertworm: Use 1 1/4 lbs. per 100 gals. Apply for leafroller when eggs are hatching.

GRAPES (7): Cutworms, Eight spotted forester, European fruit lecanium, Grape berry moth, grape leafroller, Grape leafhopper, Japanese beetle, June beetles, Orange tortrix, Omnivorous leafroller, Redbanded Leafroller, Saltmarsh caterpillar, Western grape leaf skeletonizer, Western yellow-striped armyworm: Use 1 1/4 - 2 1/2 lbs. per acre.

LOGANBERRIES: See BLACKBERRIES.

NECTARINES: See PEACHES.

OLIVES (14): Olive scale: Use 1 – 1¼ lbs. per 100 gals. Do not exceed 2 applications per year. Do not apply more than 18 ¾ lbs. of this product per acre per application.

PEACHES (3), APRICOTS (3), NECTARINES (3): Apple pendentis, Codling moth, Cucumber beetles, European earwig, Fruittree leafroller, Japanese beetle, June beetles, Lesser peachtree borer, Olive scale, Orange tortrix, Oriental fruit moth, Peach twig borer, Periodical cicada, Plum curculio, Redbanded leafroller, Scale insects, Tarnished plant bug, Tussock moth, Variegated leafroller: Use 1¼ lbs. per 100 gals. For best scale control, apply when crawlers are present. Do not apply more than 7½ lbs. per acre per application to apricots.

PEARS (3): See APPLES.

PECANS (14): Black margined aphid, Fall webworm, Hickory shuckworm, Lesser webworm, Pecan leaf phylloxera, Pecans Nut casebearer, Pecan spittlebug, Pecan stem phylloxera, Pecan weevil, Twig girdler, Walnut Caterpillar: Use 1¼ - 3 lbs. per 100 gals. Do not apply more than 9 lbs. per acre per application.

PISTACHIOS (14): Navel orangeworm: Use 2/3 – 2½ lbs. per 100 gal. Apply 150-300 gals. of spray per acre.

PLUMS: See PRUNES.

PRUNES (1), PLUMS (1), CHERRIES (1): Black cherry aphid, Brown soft scale, Cherry fruitworm, Cherry maggot, Codling moth, Eastern tent caterpillar, European earwig, Eyespotted bud moth, Forbes scale, Fruittree leafroller, Green fruitworm, Japanese beetle, Lesser peach tree borer, Mealy plum aphid, Orange tortrix, Peach twig borer, Plum curculio, Prune leafhopper, Redbanded leafroller, Rose chafer, Scale insects, Tussock moth, Variegated leafroller: Use 1 - 1 ¼ lbs. per 100 gals. Do not apply more than 7½ lbs. per acre per season.

RASPBERRIES (7): See BLACKBERRIES.

STRAWBERRIES (7): Flea beetle, Meadow spittlebug, Omnivorous leaftier, Strawberry leafroller, Strawberry weevil: Use 1½ - 2½ lbs. per acre. Do not use on Early Dawn and Sunrise varieties on the Delmarva Peninsula, as injury may occur.

WALNUTS (14): Calico scale, Codling moth, European fruit lecanium, Filbertworm, Fruittree leafroller, Frosted scale: Use 2/3 lbs. per 100 gals. Apply 1000 gals. spray per acre for mature trees. For European earwig control, use 2 ½ lbs. per 100 gals. and spray tree trunks to point of runoff.

FIELD AND ROW CROPS

ALL FIELD AND ROW CROPS (See specific crop for preharvest interval):

GRASSHOPPERS: Use 2/3 – 1¼ lbs. for nymphs on small plants or sparse vegetation on wasteland, ditch banks or borders. Use 1¼ - 1 7/8 lbs. for adult grasshoppers or applications to dense vegetation.

ALFALFA (7), CLOVER (7): Blister beetles, Mexican bean beetles: 2/3 – 1¼ lbs. Alfalfa caterpillar, Bean leaf beetle, Cucumber beetles, Green cloverworm, Japanese beetle, Leafhoppers, Three-cornered alfalfa hopper, Thrips, Velvetbean caterpillar: Use 1¼ lbs. Alfalfa weevil larvae, Armyworm, Cloverhead weevil, Corn earworm, Cutworms, Egyptian alfalfa weevil larvae, Essex skipper, European alfalfa beetle, Fall armyworm, Lygus bugs, Webworms, Yellowstriped armyworm: Use 1¼ - 1 7/8 lbs. Use higher rates east of the Rocky Mountains. For alfalfa weevil larvae, if pretreatment damage is extensive, cut alfalfa and treat the stubble. **TEXAS ONLY:** Cloverhead weevil: Use 2 lbs. Application can be made to alfalfa through center pivot irrigation system. See directions at end of FIELD AND ROW CROPS section.

ASPARAGUS (1): Asparagus beetle: Use 1¼ - 2½ lbs. Asparagus beetle, Apache cicada: Use 2½ - 5 lbs. in post harvest application only. Do not treat more than once every 3 days for post harvest application.

BEANS (INCLUDING DRY BEANS, GREEN BEANS, LIMA BEANS, NAVY BEANS, SNAP BEANS, SOUTHERN PEAS, CROWDER PEAS, COWPEAS, BLACKEYED PEAS [succulent (3), forage (14), dried and hay (21,)]: Blister beetle, Mexican bean beetle: Use 2/3 – 1¼ lbs. Alfalfa caterpillar, Bean leaf beetle, Cucumber beetles, Flea beetles, Green cloverworm, Japanese beetle, Leafhoppers, Three-cornered alfalfa hopper, Thrips, Velvetbean caterpillar, Western bean cutworm: Use 1¼ lbs. Armyworm, Corn Earworm, Cutworms, European corn borer, Fall armyworm, Stink bugs, Tarnished plant bug, Webworms: Use 1¼ - 1 7/8 lbs. Cowpea curculio: Use 2½ lbs. **CALIFORNIA ONLY—**Corn earworm, Lima bean pod borer, Lygus bugs, Stink bugs: Use 2½ lbs.

BROCCOLI (3): See CABBAGE.

BRUSSELS SPROUTS (3): See CABBAGE.

CABBAGE (3), BROCCOLI (3), BRUSSELS SPROUTS (3), CAULIFLOWER (3), KOHLRABI (3): Flea beetles, Harlequin bug: Use 2/3 – 1¼ lbs. Armyworm, Imported cabbageworm, Corn earworm: Use 1¼ - 2½ lbs.

CARROTS (7) CELERY (14) PARSLEY (14), PARSNIPS (3): Flea beetles, leafhoppers: Use 2/3 – 1¼ lbs. Aster leafhopper, Lygus bugs: Use 1¼ - 2½ lbs. Treat on a 5-7 day schedule. **DO NOT USE ON CELERY IN CALIFORNIA.**

CAULIFLOWER: See CABBAGE.

CELERY: See CARROTS.

CHINESE CABBAGE (14-15), COLLARDS (14), HANOVER SALAD (14-15), HORSERADISH (7), KALE (14), MUSTARD GREENS (13-14), RADISHES (7), RUTABAGAS (7), TURNIPS (roots (7), tops (14-15)): Flea beetle, Harlequin bug, Leafhoppers: Use 2/3 – 1¼ lbs. Aster leafhopper: Use 1¼ - 1 7/8 lbs. Armyworm, Imported cabbageworm, Corn earworm, Tarnished plant bug: Use 1¼ - 2½ lbs.

CLOVER (7): See ALFALFA.

COLLARDS(14): See CHINESE CABBAGE.

CORN (FIELD AND POP)(forage and silage (14), grain and fodder (48); (SWEET) (ears (2), forage (14), fodder (48): Corn earworm, Corn rootworm adults, European corn borer, Fall armyworm, Flea beetles, Japanese beetle, Leafhoppers, Southwestern corn borer: Use 1¼ - 2½ lbs. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Western bean cutworm: Use 2½ lbs. Treat between 90% and 100% tassel emergence. Cutworms: Use 2½ - 8 lbs. Spray in a 12 inch band over row. **OBSERVE BEE CAUTION ON FRONT OF LABEL. APPLICATION DURING POLLEN SHED WILL REDUCE BEE POPULATION.** Application can be made to corn through center pivot irrigation system. See directions at end of FIELD AND ROW CROPS section.

CUCUMBER (3), MELONS (3), PUMPKINS (3), SQUASH (3): Cucumber beetles*, Flea beetles*, Leafhoppers*, Melonworm, Pickleworm, Squash bugs*: Use 2/3 – 1¼ lbs. (*Use higher rates for these pests.)

DANDELION (14-15), ENDIVE (ESCAROLE) (14-15), LETTUCE (head and leaf (14-15), SALSIFY (7): Flea beetles, Harlequin bug, Leafhoppers: Use 2/3 – 1¼ lbs. Aster leafhopper, Lygus bugs, Spittlebugs: Use 1¼ - 1 7/8 lbs. Armyworm, Corn earworm, Fall armyworm, Imported cabbageworm, Stink bugs, Tarnished plant bug: Use 1¼ - 2½ lbs.

EGGPLANT (3): See POTATO.

ENDIVE(14-15): See DANDELION.

FORAGE GRASS (3-aerial application, 14-ground application) and PASTURE (3-aerial application, 14-ground application): Armyworm, Black grass bugs, Chinch bugs, Essex skipper, Range caterpillars, Thrips: Use 1¼ - 1 7/8 lbs. White grubs: Use 1 7/8 – 2½ lbs.

GARDEN BEET (roots (7), tops (14-15)) SPINACH (14-15), SWISS CHARD (14-15): Flea beetles, Harlequin bug, Leafhoppers: Use 2/3 – 1¼ lbs. Aster leafhopper: Use 1¼ to 1 7/8 lbs. Armyworm, Corn earworm, Tarnished plant bug, Stinkbug: Use 1¼ - 2½ lbs.

HANOVER SALAD (14-15): See CHINESE CABBAGE.

HORSERADISH (7): See CHINESE CABBAGE.

KALE(14): See CHINESE CABBAGE.

KOHLRABI (3): See CABBAGE.

LETTUCE (14-15): See DANDELION.

MELONS (3): See CUCUMBER.

MUSTARD GREENS (13-14): See CHINESE CABBAGE.

OKRA (3): Corn earworm, Stink bugs: Use 1¼ - 2½ lbs.

PARSLEY (14-15): See CARROTS.

PARSNIPS (7): See CARROTS.

PASTURE: See FORAGE.

PEANUTS (14): Blister beetles, Mexican bean beetle: Use 2/3 – 1¼ lbs. Alfalfa caterpillar, Bean leaf beetle, Cucumber beetles, Green cloverworm, Japanese beetles, Leafhoppers, Three-cornered alfalfa hopper, Thrips, Velvetbean caterpillar: Use 1¼ lbs. Armyworm, Corn earworm, Stink bugs, Webworms: Use 1¼ - 1 7/8 lbs. Whitefringed beetle adults, Cutworms: Use 2½ lbs. To avoid possible injury to tender foliage, do not apply when foliage is wet or rain or excessive humidity is expected during the next two days.

PEAS (succulent (3), dried (21), forage (14), hay (21)): Alfalfa caterpillar*, Armyworm, Cutworms*, Colorado potato beetle*, Leafhoppers, Pea leaf weevil*:
Use 1¼ - 1 7/8 lbs. (*Use higher rate for these pests.)

PEPPERS: See POTATOES.

POTATO (7), TOMATO (3), EGGPLANT (3), PEPPER (3): Flea beetles, Leafhoppers: Use 2/3 – 1¼ lbs. Colorado potato beetle, Cutworms, European corn borer, Fall armyworm, Tomato fruitworm, Tomato hornworm, Tarnished plant bug, Stink bugs: Use 1½ - 2½ lbs. Use higher rate for cutworms. Application can be made to potatoes through center pivot irrigation system. See directions at end of FIELD AND ROW CROPS section.

PUMPKINS (3): See CUCUMBER.

RADISHES(7): See CHINESE CABBAGE.

RICE (14): MISSISSIPPI DELTA AREA AND TEXAS: Armyworm, Chinch bugs, Stink bugs: Use 1¼ - 1 7/8 lbs. CALIFORNIA ONLY: Tadpole shrimp: Use 2½ lbs applied to water when pest first appears. Armyworm, Leafhopper: Use 2½ lbs. SEE ENVIRONMENTAL HAZARDS SECTION ON MAIN LABEL FOR PRECAUTIONS REGARDING AQUATIC APPLICATION. DO NOT APPLY PROPANIL HERBICIDES WITHIN 15 DAYS BEFORE OR AFTER APPLICATION, OR CROP INJURY WILL RESULT.

RUTABAGAS (7): See CHINESE CABBAGE.

SALSIFY (7): See DANDELION.

SORGHUM [grain or straw (21), forage (7)] (INCLUDING MILO, GRAIN SORGHUM, AND HYBRIDS): Armyworm, Corn earworm, Stink bugs, Webworms: Use 1¼ - 2½ lbs. Sorghum midge: Use 1 7/8 lbs. Apply when about ¼ of heads have emerged and are in bloom. Repeat application 3-5 days later if adults are still active. A third application may be necessary. Cutworms: Use 2½ lbs. Direct spray into forming heads.

SOYBEANS [forage (14), hay (21)]: Bean leaf beetles, Blister beetles, Cucumber beetles, Grape colaspis, Green cloverworm, Mexican bean beetle, Velvetbean caterpillar: Use 2/3 – 1¼ lbs. Corn earworm: Use 2/3 – 1 7/8 lbs. Vary rate depending on degree of infestation. Alfalfa caterpillar, Japanese beetle, Leafhoppers, Three-cornered alfalfa hopper: Use 1¼ lbs. Armyworm, Cutworms, Fall Armyworm, Stink bugs, Webworms: Use 1¼ - 1 7/8 lbs. Thistle caterpillar, Saltmarsh caterpillar, Yellow striped armyworm: Use 2½ - 3 lbs.
DO NOT APPLY CARBARYL COMBINED WITH 2,4-D HERBICIDES TO SOYBEANS OR CROP INJURY WILL RESULT.

SPINACH: See GARDEN BEETS.

SQUASH: See CUCUMBERS.

SWISS CHARD: See GARDEN BEETS.

SUGAR BEETS [roots, forage (28)]: Armyworm, Cutworms, Fleabeetles, Leafhoppers, Webworms: Use 1¼ - 1 7/8 lbs. Use higher rate for cutworms.

SUNFLOWER (60): Armyworms, Cutworms: Use 1 7/8 – 2½ lbs. Stem weevil, Sunflower beetle: Use 1¼ - 2½ lbs. Do not graze treated crops.

SWEET POTATOES (0): Corn earworm, Cucumber beetle, Flea beetles, Sweet potato hornworm, Sweet potato weevil, Tortoise beetles: Use 1¼- 2½ lbs. Use lower rate for young plants.

TOBACCO (3): IN PLANT BEDS—Tobacco flea beetle: Use 1 lb. per 40 gals. Apply 6 gals. per 100 square yards. Green June beetle grubs: Use 1 lb. per 130 gals. Apply to areas uprooted by sprinkling 50-100 gals. per square yards. May be applied before or after seeding. **IN FIELDS**—Budworms, Fall armyworm, Flea beetles, Japanese beetles, June beetles, Hornworms: Use 1¼-2½ lbs. per acre. Avoid excessive application to prevent crop injury. Also, avoid excessive application to bud of plant.

TOMATO (3): See POTATO.

TURNIPS [roots (7), tops (14-15)]: See CHINESE CABBAGE.

WHEAT [forage (7), grain or straw (21)]: EXCEPT CALIFORNIA: Cereal leaf beetles: Use 1¼ lbs. Armyworm: use 1¼ - 1 7/8 lbs.

DIRECTIONS FOR APPLICATION THROUGH CENTER-PIVOT IRRIGATION SYSTEMS

Application of this product through sprinkler irrigation systems is permitted for **potatoes, corn and alfalfa**. Directions, including rates and pre-harvest intervals, are given in the crop use directions above. Specific directions which must be followed for application by chemigation system are given here.

General Precautions

Apply this product only through center pivot irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness or illegal pesticide residues on the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

DO NOT connect an irrigation system (including greenhouse system) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Precautions required for Sprinkler Chemigation

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond area intended for treatment.

Additional Precautions required for Chemigation systems connected to public water systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of 15 individuals at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction.

There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Further directions for Chemigation systems

Fill mix tank half to three-quarters full of water. Begin agitation. Add wettable powder and maintain agitation. Once powder is suspended, add water to fill tank. Maintain agitation until application is complete. The desired rate per acre may be suspended in less than five (5) gallons of water, then injected into the irrigation line.

Uniform application is essential. Equipment must be calibrated to ensure steady flow. The sprinkler system must follow a pattern which will not over expose or under expose any part of the crop.

Begin carbaryl application in order to complete application when irrigation is completed. Do not irrigate for at least two days to avoid washing off the product.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Store in a dry place, in original container.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300
For other product information, contact Gowan Company or see Material Safety Data Sheet

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

EPA Text Pending: Gowan Carbaryl 80 WSB (to EPA 02-08-00)