

5417936 345
17

Ms. Bethany G. Hulcy
Gowan Company
P. O. Box 5569
Yuma, AZ 85366-5569

JUN 25 1992

Dear Ms. Hulcy:

Subject: Gowan Dimethoate 4
EPA Registration No. 10163-160
Your Application for Pesticide Amendment
Dated April 23, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. On the label, under the ENVIRONMENTAL HAZARDS section, change the first sentence of the second paragraph to read:

"This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds."


2. Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

The Confidential Statement of Formula (CSF), dated April 23, 1992, submitted with this application is now considered the Basic Formulation for the subject product and all previous CSFs are considered obsolete. Please note the first listed inert (solvent) is a List 2 Inert of Toxicological Concern under FIFRA and as such we strongly encourage you to substitute for this component in its entirety in the formulation for the subject product.

Sincerely yours,



Robert A. Forrest
Product Manager (14)
Insecticide-Rodenticide Branch
Registration Division (H7504C)

enclosure

MADDEN Disk #3 6/24/92

Front panel - P91
D.2

GOWAN DIMETHOATE 4

Active Ingredient: Dimethoate- (O,O-dimethyl S-[(methylcarbamoyl) methyl] phosphorodithioate).....	43.5%
Inert Ingredients.....	56.5%
TOTAL.....	100.0%

* Contains 4 pounds of Dimethoate per gallon

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, do not induce vomiting. Call a Poison Control Center immediately. Get medical attention.

IF INHALED, remove victim to fresh air. Apply artificial respiration if necessary.

IF ON SKIN, wash with plenty of soap and water. Get medical attention.

IF IN EYES, flush eyes with plenty of water. Seek medical attention.

NOTE TO PHYSICIAN: Atropine is antidotal if symptoms of cholinesterase inhibition are present. 2-PAM: PROTOPAM chloride may be effective as an adjunct to atropine. Use according to label directions.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning - Harmful or fatal if swallowed. Vapor harmful- concentrated material. Avoid breathing vapor or spray mist. Use only with adequate ventilation. Avoid contact with eyes as this product causes eye irritation. Avoid contact with skin, and clothing. Wash thoroughly after handling and before eating, drinking, toileting or using tobacco. Keep container lid closed. Do not contaminate food or feed products.

~~REMOVED~~ **BEST AVAILABLE COPY**
with COMMENTS
In EPA Letter Dated:

JUN 25 1992
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
10163-160

REQUIRED CLOTHING AND EQUIPMENT FOR APPLICATION

All applicators, including flaggers and all personnel associated with the mixing, loading, and transferring operations must wear the protective clothing and equipment enumerated below. Pilots are exempt from this requirement. The protective clothing and equipment to be worn are as follows:

1. Impermeable gloves (for example- rubber or plastic covered reinforced gloves).
2. Boots or boot covers.
3. Long-sleeved shirt and long pants.
4. Wide-brimmed hat.
5. Respirators must be worn by flaggers and mixers/loaders.

AERIAL APPLICATIONS: AUTOMATIC FLAGGING DEVICES SHOULD BE USED WHENEVER FEASIBLE. IF HUMAN FLAGGERS ARE EMPLOYED, THEY MUST WEAR THE PROTECTIVE CLOTHING AND RESPIRATOR SPECIFIED ON THIS LABEL.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

This pesticide is toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Agricultural Extension Service.

PHYSICAL AND CHEMICAL HAZARDS

Combustible liquid and vapor. Do not use, pour, spill, or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not enter treated areas without appropriate protective clothing until sprays have dried. Protective clothing means, at least, a hat or other suitable head covering, a long-sleeved shirt and long-legged pants or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if

BEST AVAILABLE COPY

there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written or oral warnings must include the following information: Warning - Area treated with Dimethoate on (date of application). Do not enter treated area without appropriate protective clothing until sprays have dried (or other reentry interval if your State has a more restrictive interval for this product). For citrus do not enter treated area without appropriate protective clothing for four (4) days. In case of accidental exposure, follow instructions under "STATEMENT OF PRACTICAL TREATMENT" section

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container. DO NOT STORE BELOW 45° F

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: **Plastic**- triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Metal**- triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

See attached booklet for Directions For Use.

GOWAN COMPANY

P.O. BOX 5569

YUMA, AZ 85366

EPA Reg. No. 10163-160

EPA Est. No. 10163-AZ-1

BEST AVAILABLE COPY

GOWAN DIMETHOATE 4

DIRECTIONS FOR USE

GOWAN COMPANY

P.O. BOX 5569
YUMA, AZ 85366-5569

EPA Reg. No. 10163-160
EPA Est. No. 10163-AZ-1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR APPLICATION

This product is intended for use in conventional hydraulic sprayers, ground applicators, or aerial sprayers. Do not apply when weather conditions favor drift of spray from treated areas. Repeat applications as necessary unless otherwise specified. Consult your State Experiment Station or State Extension Service for proper timing of application.

COMPATIBILITY

DIMETHOATE 4 is formulated for application in water suspensions. It is compatible with most insecticides, fungicides, and miticides. It should not be used, however, with alkaline materials such as Bordeaux mixture and lime.

Gowan Dimethoate 4 can be mixed with carbaryl, diazinon, pyrethroids, methyl azinphos, malathion, parathion, dicofol, dodine, captan, zineb, and thiram. Because uniform dispersability and sprayability may be influenced by pesticide combinations used, it is recommended that compatibility be determined before adding pesticides to the spray tank. In a pint or quart jar, mix products and water proportionate to the intended tank mix. If there is any separation, we recommend that the combination not be used. The addition of a nonionic general purpose spreader-activator will usually eliminate any incompatibility noted.

DILUTION DIRECTIONS

The rate required for thorough, uniform coverage varies with plant growth at the time of application. The following rates are therefore intended to cover a broad range of conditions.

Dilute Applications: Field and Vegetable Crops: Apply specified rate in 20 to 75 gallons of water per acre. Fruits and Nuts: Apply specified rate in 100 to 800 gallons of water per acre. For citrus use up to 2000 gallons of water per acre.

BEST AVAILABLE COPY

Concentrate Applications: Field and Vegetable Crops: Apply specified rate in not less than 5 gallons of water per acre. Fruits and Nuts: Apply specified rate in 20 to 100 gallons of water per acre. These applications require special concentrate equipment.

Air Applications: Field and Vegetable Crops: Apply specified rate in a minimum of 1 gallon of water per acre. Fruits and Nuts: Apply specified rate in a minimum of 5 gallons of water per acre.

CHEMIGATION STATEMENT

Refer to supplemental labeling entitled "APPLICATION THROUGH IRRIGATION SYSTEMS, CHEMIGATION" for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

CROP USE DIRECTIONS

CAUTION: DO NOT USE ON SEED ONIONS, SEED CARROTS, OR SEED BERMUDA GRASS.

The interval between last application and harvest is given in () following the crop name.

FIELD CROPS

Alfalfa: (10) Aphids, Grasshoppers, Leafhoppers, Plant Bugs including Lygus, reduction of Alfalfa Weevil larvae; Use 1/2 to 1 pint per acre. Do not apply to alfalfa in the bloom period. Do not graze livestock in the treated crops, hay, threshings, or stubble within 10 days of application. Effective only on cutting to which applied. Limit use to one application per cutting.

Alfalfa (seed crop only): Aphids, Grasshoppers, Leafhoppers, Lygus Bugs, reduction of Alfalfa Weevil larvae; Use 1/2 to 1 pint per acre. Do not apply if the crop or weeds in the treatment area are in bloom. Do not feed livestock in treated crops, hay, threshings, or stubble within 10 days of application.

Field Corn: (14) Banks Grass Mites (excluding Trans-Pecos area of Texas), Aphids, Bean Beetle, Corn Rootworm adult; Use 2/3 to 1 pint per acre. Grasshoppers; Use 1 pint per acre. Do not apply to corn during the pollen-shed period if bees are actively foraging in the treatment area. Apply as necessary but make no more than three applications per year. Do not feed or graze within 14 days of last application.

Cotton(Arizona and California): (14) Leafhoppers, Fleahoppers, Plant Bugs including Lygus; Use 1/2 to 1 pint per acre. Repeat applications should not be made at intervals closer than 14 days. Make only 2 applications per season at the higher rate. Do not feed treated forage or graze livestock on treated fields.

BEST AVAILABLE COPY

Cotton (Except Arizona and California): (14 if water is used for dilution, 40 if once-refined vegetable oil is used for dilution) Aphids, Mites, Thrips, Fleahoppers; Use 1/4 to 1/2 pint per acre. Plant Bugs including Lygus; Use 1/2 pint per acre. If water is used for dilution, repeat applications should not be made at intervals closer than 14 days. Do not feed treated forage or graze livestock on treated fields. If once-refined vegetable oil is used for dilution, repeat applications should not be made at intervals closer than 40 days. Make only 2 applications per season at the higher rate. Apply at least 1 quart of finished spray per acre. Do not feed treated forage or graze livestock on treated fields.

Safflower (Arizona and California): (14) Aphids, Leafhoppers, Plant Bugs including Lygus, Thrips; Use 1/2 to 1 pint per acre. Repeat applications should not be made at intervals closer than 14 days. Make only 2 applications per season at the higher rate.

Sorghum (milo): Aphids; Use 1/2 to 1 pint per acre. Banks Grass Mites (excluding Trans-Pecos area of Texas), Grasshoppers, Spider Mites; Use 1 pint per acre. Sorghum Midge; Use 1/4 to 1/2 pint per acre. Do not apply after heading. Do not feed or graze within 28 days of last application. Apply as needed but not more than 3 applications per season.

Soybeans: (21) Mexican Bean Beetle, Spider Mites, Bean Leaf Beetle, Leafhoppers, Three-cornered Alfalfa Hopper *. Grasshoppers; Use 1 pint per acre. Do not feed or graze within 5 days of last application.

*Not registered in California

Wheat: (35) Aphids; Use 1/2 to 3/4 pint per acre. Brown Wheat Mite; Use 1/3 to 1/2 pint per acre. Grasshoppers; Use 3/4 pint per acre. Do not apply within 14 days of grazing immature plant. Do not make more than 2 applications per season.

CAUTION: DO NOT USE ON SEED ONIONS, SEED CARROTS, OR SEED BERMUDA GRASS.

VEGETABLE CROPS

Beans (green, lima, snap, dry): (0) Aphids, Grasshoppers, Leafhoppers, Leaf Miners, Lygus Bugs, Mites, Bean Leaf Beetle, Mexican Bean Beetle; Use 1/2 to 1 pint per acre. Do not apply if the crop or weeds in treatment area are in bloom. Do not feed treated vines.

Broccoli, Cauliflower: (7) Aphids; Use 1/2 to 1 pint per acre.

Cabbage: (3) Aphids; Use 1/2 to 1 pint per acre.

Celery (Florida): (7) Leaf Miners; Use 1 pint per acre

Head Lettuce: (7) Aphids, Leafhoppers, Leaf Miners; Use 1/2 pint per acre.

BEST AVAILABLE COPY

Leaf lettuce, Spinach, Collards, Kale, Turnip (greens and roots), Mustard Greens, Swiss Chard, Endive (Escarole): (14) Aphids, Leafhoppers, Leaf Miners; Use 1/2 pint per acre.

Melons (except Watermelons): (3) Aphids, Leafhoppers, Leaf Miners, Thrips; Use 1 pint per acre.

Watermelons: (3) Aphids, Leafhoppers, Leaf Miners; Use 1/2 to 1 pint per acre.

Lupine: (0) Aphids, Lygus Bugs; Use 1/2 to 1 pint per acre. Apply when Aphids first appear. Do not apply if crops or weeds in the treatment area are in bloom. Do not make more than 2 applications per crop season. Do not feed or graze forage or hay.

Peas, Lentils: (0) Aphids; Use 1/3 pint per acre. Do not feed or graze hay within 21 days after last application. Do not make more than one application per growing season. Do not apply if the crop or weeds in the treatment area are in bloom.

Peppers: (0) Aphids, Leaf Miners, Maggots; Use 1/2 to 2/3 pint per acre.

Potatoes: (0) Aphids, Grasshoppers, Leafhoppers, Leaf Miners; Use 1/2 to 1 pint per acre.

Tomatoes: (7) Aphids, Leafhoppers, Leaf Miners; Use 1/2 to 1 pint per acre.

FRUIT AND NUT CROPS

Apples: (28) Apple Maggot, Codling Moth*; Use 3 to 4 pints per acre (or Northeast and North Central states : Use 1 pint per 100 gallons based on 300 to 400 gallons of water per acre). Apply at petal-fall and every 10-14 days thereafter until control is achieved. Under heavy infestations, some sting injury may occur. Do not apply when trees or substantial number of weeds in the orchard are in bloom. Do not graze livestock in treated orchards.

* Midwest and Eastern states only.

Apples, Pears: (28) Aphids, Leafhoppers, Mites (except Rust Mites): Use 1 1/2 to 3 pints per acre (or Northeast and North Central states; Use 1/2 to 1 pint per 100 gallons based on 300 gallons of water per acre). Do not apply when trees or substantial number of weeds in the orchard are in bloom. Do not graze livestock in treated orchards.

Grapefruit, Lemons, Oranges, Tangerines: (see text below for harvest intervals) Aphids; Use 1 to 2 quarts per acre in 5 to 10 gallons of water. Apply as an outside coverage spray. The interval between last application and harvest is 15 days. Mites (except Rust Mites); Use 1/2 to 1 pint per 100 gallons of water with a maximum of 2 quarts per acre. The interval between last application and harvest is 15 days. Scales (except Black or Snow); Use 1 to 1 1/2 pints per 100 gallons or equivalent amount in concentrated spray. Apply as thorough coverage spray. The interval between last application and harvest is 45 days. Thrips; Use 1 to 2 quarts per acre in 5 to 10 gallons of water applied as a mist spray. The interval between last application and harvest is

15 days. Whiteflies; Use 3 to 4 pints per acre applied as a thorough distribution coverage spray. The interval between last application and harvest is 15 days. Do not apply during bloom period. Do not use on citrus seedlings. Make no more than 2 applications to mature fruit. Do not enter treated groves within 4 days of last application. Do not graze livestock in treated orchards.

Citrus (Arizona and California)(Non-bearing and nursery stock): Aphids, Thrips; Use 3 to 4 pints per acre as a foliar spray. Repeat applications as necessary. May be applied in the year grapefruit, lemon, orange, and tangerine trees begin to bear fruit. Do not enter treated groves within 4 days of last application without appropriate protective equipment. Do not graze livestock in treated groves. For soil drench (1-3 year old trees); Use 2 quarts per acre applied in the furrow or basin around the base of the tree. Apply when insect injury to new growth appears. Do not apply to trees that will bear fruit within one year. Do not graze livestock in treated groves.

Pecans: (21) Aphids, Mites, Leafhoppers; Use 2/3 pint per acre. Do not graze livestock in treated groves.

BEST AVAILABLE COPY

APPLICATION THROUGH IRRIGATION SYSTEMS—CHEMIGATION

Apply this product only through sprinkler, including center pivot, lateral move, and tow side (wheel) roll, traveler, big gun, solid set, or hand move, flood (basin), furrow, border or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Mix in clean supply tank the recommended amount of this product for acreage to be covered, and needed quantity of water.

This product should not be tank-mixed with other pesticides, surfactants or fertilizers unless prior use has shown the combination noninjurious under your conditions of use. Follow precautionary statements and directions for oil tank-mix products.

On all crops, use sufficient gaulonage of water to obtain thorough and uniform coverage, but not cause runoff or excessive leaching. This will vary depending on equipment, pest problem and stage of crop growth. Application of more or less than optimal quantity of water may result in decreased chemical performance, crop injury or illegal pesticide residues.

Enter this product into the irrigation water uniformly during the period of operation. Do not overlap application. Follow recommended label rates, application timing, and other directions and precautions for crop being treated.

Continuous mild agitation of pesticide mixture may be needed to assure a uniform application, particularly if the supply tank requires a number of hours to empty.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Note: Gowan Company does not encourage connecting chemigation systems to public water supplies. The following information is provided for users who have diligently considered all other application and water supply options before electing to make such a connection.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily or least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

SPRINKLER CHEMIGATION (FOLIAR SPRAY USES)

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

FLOOD (BASIN) FURROW AND BORDER CHEMIGATION (SOIL DRENCH USES)

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a dike structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

a. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

b. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

c. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

d. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

e. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

f. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

DRIP (TRICKLE) CHEMIGATION (SOIL DRENCH USES)

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pipe and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

BEST AVAILABLE COPY

Booklet ^{1104-L} - p. 7

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

ALL APPLICABLE RESTRICTIONS, PRECAUTIONS, AND DIRECTIONS ON THE EPA REGISTERED PRODUCT LABEL MUST BE FOLLOWED

4/92

BEST AVAILABLE COPY