

3/21/97

F3C

PM 14

10163-160

pg 1/10

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-31-98.



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input type="checkbox"/>	Other

OPP Identifier Number
246432

Application for Pesticide - Section I

1. Company/Product Number Gowan Company / 10163-160	2. EPA Product Manager Bill Jacobs	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Gowan Company / Gowan Dinosebure 4	PM# 14	
5. Name and Address of Applicant (Include ZIP Code) Gowan Company P.O. Box 5569 Yuma, AZ 85366-5569 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input checked="" type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification per PR Notice 95-2 to add advisory statements.

NOTIFICATION

MAR 21 1997

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify)	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Cindy Baker	Title Registration Specialist	Telephone No. (Include Area Code) 520-783-8844
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature Cindy Baker	3. Title Registration Specialist	
4. Typed Name Cindy Baker	5. Date 2/25/97	

2/10

GOWAN DIMETHOATE 4

Active Ingredient: Dimethoate	% By Wt.
(O,O-dimethyl S-[(methylcarbamoyl) methyl] phosphorodithioate).....	43.5%
Inert Ingredients.....	56.5%
	Total 100.0%

* Contains 4 pounds of Dimethoate per gallon

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

Organophosphate Insecticide

IF SWALLOWED, do not induce vomiting. Call a Poison Control Center immediately. Get medical attention.

IF INHALED, remove victim to fresh air. Apply artificial respiration if necessary.

IF ON SKIN, wash with plenty of soap and water. Get medical attention.

IF IN EYES, flush eyes with plenty of water. Seek medical attention.

NOTE TO PHYSICIAN: Atropine is antidotal if symptoms of cholinesterase inhibition are present. 2-PAM (PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning - Harmful or fatal if swallowed. Vapor harmful- concentrated material. Avoid breathing vapor or spray mist. Use only with adequate ventilation. Avoid contact with eyes as this product causes eye irritation. Avoid contact with skin and clothing. Keep container lid closed. Do not contaminate food or feed products.

NOTIFICATION
MAR 21 1997

ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and aquatic invertebrates. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Agricultural Extension Service.

NET CONTENTS _____ GALLONS

EPA Reg. No. 10163-160
EPA Est. No. 67545-AZ-1

Gowan Company
P.O. Box 5569
Yuma, AZ 85366

3/10

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers (other than mixers and loaders) must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils or Viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective Eyewear
- Chemical-resistant headgear for overhead exposure

Mixers and Loaders must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils or Viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear
- For exposures in enclosed areas - A respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G)
- For exposures outdoors - Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements: When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. AERIAL APPLICATIONS: AUTOMATIC FLAGGING DEVICES SHOULD BE USED WHENEVER FEASIBLE.

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

PHYSICAL AND CHEMICAL HAZARDS

Combustible liquid and vapor. Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

PHYTOTOXICITY STATEMENT

As is common with most emulsifiable concentrate formulations, adverse effects such as spotting or discoloration of the fruit or foliage can occur. Some conditions known to contribute to phytotoxicity include, but are not limited to: high temperatures, poor spray drying conditions, excessive spray deposit or runoff, certain spray mixtures, stage of crop development or tank mixes with other pesticides.

RESISTANCE MANAGEMENT

Based on historical use patterns in some area, certain pest species listed on this label may have developed resistance to Dimethoate 4. Consult your local agricultural advisor, State Cooperative Extension Service or regional Gowan Company representative for recommendations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils or Viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

DIRECTIONS FOR APPLICATION

This product is intended for use in conventional hydraulic sprayers, ground applicators, or aerial sprayers. Do not apply when weather conditions favor drift of spray from treated areas. Repeat applications as necessary unless otherwise specified. Consult your State Experiment Station or State Extension Service for proper timing of application.

COMPATIBILITY

DIMETHOATE 4 is formulated for application in water suspensions. It is compatible with most insecticides, fungicides, and miticides. It should not be used, however, with alkaline materials such as Bordeaux mixture and lime.

Gowan Dimethoate 4 can be mixed with carbaryl, diazinon, pyrethroids, methyl azinphos, malathion, parathion, dicofol, dodine, captan, zineb, and thiram. Because uniform dispersability and sprayability may be influenced by pesticide combinations used, it is recommended that compatibility be determined before adding pesticides to the spray tank. In a pint or quart jar, mix products and water proportionate to the intended tank mix. If there is any separation, we recommend that the combination not be used. The addition of a nonionic general purpose spreader-activator will usually eliminate any incompatibility noted.

DILUTION DIRECTIONS

The rate required for thorough, uniform coverage varies with plant growth at the time of application. The following rates are therefore intended to cover a broad range of conditions.

Dilute Applications: Field and Vegetable Crops: Apply specified rate in 20 to 75 gallons of water per acre. Fruits and Nuts: Apply specified rate in 100 to 800 gallons of water per acre. For citrus, use up to 2,000 gallons of water per acre.

Concentrate Applications: Field and Vegetable Crops: Apply specified rate in not less than 5 gallons of water per acre. Fruits and Nuts: Apply specified rate in 20 to 100 gallons of water per acre. These applications require special concentrate equipment.

Air Applications: Field and Vegetable Crops: Apply specified rate in a minimum of 1 gallon of water per acre. Fruits and Nuts: Apply specified rate in a minimum of 5 gallons of water per acre.

5/10

CHEMIGATION STATEMENT

Refer to the section "APPLICATION THROUGH IRRIGATION SYSTEMS , CHEMIGATION" for use directions for chemigation. Do not apply this product through any irrigation system unless the instructions for chemigation are followed.

CROP USE DIRECTIONS

The interval between last application and harvest is given in () following the crop name.

FIELD CROPS

Alfalfa: (10) Aphids, Grasshoppers, Leafhoppers, Plant Bugs including Lygus, reduction of Alfalfa Weevil larvae; Use 1/2 to 1 pint per acre. This pesticide is highly toxic to bees, do not apply if bees are visiting the areas to be treated when the crop or weeds are in bloom. Do not graze livestock in the treated crops, hay, threshings, or stubble within 10 days of application. Effective only on cutting to which applied. Limit use to one application per cutting.

Alfalfa (seed crop only): Aphids, Grasshoppers, Leafhoppers, Lygus Bugs, reduction of Alfalfa Weevil larvae; Use 1/2 to 1 pint per acre. This pesticide is highly toxic to bees, do not apply if bees are visiting the areas to be treated when the crop or weeds are in bloom. Do not graze livestock in treated crops, hay, threshings, or stubble within 10 days of application.

Field Corn: (14) Banks Grass Mites (excluding Trans-Pecos area of Texas), Aphids, Bean Beetle, Corn Rootworm adult; Use 2/3 to 1 pint per acre. Grasshoppers; Use 1 pint per acre. Do not apply to corn during the pollen-shed period. Apply as necessary but make no more than three applications per year. Do not feed or graze within 14 days of last application.

Cotton (Arizona and California): (14) Leafhoppers, Fleahoppers, Plant Bugs including Lygus; Use 1/2 to 1 pint per acre. Repeat applications should not be made at intervals closer than 14 days. Make only 2 applications per season at the higher rate. Do not feed treated forage or graze livestock on treated fields.

Cotton (Except Arizona and California): (14 if water is used for dilution, 40 if once-refined vegetable oil is used for dilution) Aphids, Mites, Thrips, Fleahoppers; Use 1/4 to 1/2 pint per acre. Plant Bugs including Lygus; Use 1/2 pint per acre. If water is used for dilution, repeat applications should not be made at intervals closer than 14 days. If once-refined vegetable oil is used for dilution, repeat applications should not be made at intervals closer than 40 days. Make only 2 applications per season at the higher rate. Apply at least 1 quart of finished spray per acre. Do not feed treated forage or graze livestock on treated fields.

Grass, Grown for seed (For use in Oregon only) (14) Aphids, Plant bugs, Winter grain mites: Use 1/2 to 2/3 pint per acre. Apply in a minimum of 2 gallons of water per acre. Do not graze or use seed or seed screenings for livestock feed or food purposes.

Safflower (Arizona and California): (14) Aphids, Leafhoppers, Plant Bugs including Lygus, Thrips; Use 1/2 to 1 pint per acre. Repeat applications should not be made at intervals closer than 14 days. Make only 2 applications per season at the higher rate.

Sorghum (milo): (28) Aphids; Use 1/2 to 1 pint per acre. Banks Grass Mites (excluding Trans-Pecos area of Texas), Grasshoppers, Spider Mites; Use 1 pint per acre. Sorghum Midge; Use 1/4 to 1/2 pint per acre. Do not apply after heading. Do not feed or graze within 28 days of last application. Apply as needed but not more than 3 applications per season.

Soybeans: (21) Mexican Bean Beetle, Spider Mites, Bean Leaf Beetle, Leafhoppers, Three-cornered Alfalfa Hopper *, Grasshoppers; Use 1 pint per acre. Do not feed or graze within 5 days of last application.

*Not registered in California

Wheat: (35) Aphids; Use 1/2 to 3/4 pint per acre. Brown Wheat Mite; Use 1/3 to 1/2 pint per acre. Grasshoppers; Use 3/4 pint per acre. Do not apply within 14 days of grazing immature plant. Do not make more than 2 applications per season.

CAUTION: DO NOT USE ON SEED ONIONS OR SEED CARROTS

VEGETABLE CROPS

Beans (green, lima, snap, dry): (0) Aphids, Grasshoppers, Leafhoppers, Leafminers, Lygus Bugs, Mites, Bean Leaf Beetle, Mexican Bean Beetle; Use 1/2 to 1 pint per acre. This pesticide is highly toxic to bees, do not apply if bees are visiting the areas to be treated when the crop or weeds are in bloom. Do not feed treated vines.

Broccoli, Cauliflower: (7) Aphids; Use 1/2 to 1 pint per acre.

Brussels Sprouts (For use in California Only): (10) Aphids. Use 1-2 pints per acre. Apply in a minimum of 100 gallons of water by ground equipment only. Apply when insects first appear and repeat as needed. Do not feed or graze livestock in treated fields. Do not apply by air. Do not exceed 6 applications per growing season.

Cabbage: (7) Aphids; Use 1/2 to 1 pint per acre.

Celery (Florida): (7) Leafminers; Use 1 pint per acre.

Head Lettuce: (7) Aphids, Leafhoppers, Leafminers; Use 1/2 pint per acre.

Leaf lettuce, Spinach, Collards, Kale, Turnip (greens and roots), Mustard Greens, Swiss Chard, Endive (Escarole): (14) Aphids, Leafhoppers, Leafminers; Use 1/2 pint per acre.

Melons (except Watermelons): (3) Aphids, Leafhoppers, Leafminers, Thrips; Use 1 pint per acre.

Watermelons: (3) Aphids, Leafhoppers, Leafminers; Use 1/2 to 1 pint per acre.

Lupine: (0) Aphids, Lygus Bugs; Use 1/2 to 1 pint per acre. Apply when Aphids first appear. This pesticide is highly toxic to bees, do not apply if bees are visiting the areas to be treated when the crop or weeds are in bloom. Do not make more than 2 applications per crop season. Do not feed or graze forage or hay.

Peas, Lentils: (0) Aphids; Use 1/3 pint per acre. Do not feed or graze hay within 21 days after last application. Do not make more than one application per growing season. This pesticide is highly toxic to bees, do not apply if bees are visiting the areas to be treated when the crop or weeds are in bloom.

Peppers: (0) Aphids, Leafminers, Maggots; Use 1/2 to 2/3 pint per acre.

Potatoes: (0) Aphids, Grasshoppers, Leafhoppers, Leafminers; Use 1/2 to 1 pint per acre.

Tomatoes: (7) Aphids, Leafhoppers, Leafminers; Use 1/2 to 1 pint per acre.

FRUIT AND NUT CROPS

Apples: (28) Apple Maggot, Codling Moth*; Use 3 to 4 pints per acre (or Northeast and North Central states ; Use 1 pint per 100 gallons based on 300 to 400 gallons of water per acre). Apply at petal fall and every 10-14 days thereafter until control is achieved. Under heavy infestations, some sting injury may occur. Do not apply when trees or substantial number of weeds in the orchard are in bloom. Do not graze livestock in treated orchards.

* Midwest and Eastern states only.

Apples, Pears: (28) Aphids, Leafhoppers, Mites (except Rust Mites); Use 1 1/2 to 3 pints per acre (or Northeast and North Central states; Use 1/2 to 1 pint per 100 gallons based on 300 gallons of water per acre). Do not apply when trees or substantial number of weeds in the orchard are in bloom. Do not graze livestock in treated orchards.

Grapefruit, Lemons, Oranges, Tangerines: (see text below for harvest intervals) Aphids; Use 1 to 2 quarts per acre in 5 to 10 gallons of water. Apply as an outside coverage spray. The interval between last application and harvest is 15 days. Mites (except Rust Mites); Use 1/2 to 1 pint per 100 gallons of water with a maximum of 2 quarts per acre. The interval between last application and harvest is 15 days. Scales (except Black or Snow); Use 1 to 1 1/2 pints per 100 gallons or equivalent amount in concentrated spray. Apply as thorough coverage spray. The interval between last application and harvest is 45 days. Thrips; Use 1 to 2 quarts per acre in 5 to 10 gallons of water applied as a mist spray. The interval between last application and harvest is 15 days. Whiteflies; Use 3 to 4 pints per acre applied as a thorough distribution coverage spray. The interval between last application and harvest is 15 days. Do not apply during bloom period. Do not use on citrus seedlings. Make no more than 2 applications to mature fruit. Do not graze livestock in treated orchards.

Citrus (Arizona and California)(Non-bearing and nursery stock): Aphids, Thrips; Use 3 to 4 pints per acre as a foliar spray. Repeat applications as necessary. May be applied in the year grapefruit, lemon, orange, and tangerine trees begin to bear fruit. Do not graze livestock in treated groves. For soil drench (1-3 year old trees); Use 2 quarts per acre applied in the furrow or basin around the base of the tree. Apply when insect injury to new growth appears. Do not apply to trees that will bear fruit within one year. Do not graze livestock in treated groves.

Pecans: (21) Aphids, Mites, Leafhoppers; Use 2/3 pint per acre. Do not graze livestock in treated groves.

SHADE AND ORNAMENTAL TREES AND PLANTS

ORNAMENTAL STATEMENT

Before treating a large number of ornamental plants with Dimethoate 4 alone or as a tank mixture with any other material, make a test application on a few plants and observe for 7-10 days prior to treating large areas to reduce the possibility of plant injury.

Azaleas (Outdoor): Lace bugs, Leafminers, Mites, Tea Scale, Whiteflies. Use 1 oz. Per 6 gallons of water as a foliar spray.

Camellia: Aphids, Camellia Scale, Tea Scale, Mites: Foliar Spray: Use 1 oz. Per 6 gallons of water. Apply 2 sprays, 6 weeks apart at the first year followed by annual applications soon after first growth begins in the spring. Soil Drench: Apply as a soil drench around the base of plants in early spring at the rate of 1 1/3 ozs. per gallon of water per plant up to 6 feet tall. Increase the rate proportionally for larger plants.

Carnations: Aphids, Thrips, Mites: **Foliar Spray:** Use 1 oz. Per 6 gallons of water. **Soil drench:** Apply as a soil drench at the rate of 2 2/3 ozs. per 500 sq. feet of bed or bench (6 2/3 quarts per acre) in sufficient water for even distribution. Water in thoroughly after application.

Cyperus: Bactra Moth Larvae: Use 1 oz per 6 gallons of water as a drenching spray.

Day Lilies: Aphids, Thrips: Use 2 ozs. per 6 gallons of water as a foliar spray.

Arborvitae: Aphids, Bagworms, Mites: Use 2 ozs. per 6 gallons of water as a foliar spray.

Birch: Aphids, Leafminers: Use 1/2 oz. Per 6 gallons of water as a foliar spray. For leafminers, apply when leaves are expanded (about mid-May) and repeat in early July.

Boxwood: Leafminers, Mealybugs, Mites: Use 1 oz. per 6 gallons of water as a foliar spray. For leafminers, apply in spring when leafminer flies first appear or in early summer to control larvae in infested leaves.

Cedar: Mites: Use 2 ozs. per 6 gallons of water as a foliar spray.

Euonymus: Aphids, Scale: Use 2 ozs. per 6 gallons of water as a foliar spray.

Gardenias: Tea Scale, Whiteflies: Use 1 oz. per 6 gallons of water as a foliar spray.

Gerberas: Thrips: Use 1 oz. per 6 gallons of water as a foliar spray.

Gladiolas: Aphids, Thrips: Use 1 oz. per 6 gallons of water as a foliar spray.

Iris: Aphids, Iris borer, Thrips: Use 2 ozs. per 6 gallons of water as a foliar spray. For borer control, spray when new leaves are 5-6 inches tall.

Poinsettas (outdoor): Mites, Whiteflies, Mealybugs, Aphids: Use 1 oz. per 6 gallons of water as a foliar spray.

Ficusnitida (Outdoor): Thrips: Use 1 oz. per 6 gallons of water as a foliar spray. Do not use on potted plants.

Holly (English and Amercian, not Buford variety). Leafminers, Mites, Soft Scale: Use 1 oz. per 6 gallons of water as a foliar spray. For leafminers, apply in spring when leafminer flies first appear or in early summer for control of larvae in the infested leaves.

Hemlock: Mites, Scales: Use 1 oz. per 6 gallons of water as a foliar spray.

Juniper: Aphids, Bagworms, Midges, Mites: Use 2 ozs. per 6 gallons of water as a foliar spray.

Oak: Golden Oak Scale: Use 2 ozs. per 6 gallons of water as a foliar spray.

Pine: Aphids, Bagworms, European pine shoot moth, Nantucket pine tip moth, Zimmerman pine moth: Use 2 ozs. per 6 gallons of water as a foliar spray.

Roses (Outdoor): Aphids, Leafhoppers, Mites, Thrips: Use 1 oz. per gallon of water as a foliar spray. For commercial fields, use 2/3 pint per acre in 5-10 gallons of water by air or 2/3 pint per acre in 100 gallons of water by ground.

Taxus: Fletcher Scale, Mealybugs, Mites: Use 2 ozs. per 6 gallons of water as a foliar spray.

APPLICATION THROUGH IRRIGATION SYSTEMS - CHEMIGATION

Apply this product only through sprinkler, including center pivot, lateral move, end tow side (wheel) roll, traveler, big gun, solid set, or hand move, flood (basin) furrow, border, or drip (including surface and subsurface) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Mix in clean supply tank the recommended amount of this product for acreage to be covered, and needed quantity of water.

This product should not be tank-mixed with other pesticides, surfactants or fertilizers unless prior use has shown the combination noninjurious under your conditions of use. Follow precautionary statements and directions for all tank-mix products.

On all crops, use sufficient gallonage of water to obtain thorough and uniform coverage, but not cause runoff or excessive leaching. This will vary depending on equipment, pest problem and stage of crop growth. Application of more or less than optimal quantity of water may result in decreased chemical performance, crop injury or illegal pesticide residues.

Meter this product into the irrigation water uniformly during the period of operation. Do not overlap application. Follow recommended label rates, application timing, and other directions and precautions

for crop being treated. Continuous mild agitation of pesticide mixture may be needed to assure a uniform application, particularly if the supply tank requires a number of hours to empty. Do not apply when wind speed favors drift beyond the area intended for treatment.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Note: *Gowan Company does not encourage connecting chemigation systems to public water supplies. The following information is provided for users who have diligently considered all other application and water supply options before electing to make such a connection.*

Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of a least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from a point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of a least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of material that are compatible with pesticides and capable of being fitted with a system interlock.

SPRINKLER CHEMIGATION (FOLIAR SPRAY USES)

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

FLOOD (BASIN) FURROW AND BORDER CHEMIGATION (SOIL DRENCH USES)

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- a. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

10710

- b. The pesticide injection pipeline must contain functional automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- c. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- d. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- e. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- f. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

**DRIP (INCLUDING SURFACE AND SUBSURFACE)
CHEMIGATION (SOIL DRENCH USES)**

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pipe and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container. DO NOT STORE BELOW 45° F

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

BEST AVAILABLE COPY