

PM13

10163-169

1/6

Please read instructions on reverse before completing form.

Form Approved: OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number
220253

Application for Pesticide - Section I

1. Company/Product Number 10163-169	2. EPA Product Manager Lakocca	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Imidan 70 WP Imidan 70 WSB	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) Gowan Company PO Box 5564 Yuma, AZ 85366 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Per PR Notice 95-2, Pest added to label

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Anne Stout	Title Reg. Specialist	Telephone No. (Include Area Code) 520-783-8844	
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature A. Stout	3. Title Reg. Specialist		
4. Typed Name Anne Stout	5. Date 2/7/96		



Imidan[®] 70-WP
Agricultural Insecticide

EPA Reg. No. 169

Imidan[®] 70-WSB
Agricultural Insecticide

EPA Reg. No. 10163-184

Wettable Powder...For Multi-Crop Pest Control

ACTIVE INGREDIENT: Phosmet	% By Wt.
N-(Mercaptomethyl) phthalimide, S-(O,O-dimethyl phosphorodithioate)	70%
INERT INGREDIENTS	30%
TOTAL	100%

SUPPLEMENTAL LABELING FOR USE ON CHRISTMAS TREE PLANTATIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.


This labeling must be in the possession of the user at the time of pesticide application.

CHRISTMAS TREE PLANTATIONS	Sawfly species, Scale species, European Pine Shoot Moth Eastern Pine Shoot Moth, Gypsy Moth, Nantucket Pine Tip Moth, Pitch Eating Weevil, Pales Weevil, adult Root Collar Weevil	1 1/3 lb. per acre	Apply in sufficient water to achieve thorough and complete coverage through aerial or ground application equipment. Apply when pest populations reach economic threshold levels as determined by the local Extension Service, Forest Service, or other monitoring system.
----------------------------------	--	-----------------------	--

®Imidan is a registered trademark of ZENECA Inc.

Gowan

GOWAN COMPANY • 1644 ENGLER AVENUE • P.O. BOX 5569 • YUMA, AZ 85366-5569
(520) 783-8844 • FAX (520) 343-9255



3/6

Imidan® 70-WP

Agricultural Insecticide

Wettable Powder...For Multi-Crop Pest Control

ACTIVE INGREDIENT:

Phosmet: N-(Mercaptomethyl)phthalimide, S-(O,O-dimethyl phosphorodithioate) 70%

INERT INGREDIENTS 30%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT Organophosphate Insecticide

Call a Poison Control Center or a physician immediately. If a known exposure occurs or is suspected, immediately start the recommended procedures below. Simultaneously contact a Poison Center, a physician, or the nearest hospital. Inform the person contacted of the type and extent of exposure, describe the victim's symptoms, and follow the advice given. NOTE: Be sure to advise the person contacted that the compound is a cholinesterase inhibitor.

NOTE TO MEDICAL PERSONNEL: Exposure may cause cholinesterase inhibition. Atropine by injection is antidotal. 2-PAM (Protopam Chloride) is also antidotal when administered early and in conjunction with atropine.

IF SWALLOWED: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Give fluids until vomitus is clear. If victim is unconscious or convulsing, do not induce vomiting or give anything by mouth.

IF INHALED: Remove to fresh air. If not breathing, clear victim's airway and start mouth-to-mouth artificial respiration. If breathing is difficult, give oxygen, preferably with a physician's advice.

IF ON SKIN: Flush all affected areas with plenty of water for several minutes. Seek medical attention if skin irritation occurs.

IF IN EYES: Hold eyelids apart and flush eyes with plenty of water. Get medical attention if irritation persists.

FOR 24 HOUR EMERGENCY ASSISTANCE, CALL CHEMTREC: (800) 424-9300.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING-AVISO

May be fatal if swallowed, inhaled, or absorbed through the skin. Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Chemical-resistant headgear for overhead exposure
- Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops while bees are actively visiting the treatment areas.

USE PRECAUTIONS

Read all precautions and directions before using. Apply this product only as specified on this label. Imidan 70-WP is compatible with most commonly used insecticides and fungicides, but is incompatible with alkaline materials such as spray lime, lime sulfur, and bordeaux mixtures. These materials will reduce the insecticidal activity of Imidan 70-WP. Insecticidal activity may also be reduced when the spray solution has a pH of 7 or higher. The pH of the spray solution must be corrected by the addition of a suitable buffering or acidifying agent for optimum insecticidal activity.

Gowan

EPA Reg. No. 10163-169
EPA Est. No. 67545-AZ-1

Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Imidan® 70-WP

Agricultural Insecticide

Wettable Powder...
For Multi-Crop Pest Control

Gowan

P.O. Box 5569
Yuma, AZ 85366-5569

EPA Reg. No. 10163-169
EPA Est. No. 67545-AZ-1

DIRECTIONS FOR USE

is a violation of Federal law to use this product in a manner inconsistent with its labeling.

or any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Chemical-resistant headgear for overhead exposure

DILUTION DIRECTIONS

The rate required for thorough, uniform coverage varies with plant growth at time of application. Except as specified for certain uses, the following rates are intended to cover a broad range of conditions.

Dilute Application

Field and Row Crops: Apply specified rate in 20 to 75 gallons of water per acre.

Tree and Vine Crops: Apply specified rate in 100 to 800 gallons of water per acre.

Concentrate Application

Field and Row Crops: Apply specified rate in not less than 5 gallons of water per acre.

Tree and Vine Crops: Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these applications.

Air Application

Field and Row Crops: Apply specified rate in a minimum of 3 gallons of water per acre.

Tree and Vine Crops: Apply specified rate in a minimum of 5 gallons of water per acre.

MIXING DIRECTIONS

Pour recommended amount of this material on surface of water in nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand.

USE LIMITATIONS

Do not exceed the maximum rate of Imidan 70-WP per acre or the time limitations specified for the individual crops.

CHEMIGATION STATEMENT

Apply this product only through one or more of the following types of systems: sprinkler (including center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move). Do not apply this product through any other type of irrigation system.

Refer to section entitled GENERAL CHEMIGATION INSTRUCTIONS for specific directions for use.

RECOMMENDATIONS FRUIT AND NUT CROPS

NOTE: Imidan 70-WP may be used during dormancy to control specified insects listed in each crop grouping which may overwinter on the tree and vine crops. Imidan 70-WP may be used in combination with spray oils; always follow spray oil manufacturer's label recommendations.

SPLIT APPLICATION SPRAY SCHEDULES: Applications to tree fruits and nuts may be made using a split application spray schedule as recommended below. Each schedule counts for one application. See crop for more specific application directions (if applicable).

Schedule #1: Apply one-half the approved rate per acre followed 5 to 7 days later with a second half of the approved rate. Both applications must be made to achieve control. Unless otherwise indicated, applications may be made by ground or air.

Schedule #2: Apply recommended rate per acre to every other row, followed with the recommended rate per acre 5 to 7 days later to alternate rows. Both applications must be made to achieve control.

ALMONDS: For control of Peach Twig Borer; Use 4 1/3 lbs. per acre. Limit use on bearing almonds to one foliar application per season. Do not apply within 30 days of harvest.

APPLES (EAST OF THE ROCKIES): For control of Apple Maggot, Codling Moth, Elm Spanworm, European Corn Borer, European Sawfly, Fruitree Leafroller, Green Fruitworm, Gypsy Moth, Japanese Beetle, Obliquebanded Leafroller*, Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Redbanded Leafroller, Redhumped Caterpillar, and San Jose Scale; Use 2 1/8 to 5 1/3 lbs. per acre (or 3/4 to 1 lb. per 100 gals. not to exceed 5 1/3 lbs. per acre) per application. For heavy insect infestations, use higher dosage rates. Repeat as necessary in accordance with insect infestations and local and State spray programs. May suppress European Red and Two-Spotted Mites when used in a seasonal program. Do not apply more than 30 lbs. per acre per crop season. Do not apply within 7 days of harvest.

*In some areas, these species may have developed resistance to organophosphate insecticides. Consult your local agricultural advisor or Cooperative Extension Service for recommendations.

TANK MIX (NORTHEAST ONLY): For control of the insects listed above for apples, plus Apple Aphid, Rosy Apple Aphid, Tarnished Plant Bug, Spotted Tentiform Leafminer, Tufted Apple Budmoth, and White Apple Leafhopper, a tank mix of Imidan 70-WP and methomyl (Lannate®) may be used as follows: Mix 1/3 to 2/3 lb. of Imidan 70-WP plus 1/2 to 1 pt. of methomyl 1.8L or 2 to 4 ozs. of methomyl 90% S.P. per 100 gals. of water. Apply as a full cover spray using up to 400 gals. per acre. Repeat as necessary in accordance with insect infestations and local or State spray programs. Do not use on early McIntosh or Wealthy varieties. Do not apply within 8 days of harvest. Do not graze/feed livestock under treated trees for 10 days after application. Do not apply more than 30 lbs. per acre of Imidan 70-WP per crop season.

®Lannate is a registered trademark of E.I. Du Pont de Nemours and Company, Inc.

APPLES (WEST OF THE ROCKIES): For control of Apple Maggot, Codling Moth, European Sawfly, Green Apple Aphid, Japanese Beetle, Orange Tortrix, Peach Twig Borer, Plum Curculio, Redbanded Leafroller, Redhumped Caterpillar, Rosy Apple Aphid, and San Jose Scale; Use 3 1/2 to 5 1/3 lbs. per acre per application. For heavy insect infestations, use higher dosage rates. Repeat as necessary in accordance with insect infestations and local and State spray programs. Do not apply more than 30 lbs. per acre per crop season. Do not apply within 7 days of harvest.

APRICOTS (EAST OF THE ROCKIES): For control of Apple Maggot, Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, and Redbanded Leafroller; Use 2 1/8 to 4 1/4 lbs. per acre (or 3/4 to 1 lb. per 100 gals. not to exceed 4 1/4 lbs. per acre) per application. For heavy insect infestations, use higher dosage rates. Repeat as necessary in accordance with insect infestations and local and State spray programs. May suppress European Red and Two-Spotted Mites when used in a seasonal program. Do not apply within 14 days of harvest.

APRICOTS (WEST OF THE ROCKIES): For control of Apple Maggot, Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, and Redbanded Leafroller; Use 4 1/4 lbs. per acre per application. Repeat as necessary in accordance with insect infestations and local and State spray programs. Do not apply within 14 days of harvest.

BLUEBERRIES (NORTHEASTERN U.S. ONLY): For control of Blueberry Maggot, Cherry Fruitworm, Cranberry Fruitworm, Japanese Beetle, Plum Curculio, Obliquebanded Leafroller, Redbanded Leafroller; Use 1 1/3 lbs. per acre as a foliar spraying a minimum of 2 gals. of water by aircraft. A second application may be made when indicated by insect infestations and local and State spray programs. May be applied up to 3 days before harvest.

CHERRIES, SOUR (TART): For control of Cherry Fruit Fly, Fruitree Leafroller, Japanese Beetle, Peach Twig Borer, Plum Curculio, and San Jose Scale; Use 2 1/8 to 2 1/2 lbs. per acre (or 3/4 lb. per 100 gals. not to exceed 2 1/2 lbs. per acre) per application. Repeat as necessary in accordance with insect infestations and local and State spray programs. May suppress European Red and Two-Spotted Mites when used in a seasonal program. Do not apply within 7 days of harvest.

GRAPES (EAST OF THE ROCKIES): For control European Rose Chafer, Fleabeetle, Grape Berry Moth, Grape Cane Borer, Grape Cane Girdler, Grape Leafhopper, Grape Mealybug, Japanese Beetle, and Redbanded Leafroller; Use 1 1/3 to 2 1/8 lbs. per acre. For grape berry moth, apply prebloom, postbloom, first and late cover sprays as needed. For grape leafhopper, apply when most nymphs hatch (generally coincides with grape berry moth). Use higher rates for control of Japanese beetle. Spray both sides of each row, and tops of vines to assure adequate coverage of fruit and foliage. Make applications as necessary to maintain control in accordance with insect infestations and local and State spray programs. Do not apply within 14 days of harvest.

GRAPES (WEST OF THE ROCKIES): For control of Grape Leafroller, Omnivorous Leafroller, and Western Grapeleaf Skeletonizer; Use 1 1/3 lbs. per acre. Spray both sides of each row and tips of vines to assure adequate coverage of fruit and foliage. Applications may be made at any time between egg hatch and pupation for leafroller, leafroller, and western grapeleaf skeletonizer. Do not apply within 7 days of harvest. (In California, the reentry interval is 5 days.)

NECTARINES (EAST OF THE ROCKIES): For control of Apple Maggot, Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, and Redbanded Leafroller; Use 2 1/8 to 4 1/4 lbs. per acre (or 3/4 to 1 lb. per 100 gals. not to exceed 4 1/4 lbs. per acre) per application. For heavy insect infestations, use higher dosage rates. Repeat as necessary in accordance with insect infestations and local and State spray programs. May suppress European Red and Two-Spotted Mites when used in a seasonal program. Do not apply within 14 days of harvest. Do not use Omite® in combination with Imidan 70-WP on late maturing nectarine varieties as fruit injury may result.

®Omite is a registered trademark of Uniroyal Chemical.

NECTARINES (WEST OF THE ROCKIES): For control of Apple Maggot, Omnivorous Leafroller, Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, and Redbanded Leafroller; Use 4 1/4 lbs. per acre per application. Repeat as necessary in accordance with insect infestations and local and State spray programs. Do not apply within 14 days of harvest. Do not use Omite® in combination with Imidan 70-WP on late maturing nectarine varieties as fruit injury may result. (In California, the reentry interval is 5 days.)

®Omite is a registered trademark of Uniroyal Chemical.

PEACHES (EAST OF THE ROCKIES): For control of Japanese Beetle, Oriental Fruit Moth, Peach Twig Borer, and Plum Curculio; Use 2 1/8 to 4 1/4 lbs. per acre (or 3/4 to 1 lb. per 100 gals. not to exceed 4 1/4 lbs. per acre) per application. For heavy insect infestations, use higher dosage rates. Repeat as necessary in accordance with insect infestations and local and State spray programs. May suppress European Red and Two-Spotted Mites when used in a seasonal program. Do not apply more than 17 lbs. per acre per crop season. Do not apply within 14 days of harvest.

PEACHES (WEST OF THE ROCKIES): For control of Japanese Beetle, Omnivorous Leafroller, Oriental Fruit Moth, Peach Twig Borer, and Plum Curculio; Use 4 1/4 lbs. per acre per application. Repeat as necessary in accordance with insect infestations and local and State spray programs. Do not apply more than 17 lbs. per acre per crop season. Do not apply within 14 days of harvest. (In California, the reentry interval is 5 days.)

PEARS (EAST OF THE ROCKIES): For control of Codling Moth, Elm Spanworm, Fruitree Leafroller, Gypsy Moth, Plum Curculio, and Redbanded Leafroller; Use 2 1/8 to 7 1/8 lbs. per acre (or 3/4 to 1 lb. per 100 gals. not to exceed 7 1/8 lbs. per acre) per application. For heavy insect infestations, use higher dosage rates. Repeat as necessary in accordance with insect infestations and local and State programs. May suppress European Red and Two-Spotted Mites when used in a seasonal program. Do not apply within 7 days of harvest.

PEARS (WEST OF THE ROCKIES): For control of Codling Moth, Elm Spanworm, Fruitree Leafroller, Gypsy Moth, Pear Psylla, Plum Curculio, and Redbanded Leafroller; Use 3 1/2 to 7 1/8 lbs. per acre per application. For heavy insect infestations, use higher dosage rates. Repeat as necessary in accordance with insect infestations and local and State programs. Do not apply within 7 days of harvest.

PECANS: For control of Pecan Nut Casebearer, Pecan Weevil®, and Hickory Shuckworm; Use 3 1/8 lbs. per acre (or 1 lb. per 100 gals. not to exceed 3 1/8 lbs. per acre) applied when infestation starts. Repeat as necessary in accordance with insect infestations and local or State programs. For Pecan Weevil, repeat applications should be made at 7 day intervals. Do not apply within 14 days of harvest. Do not graze or feed livestock on cover crops grown in treated pecan groves.

*For control of low to moderate populations of Pecan Weevils; adequate control may not be achieved when heavy populations are present.

PEACHES, PRUNES (EAST OF THE ROCKIES): For control of Apple Maggot, Codling Moth, Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Redbanded Leafroller, and Redhumped Caterpillar; Use 2 1/8 to 4 1/4 lbs. per acre (or 3/4 to 1 lb. per 100 gals. not to exceed 4 1/4 lbs. per acre) per application. For heavy insect infestations, use higher dosage rates. Repeat as necessary in accordance with insect infestations and local and State spray programs. May suppress European Red and Two-Spotted Mites when used in a seasonal program. Do not apply within 7 days of harvest.

PLUMS, PRUNES (WEST OF THE ROCKIES): For control of Apple Maggot, Codling Moth, Omnivorous Leafroller, Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Redbanded Leafroller, and Redhumped Caterpillar; Use 3 1/2 to 4 1/4 lbs. per acre per application. For heavy insect infestations, use higher dosage rates. Repeat as necessary in accordance with insect infestations and local and State spray programs. Do not apply within 7 days of harvest.

FIELD, FORAGE, AND VEGETABLE CROPS

ALFALFA (ARIZONA, CALIFORNIA AND NEVADA ONLY): For control of Common Alfalfa Weevil Larvae and Egyptian Alfalfa Weevil Larvae; Use 1 lb. of Imidan 70-WP per acre in a minimum of 10 gals. of water by ground equipment (20 gals. for dense stands) or 5 gals. of water by aircraft. Consult your local farm advisor regarding the proper timing of application. Larvae should be sprayed when they are actively feeding. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the CHEMIGATION section of this label. Do not apply more than once per cutting. Do not graze or cut for hay within 14 days of application. Do not apply to alfalfa in the bloom period. Do not use with latex or pinolene based adjuvants or any agricultural sticker or extender.

For Pea Aphids, apply the recommended rate of Imidan 70-WP in combination with 1/3 to 2/3 pt. Dimethoate 2.67E or 1/4 to 1/2 pt. Dimethoate 4E. Do not apply more than once per cutting. Do not graze or cut for hay within 14 days of application. Do not apply to alfalfa in the bloom period. Wear the protective clothing specified on the Dimethoate label when preparing and using this combination.

ALFALFA (EXCEPT ARIZONA, CALIFORNIA AND NEVADA): For control of Common and Egyptian Alfalfa Weevil Larvae and Adults, Potato Leafhopper, Meadow Spittlebug, and Alfalfa Blotch Leafminer; Use 1 1/3 lbs. per acre in a minimum of 10 gals. of water by ground equipment (20 gals. for dense stands) or a minimum of 2 gals. of water by

aircraft. Consult your local County Agent or Extension Service Representative regarding the proper timing of application. Larvae should be sprayed when they are actively feeding. Application for adult weevils should be timed to periods of visible activity. For alfalfa blotch leafminer and meadow spittlebug, application should be made when first signs of infestation are visible. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the CHEMIGATION section of this label. Do not apply more than once per cutting. Do not graze or cut for hay within 7 days of application. Do not apply to alfalfa during the bloom period.

*Recommended for Potato Leafhopper in the Northeast and North Central States only.

For Pea Aphids, apply the recommended rate of Imidan 70-WP in combination with 1/3 to 2/3 pt. Dimethoate 2.67E or 1/4 to 1/2 pt. Dimethoate 4E. Do not apply more than once per cutting. Do not graze or cut for hay within 10 days of application. Do not apply to alfalfa in the bloom period. Wear the protective clothing specified on the Dimethoate label when preparing and using this combination.

COTTON (EXCEPT SAN JOAQUIN VALLEY, CALIFORNIA): For control of overwintering generation of Boll Weevils; Apply 1/3 to 3/4 lb. per acre in a minimum of 5 gals. of water by ground equipment or apply 3/4 lb. per acre in a minimum of 3 gals. of water by aircraft. Make 2 applications. The first application should be made at the 1/3 grown square stage and the second 5 to 7 days later. Use the higher rate under heavy infestations. For control of first, second, or third generation Boll Weevils, apply 1 to 1 1/3 lbs. Imidan 70-WP in the same amount of water as specified above. Make applications at intervals of 3 to 7 days depending upon weevil population and weevil migration in fields. Use the higher rate for heavy infestations. Check infestations regularly. Do not exceed 14 lbs. of Imidan 70-WP per acre per season. Do not apply within 21 days of harvest. Do not graze or feed forage to livestock. Use on cotton prohibited in Copiah and Clairborne Counties, MS; Lauderdale and Madison Counties, AL; and Lawrence County, TN. Do not apply within one mile of any coastal or estuarine waters. Do not apply within 100 feet of aquatic habitats.

PEAS, FRESH AND DRY (PACIFIC NORTHWEST ONLY): For control of Pea Weevil and Pea Leaf Weevil; Use 1 to 1 1/3 lbs. per acre in a minimum of 5 gals. of water per acre by aircraft or 20 gals. of water by ground equipment. Apply between emergence and early pod formation when adult populations are present but before eggs are laid. Consult your local County Agent or Extension Service Representative regarding proper timing of application. Do not apply within 7 days of harvest. Do not graze or feed forage to livestock within 7 days of harvest. Do not cut treated fresh pea forage for hay within 10 days of application.

POTATOES: For control of Colorado Potato Beetle, Potato Flea Beetle, and Potato Leafhopper; Use 1 1/3 lbs. per acre in sufficient water to provide good coverage. Repeat applications as necessary throughout the growing season. For application by irrigation systems, apply specified dosage per acre. Follow all directions under the CHEMIGATION section of this label. Do not apply within 7 days of harvest. Use only on potatoes to be harvested by machine. Not recommended for use on potatoes in California.

FIELD MARGINS: For control of Grasshoppers in margins of cultivated fields and forage crop sites listed on this label, apply 2 1/2 to 2 3/4 lbs. Imidan 70-WP in 10 to 50 gals. of water per acre (20 to 50 gals. of water for dense stands) or in 5 to 10 gals. of water by aircraft. Do not graze livestock in treated area. Do not harvest for food or feed.

DECIDUOUS SHADE AND ORNAMENTAL TREES AND WOODY EVERGREENS

Imidan 70-WP is recommended for use by commercial applicators on deciduous shade and ornamental trees and woody evergreens in parks, residential and recreational areas, along thoroughfares and other localized areas where infestations of Gypsy Moth, Spring Cankerworm, Elm Spanworm, Birch Leafminer, Eastern Tent Caterpillar, Elm Leaf Beetle Larvae, Japanese Beetle, or Redhumped Caterpillar occur. When such insects or their damage occur, apply at a rate of 3/4 to 1 lb. per 100 gals. of water thoroughly wetting all parts of the affected plants to the point of runoff. For heavy insect infestations, use the higher dosage rate. Repeat application as necessary to maintain insect control.

• Recommended on Deciduous Shade and Ornamental Trees (such as Ash, Beech, Oak, Dogwood, Willow, Hickory, Locust, Liquidambar, Hawthorn, Birch, Elm, Maple) to control Gypsy Moth, Spring Cankerworm, Elm Spanworm, Eastern Tent Caterpillar, Elm Leaf Beetle Larvae, Japanese Beetle, and Redhumped Caterpillar.

• Recommended on Woody Evergreens (such as Arborvitae, Cedar, Fir, Hemlock, Juniper, Pine, Spruce, Yew) to control Gypsy Moth.

• Recommended on Birch Trees to control Birch Leafminer.

Choose a cool, calm period, preferably in early morning or evening. Do not apply if rain is expected or before leaf surfaces are dry.

CHRISTMAS TREE PLANTATIONS (EXCEPT CALIFORNIA): For control of Sawfly species, Scale species, European Pine Shoot Moth (*Rhyacionia buoliana*), Eastern Pine Shoot Moth, Nantucket Pine Tip Moth, Pitch Eating Weevil, Peas Weevil, adult Root Collar Weevil; Use 1 1/3 lbs. per acre in sufficient water to achieve thorough and complete coverage through ground or aerial application equipment. Apply when pest populations reach economic threshold levels as determined by the local Extension Service, Forest Service, or other monitoring systems.

PINE TREES (PACIFIC NORTHWEST ONLY): For control of European Pine Shoot Moth (*Rhyacionia buoliana*); Use 1 1/3 lbs. Imidan 70-WP per acre or for individual trees at a rate of 1 1/3 lbs. per 100 gals. of water. Spray trees for thorough coverage of foliage. Repeat at 2 week intervals throughout flight period of the moth.

PINE SEEDLINGS (WHITE, SLASH, AND LOBLOLLY): For control of Fates Weevil (*Hylobius pales*) and Pitch-eating Weevil (*Pachylobius picivorus*); Use Imidan 70-WP as a 4% top dip, dipping down to and including root collar only. Dip in bundles loose enough to allow solution to penetrate the bundles. Avoid coverage of roots. Soak tops in solution for 10 to 15 seconds to assure adequate coverage of all top growth. Drain and allow seedlings to dry before planting. The addition of an extender such as Nu-Film 17® may help retain the Imidan 70-WP on the foliage under high rainfall conditions. Some slight needle burn and first year growth reduction may occur on treatment of loblobly pine.

®Nu-Film 17 is a registered trademark of Miller Chemical & Fertilizer Corp.