

BOTRAN® 6% Dust Fungicide

For: Apricot, Garlic, Grape, Lettuce, Onion, Snap Bean, Tomato and Ornamentals

ACTIVE INGREDIENT:		% By Wt.
2, 6-dichloro-4-nitroaniline		6.0%
OTHER INGREDIENTS		94.0%
		TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
If Swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. -Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL TOLL FREE 1-888-478-0798	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes substantial but temporary eye injury. Harmful if swallowed. Avoid inhalation of dust. Do not get in eyes, on skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Goggles or face shield

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

Dealers Must Sell in Original Packages Only

NET CONTENTS _____ POUNDS



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Goggles or face shield


DIRECTIONS FOR USE ON EDIBLE CROPS

Applications of BOTRAN 6% DUST may be made with ground or aerial equipment, as appropriate. Do not use on germinating seeds or annual seedlings except as suggested. Extreme environmental conditions may have effects on fungicide treatments. The grower should watch to note that the treatment is working and in case of poor control, should take appropriate measures.

PREHARVEST INTERVAL

The required days between last application and harvest are given in () after each crop name.

CROP	DISEASE	LBS OF PRODUCT/ACRE	COMMENTS
Apricot (10)	Fruit decay <i>Monilinia</i> brown rot <i>Rhizopus</i> rot	50	Apply 10 days prior to harvest.
	<ul style="list-style-type: none"> • Do not apply more than 66 2/3 lb formulated product per acre per year. • Do not allow animals to graze treated areas. 		
Grapes (1)	Bunch rot, Storage rot (<i>Botrytis</i> rot)	30	Apply at cluster tightening and repeat at 2-week intervals.
	<ul style="list-style-type: none"> • Do not apply more than 66 2/3 lb formulated product per acre per year. • NOTE: Use limited to grapes grown west of the Rocky Mountains only. 		
Lettuce (leaf only) (14) (Greenhouse) (Field)	Blight <i>Botrytis</i> gray mold rot <i>Botrytis</i> cinerea	30	Dust 7 days following transplanting. Repeat application when plants are half mature. Do not apply to wilted plants or seedlings.
	Drop or watery soft rot <i>Sclerotinia</i> sclerotiorum <i>Botrytis</i> gray mold rot <i>Botrytis</i> cinerea	30	Apply first BOTRAN 6% DUST application at thinning. Repeat application 7 days later.
<ul style="list-style-type: none"> • Do not apply more than 66 2/3 lb formulated product per acre per year. 			
Onion, Garlic, and Shallot (see text)	<i>Botrytis</i> rot	41 2/3 - 66 2/3	Apply only at planting. Good coverage is essential for good disease control.
	<ul style="list-style-type: none"> • Do not apply more than 66 2/3 lb formulated product per acre per year. • Do not make more than one application per year. • Do not plant spinach as a follow-up crop in treated soil. 		
Snap Bean (2)	White mold <i>Sclerotinia</i> sclerotiorum (Sclerotiniöse)	50 (pole varieties) 40 (bush varieties)	Apply when disease is anticipated.
	<ul style="list-style-type: none"> • Do not apply more than 66 2/3 lb formulated product per acre per year. • Do not feed treated foliage to livestock. 		
Tomato (greenhouse) (10)	Stem canker <i>Botrytis</i> cinerea	30	Dust stem of plant from ground level up to a height of 18-24 inches in anticipation of disease. Repeat weekly or as necessary
	<ul style="list-style-type: none"> • Do not apply more than 66 2/3 lb formulated product per acre per year. 		



DIRECTIONS FOR USE ON ORNAMENTAL CROPS

CROP	DISEASE	LBS OF PRODUCT/ACRE	COMMENTS
Rose Hydrangea	Blight and gray mold <i>Botrytis cinerea</i>	30	Field: Apply BOTRAN 6% DUST prior to shipment or storage. Stored: Thoroughly dust storage area and all canes and dead foliage; Repeat application during long storage periods. • Do not apply more than 66 2/3 lb formulated product per acre per year.
Chrysanthemum Geranium Rose	Blight <i>Botrytis</i>	30	Stock cuttings or greenhouse plants: Begin applications when disease is anticipated or first appears. Dust foliage and flower at 7 to 14 day intervals. During prolonged damp, cool periods, apply every 5 to 7 days. • Do not apply more than 66 2/3 lb formulated product per acre per year.
Gladiolus	Dry Rot <i>Stromatinia gladioli</i>	50	Dust into furrow at planting time so that treated soil surrounds corms.
	Blight <i>Botrytis</i>	30	Dust thoroughly to insure coverage between spikes. Apply to bunched cut flower spikes after harvest and before shipment or storage. • Do not apply more than 66 2/3 lb formulated product per acre per year.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Store in original container and keep closed. Store in a cool, dry place.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.
CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable, return the unopened container for a full refund.

Our recommendations for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. All such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label and is reasonably fit for the intended purpose referred to on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE FULLEST EXTENT PERMITTED BY LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY AND GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Botran® is a registered trademark of Gowan Company LLC.

EPA Text Pending: Botran 6% Dust (resent to EPA 01-27-06)

ACCEPTED

MAR 10 2006

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10103-188

