10163-200 9.	-18-2006	1 1.000	1/8
	- .	Re >-* **	Print Form
Please read instructions on reverse before completing form. United States Environmental Protect Washington, DC 2	tion Agency	ed. OMB No. 2070-0060. Registration Amendment Other	Approval expires 2-28- DPP Identifier Number
Applicat	tion for Pesticide - Sectio		
1. Company/Product Number 10163-200	2. EPA Product Manage Jim Tompkins		osed Classification
4. Company/Product (Name) Prefar 4-E Selective Herbicide	PM# 25		LJ
5. Name and Address of Applicant (Include ZIP Code) Gowan Company P.O. Box 5569 Yuma, AZ 85366 Check if this is a new address	(b)(i), my product is s to:	r. In accordance with FII milar or identical in comp	osition and labeling
	Section - II		
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below. Explanation: Use additional page(s) if necessary. (For sect added "inhaled" to precautionary statement	Generation C "Me Too" Appl Other - Explain	cation. SEP	FICATION 1 8 2006
	Section - III		
Lebei Container		2. Type of Container Metal Plastic Glass Paper Other (Sper Other (Sper On Label On Label On Labeling accompanir	
Stèn	Section - IV	· · · · · · · · · · · · · · · · · · ·	
1. Contact Point <i>Complete items directly below for identificat</i>		cessary, to process this au	olication.)
Name Kyla S. Smith	Title Registration Specialist		o, (Include Area Code)
Certific I certify that the statements I have made on this form an I acknowledge that any knowlingity false or misleading s both under applicable law.	d all attachments thereto are true, ac	curate and complete.	Date Application Received (Stamped)
2. Signature	3. Title Registration Specialist		
4. Typed Name Kyla S. Smith	5. Date 8-22-06		
		The File Ocasis ford-deally	Vallaur Angliaget Co



9/ Du 18/4



Emulsifiable liquid for control of annual grasses and broadleaf weeds

ACTIVE INGREDIENT: Bensulide

	70 Gy 440.	
S-(O,O-diisopropyl phosphorodithioate) ester of N-(2-mercaptoethyl) benzenesulfonamide	46.0%	
OTHER INGREDIENTS	54.0%	
τοτ	L 100.0%	2

Contains 4 pounds of active ingredient per gallon Contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN CAUTION

	FIRST AID
	Organophosphate Pesticide
If swallowed	Immediately call a poison control center or doctor.
	Do not induce vomiting unless told to do so by the poison control center or doctor.
	Do not give ANY liquid to the person.
	Do not give anything by mouth to an unconscious person.
If inhaled	Move person to fresh air.
	If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
	Call a poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminated clothing.
	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
	Call a poison control center or doctor for treatment advice.
	HOT LINE NUMBER
Have the product containe 0798 for emergency medi	er or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-478- cal treatment information.
	NOTE TO PHYSICIAN
	phosphate herbicide. If symptoms of cholinesterase inhibition are present, atropine by injection is antidotal. 2-PAM is also inistered, but only in conjunction with atropine. Contains petroleum distillates – vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin or clothing. Causes moderate eye irritation.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- · Chemical resistant gloves such as barrier laminate or Viton > 14 mil.
- Shoes plus socks
- In addition, a NIOSH-approved dust-mist filtering respirator or with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter must be worn by: 1) mixers, loaders 2) persons participating in chemigation, and 3) commercial or for-hire applicators (due to risk from repeated exposures)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL

When handlers use closed systems or enclosed cabs, in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the Wos

NET CONTENTS _____ GALLONS

EPA Reg. No. 10163-200 EPA Est. No. 34704-MS-1



Gowan Company P O. Box 5569 Yunia, AZ 85366-5569

USER SAFETY RECOMMENDATIONS

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the tollet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This chemical is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from the treated area.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Do not apply directly to water, or to area where water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This product may impair reproduction in birds if used during the breeding season.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use, store, pour, or spill near heat or open flame. Keep container closed when not in use

DIRECTIONS FOR USE

GENERAL USE PRECAUTIONS AND RESTRICTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

- READ ALL DIRECTIONS BEFORE USING.
- MIX PREFAR 4-E WELL BEFORE DILUTING WITH WATER.
- Prefar 4-E should be used only for purposes listed on the label. Do not exceed the rates specified on the label. (DO NOT OVERDOSE.)
- USE ON MINERAL SOILS ONLY.
- Application of Prefar 4-E to soil that is cold and/or wet for an extended period of time due to rains or poorly drained soil during the germination period may result in reduction of crop stand or yields.
- Combining Prefar 4-Ewith fertilizers or other agricultural chemicals could result in incompatibility, reduced efficacy, or reduced crop safety. It is recommended that the user test the response of combination treatments with other products on a small scale prior to commercial scale use.
- Vary the rate of Prefar 4-E you apply depending on the soil texture. Apply the higher recommended label rates to heavier, fine textured soils. Apply lower rates to lighter, coarse textured soils with lower organic matter.
- This product may not be applied by aircraft.
- Do not allow this product to drift.

WEEDS CONTROLLED

ANNUAL GRASSES	
Crabgrass	(Digitaria spp.)
Fall Panicum	(Panicum dichotomiflorum)
Foxtail	(Setaria spp.)
Goosegrass	(Eleusine indica)
Jungle Rice	(Echinochloa colonum)
Sprangletop	(Leptochioa spp.)
Barnyardgrass	(Echinochioa crusgalli)
(Watergrass)	
ANNUAL BROADLEAF W	VEEDS
(AZ, NM, TX and CA only)
Lambsquarters	(Chenopodium album)
Purslane	(Portulaca oleracea)
Redroot Pigweed	(Amaranthus retroflexus)
(AZ and CA only)	
Burning Nettle	(Urtica urens)

(Chenopodium murale)

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate or viton > 14 mil.
- Shoes plus socks

Nettleleaf Goosefoot

GENERAL INFORMATION

Prefar 4-E is a selective preemergent herbicide that controls weeds by interfering with their normal germination and seedling development. Tests deemed reliable demonstrate that when Prefar is applied to labeled crops according to label directions under normal growing conditions, it is safe to the crop. Normal crop conditions include soil with good water drainage, acceptable soil fertility, and soil temperatures conducive to timely seed germination and plant emergence. Any deviation from normal growing conditions resulting from adverse environmental conditions or changes in production practices may alter a crop's response to Prefar.

The user should strictly comply with the soil prenaration and application instructions on this label. Questions concerning the use of the product should be directed to Gowan Company or local extension agents or university specialists.

SOIL PREPARATION: Prepare the soil according to good agricultural practice. Prefar 4-E dcea not control emerged weeds. All weed growth and crop stubble must be thoroughly worked into the soil prior to application.

APPLICATION INSTRUCTIONS: Spray equipment should be carefully calibrated before use and checked frequently during application to be sure nozzles are free from clogging and are delivering a uniform spray pattern. Apply uniformly the recommended rate of Prefar 4-E in 10 to 50 gallous of water per acre. See comments for specific crop use directions.

PREPLANT INCORPORATION: Apply to well-worked soil that is dry enough to permit thorough incorporation. Incorporate to a depth of 1-2 inches before planting. One-inch incorporation is optimum, while incorporating deeper than 2 inches will result in reduced weed control or crop injury.

PREEMERGENCE APPLICATION: Apply Prefar 4-E preemergence only on crops where the application is followed by immediate irrigation. Delaying irrigation more than 36 hours may result in poor weed control. For sprinkler irrigation wet the soil at least 2 to 4 inches deep. For furrow irrigation thoroughly wet the entire bed top (completely black out bed top).

ROTATIONAL CROP INSTRUCTIONS: Carrots, and other crops on this label (or crops from labeled crop groupings) may be replanted following applications of Prefar 4-E without restriction. All other crops should not be planted for 120 days and the soil must be tilled to a minimum depth of 4 inches prior to replanting. Prefar 4-E may be used on labeled crops grown through or under plastic.

CHEMIGATION USE INSTRUCTIONS

CHEMIGATION PROHIBITION – This product may only be applied by chemigation in Arizona and California.

Apply this product only through sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move; or drip (trickle), including surface and subsurface drip irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers, or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

For chemigation systems connected to public water systems:

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap

exists and that there is no blockage of the overflow of the reservoir tank.

In addition, all directions and requirements specified for Sprinkler and Drip (trickle) Chemigation must be followed. Sprinkler and Drip (Trickle) Chemigation:

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

For best results, application should be made at the beginning of the irrigation cycle, injecting over a 60 to 90 minute period after all lines and sprinklers have been cleared and are operating with correct pressure. Shut off injection equipment at the end of the application and continue to irrigate until soil is wetted to a depth of 2 to 4 inches.

CROP USE DIRECTIONS

All Prefar 4-E rates are given on a broadcast rate per acre basis. For banded applications reduce proportionately the amount of Prefar 4-E applied according to the area to be treated. Do not make applications to any crop not on this label unless the supplemental labeling for Prefar 4-E supplied by Gowan Company is followed.

Prefar 4-2 Supplied by Gowan Company is followed.		
CROP	BROADCAST	COMMENTS
	RATE	
	Quarts/Acre	
CUCURBIT VEGETABLE GROUP, including but not limited to:	5-6	Apply preplant or preemergence to flat-planted
Chayote, Chinese Waxgourd (Chinese Preserving Melon), Citron		or bedded crops. Follow application instructions
Melon, Cucumbers, Gherkin, Edible Gourd (Hechima, Chinese Okra),	[given on this label.
Muskmelons (including Muskmelon, True Cantaloupe, Cantaloupe,		
Casaba, Crenshaw Melon, Golden Pershaw Melon, Honeydew Melon,		
Honey Balls, Mango Melon, Persian Melon, Pineapple Melon, Santa		
Claus Melon, Snake Melon), Pumpkins, Summer Squash (Crookneck		
Squash, Straightneck Squash, Zucchini, Scallop Squash, Vegetable		
Marrow, Spaghetti Squash, Hyotan, Cucuzza, Balsam Apple, Balsam		
Pear, Bitter Melon, Chinese Cucumber), Winter Squash (Including		
Butternut Squash, Calabaza, Hubbard Squash, Acom Squash),		
Watermelons		
BRASSICA (COLE) LEAFY VEGETABLE GROUP, including but not	5-6	Apply preplant or preemergence. Follow
limited to: Broccoli, Broccolini, Broccoflower, Chinese Broccoli,		application instructions given on this label.
Broccoli Raab (Rapini), Brussels Sprouts, Cabbage, Chinese Cabbage	 Do not apply 	more than 6 lbs. a.i. per acre per season.
(Bok Choy, Napa), Chinese Mustard Cabbage (Gai Choy) Cauliflower,		
Cavalo Broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens,		
Mustard Spinach, Rape Greens		
FRUITING VEGETABLES	5-6	Apply prepiant or preemergence. Follow
Eggplant, Ground Cherry, Pepinos, Peppers (Bell Peppers, Chili		application instructions given on this label.
Peppers, Cooking Peppers, Pimentos, Sweet Peppers), Tomatillo		

CROP BROADCAST COMMENTS RATE Quarts/Acre LEAFY VEGETABLES (EXCEPT BRASSICA VEGETABLE) 5 - 6 Apply preplant or preemergence. Follow GROUP, including but not limited to: application instructions given on this label. Do Arugula (Roquette), Cardoon, Celery, Chinese Celery, Celtuce, Chervit, Chrysanthemum (Edible Leafed Garland), Corn Salad, Cress not use Prefar 4-E on spinach as severe phytotoxicity will occur. (Garden, Upland) Dandelion, Dock (Sorrel), Endive (Escarole), Do not apply more than 6 lbs. a.i. per acre per season Florence Fennel (Sweet Anise, Sweet Fennel, Finochio), Lettuce (Head Or Leaf), Orach, Parsley, Cilantro (Mexican Parsley, Chinese Parsley), Radicchio (Red Chicory), Swiss Chard CARROTS (Texas Only) 5 - 6 Apply preplant or preemergence. Follow application instructions given on this label. Do not feed treated carrots to livestock. Do not apply more than 5 lbs. a.i. per acre per season -DRY BULB VEGETABLES: Garlic, Bulb Onions, Shallots 5 - 6 Apply preplant or preemergence. Follow application instructions given on this label. Onions germinate and grow best when planted in fine-textured soil with good water drainage where the soil salinity value is less than 4 ECe* and soil temperature is between 55° and 75°. Use of Prefar on onions under less favorable soil and environmental conditions may reduce stand and vigor resulting in lower yield. To avoid injury, do not incorporate by mechanical methods in California or Arizona. Consult your local extension agent or university specialist for further information. (*ECe is the electrical conductivity extract.) Do not apply more than 6 lbs. a.i. per acre per season • DRY BULB VEGETABLES: For control of redroot pigweed and common 5 - 6 Garlic, Bulb Onions, Shallots (Idaho and Oregon only) lambsquarters, prepare a good, flat seed bed by standard cultural practices. Apply Prefar 4-E in the fall in 10 - 50 gals. of water in a 10 - 12 inch band and bed-up, throwing soil from the furrows over the sprayed band of Prefar 4-E. Allow to remain undisturbed until spring. Prior to planting in spring, drag-off the bed tops being careful not to drag off soil below the level of Prefar 4-E application. Plant in the center of the beds. Do not apply more than 6 lbs. a.i. per acre per season

PREFAR 4-E/ALANAP® TANK MIX DIRECTIONS

A tank mix combination of Prefar 4-E and ALANAP can be mixed (incorporated) into the soil before planting for weed control in cantaloupes, cucumbers, muskmelons and watermelons. This combination will control a broader spectrum of weeds than will either material used alone. **IMPORTANT:** Follow all general directions for use and use precautions that appear on the ALANAP label and elsewhere on this label. **Weeds controlled by PREFAR 4-E/ALANAP combinations:**

Annual Grasses: Bamyardgrass (watergrass), Crabgrass, Fall Panicum, Foxtail, Goosegrass, Sprangletop.

Annual Broadleaf Weeds: Field and hedge bindweed,

Carpetweed, Chickweed, Cocklebur, Galinsoga, Groundcherry, Lambsquarters, Mustard, Pigweed, Purslane, Ragweed, Shepherd's purse, Velvetleaf.

Soil Preparation: Prepare the soil according to good agricultural practice. All weed growth and crop stubble must be thoroughly worked into the soil.

Tank Mixing Directions: Combine Prefar 4-E and ALANAP in the spray tank in the recommended dosage proportions using 10 to 40 gallons of water per acre of area to be treated.

Spray equipment should be carefully calibrated before use and checked frequently during application to be sure nozzles are free from clogging and are delivering a uniform spray pattern.

CROP	BROADCAST RATE Quarts/Acre	COMMENTS
Cantaloupes, Cucumbers, Muskmelons, Watermelons	4-6 qts. Prefar 4-E plus 4-8 qts. ALANAP	Apply preplant in sufficient weter (19-40 gals.) to provide thereugh coverage. Incorporate lightly (½ to 1 inci) into the soil using suitable equipment just prior to planting. Use lower rates on light (sardy) hoils and higher rates on heavier (clay) soils. Reduce the rate proportionately for band applications. Incorporation is not necessary if the application is followed by immediate irrigation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Protect from temperatures below 42° F. At temperatures below 42° F product may crystallize. Dissolve crystals with quiet storage at warm temperatures (75° F). Redissolution rate: 40 - 48°F (40-60 days); 60° F (1-3 weeks); 70 - 80° F (1-3 days). Roll container and agitate contents to provide a uniform product prior to use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

SPILL OR LEAK: Small spills can be handled routinely. Use adequate ventilation and wear an air-supplied respirator to prevent inhalation. Wear suitable protective clothing to prevent

skin and eye contact. Use the following procedures:

1. Spread a suitable absorbent such as clay, sawdust or kitty litter on the spilled material.

- 2. Place sweepings in an appropriate chemical waste container.
- 3. Generously cover the contaminated areas with a common household detergent. Using a stiff brush and small amounts of water, work the detergent into the spill material forming a slurry. Do not splatter on yourself or bystanders and completely avoid skin or eye contact with this material. Brush the slurry into cracks and crevices and allow to stand for 2-3 minutes.
- 4. Spread absorbents on the slurried liquid and shovel the absorbed material into the container.

5. Repeat if necessary.

6. Flush the area with water while observing proper environmental considerations.

7. Seal container and dispose of in approved pesticide landfill.

Large spills should be handled according to a predetermined plan. For assistance in developing a plan, contact Gowan Company.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC[®](800) 424-9300.

For other product information, contact Gowan Company or

see Material Safety Data Sheet.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility including injury or damage, resulting from its misuse as such, or in combination with other materials.

This product is sold only for uses stated on its label. No expressed or implied license is granted to use or sale of this product under any patent in any country except as specified: Country: United States of America.

Prefar[®] is a registered trademark of ZENECA Inc. ALANAP® is a registered trademark of Uniroyal Chemical.

EPA Text Notification: Prefar 4E (8-22-06)

Supplemental Labeling

4

7/8



Emulsifiable liquid for control of annual grasses and broadleaf weeds

For Field Grown Flowers, Bulbs and Ornamentals*

CROP	RATE Quarts/Acre	COMMENTS
Herbaceous Plants: Including but not limited to: Alyssum, Aster, Bachelor's Button, Calendula, Candy-tuft, Coral Bell, Daisy,	5-9	Apply preplant or preemergence to flatplanted or bedded crops. Follow application instructions given on this label.
Marigold, Pansy, Primrose, Stock, Sweet Pea, Wallflower.	 Prior to treating new species or varieties, treat a few plants and observe results to reduce the possibility of crop injury. May not be used in residential areas. 	
Bulbs: Including but not limited to: Daffodil, Dahlia, Freesia, Gladiolus, Narcissus, Ranunculus, Tulip	 May not be used in resid 	





August 22, 2006

Docurnent Processing Desk (NOTIF) Jim Tompkins, PM Team 25 (703) 305-5697 U.S.E.P.A., Office of Pesticide Programs (7504P) One Potomac Yard 2777 South Crystal Drive Arlington, Virginia 22202

RE: Notification of minor changes per PR Notice 98-10 EPA Reg. No. 10163-200, Prefar 4-E

Dear Mr. Tompkins:

We are submitting notification to inform you of the following change that have been made to the Prefar 4-E label. In accordance with PR Notice 98-10, we are submitting the following to support this notification:

- EPA Form 8570-1 Application for Pesticide
- One (1) copy of the proposed label

The minor change is as follows:

"Inhaled" was added to the Precautionary Statement: "Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with eyes, skin or clothing. Causes moderate eye irritation." It was left off accidentally in our last submission.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Thank you very much for your help. If you have any questions or comments, please do not hesitate to contact me at (928) 819-1531.

Sincerely Kyla S. Smith

Regulatory Specialist

Enclosures

NOTIFICATION SEP 1 8 2006