

10163-220

01/07/2003

1/1

RESTRICTED USE PESTICIDE

Due to Reproductive Effects

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during mixing, loading, equipment repair and equipment cleaning. Certified applicators must ensure that all persons involved in these activities under their direct supervision are informed of the precautionary statements.

SUPPLEMENTAL LABELING FOR USE ON ONIONS

EPA Reg No. 10163-220

MSR™

**SPRAY CONCENTRATE
SYSTEMIC INSECTICIDE
FOR EFFECTIVE SYSTEMIC CROP PROTECTION**

ACCEPTED
JAN - 7 2003
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10163-220

ACTIVE INGREDIENT:

S-[2-(Ethylsulfinyl)ethyl] O,O-dimethyl phosphorothioate	25.0%
INERT INGREDIENTS	75.0%
	TOTAL 100.0%

Contains 2 lbs. S-[2-(Ethylsulfinyl)ethyl] O,O-dimethyl phosphorothioate per U.S. gallon
(This product contains aromatic petroleum distillates.)

KEEP OUT OF REACH OF CHILDREN

WARNING

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label must be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

CROP	PEST	PINTS*	COMMENTS
Spanish Onions (bulb) (West of the Mississippi River only)	Thrips (population suppression only)	1 ½ - 2	Apply specified dosage per acre in sufficient water for complete coverage but not less than 10 gallons per acre. For best results make 2 applications at a minimum of 10-14 day intervals. Use higher rate when pest pressure is high. For application by irrigation systems, apply specified dosage per acre. Follow all directions given under the CHEMIGATION section of this label.

RESTRICTIONS:

- Do not apply more than 1 lb ai/acre/season
- Preharvest Interval is 30 days
- Reentry Interval is 10 days
- Do not thin mature plants following an application of MSR™ Spray Concentrate
- Do not hand weed following an application of MSR™ Spray Concentrate
- Do not harvest by hand following applications of MSR™ Spray Concentrate

* This dosage of MSR™ Spray Concentrate is calculated for conventional sprayers. When lower volumes of spray are applied per acre, as with low pressure, low-volume, airplane or mist-type equipment, increase the concentration of MSR™ Spray Concentrate in the spray mixture in order to apply amount of MSR™ Spray Concentrate application per per acre to a full coverage spray but in no case less than one gallon of water per acre.

Registrant: Gowan Company
P.O. Box 5569
Yuma, AZ 85366