



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 7, 2022

Kyla S. Smith
Registration Specialist
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

Subject: Label – Voluntary Use Deletion for Removal of Geraniums and Hydrangeas
Product Name: Botran 5F Fungicide
EPA Registration Number: 10163-226
Application Date: 08/20/2019
Decision Number: 554870

Dear Ms. Smith:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

This amendment supports the notification of use termination submitted by Gowan Company to voluntarily amend the registration of Botran 5F Fungicide to terminate the following uses:

Table 1	
EPA Registration Name and Number	Use Deletion
Botran 5F Fungicide, EPA registration No. 10163-226	apricots, chrysanthemums, conifers, hydrangeas, geraniums, gladiolus, grapes, greenhouse cucumbers, greenhouse lettuce, greenhouse rhubarb, greenhouse tomato, nectarines, peaches, plums/prunes, potatoes, roses, and sweet cherries.

The Notice of Request to Voluntary Amend to Terminate Uses of the above registration and specified uses were published in the Federal Register on May 5, 2021, and the comment period ended on June 4, 2021. The Agency received seven comments on this notice; none of which would affect the Agency's action nor a written withdrawal of the use deletion from Gowan Company.

Pursuant to FIFRA section 6(f) (7 U.S.C. 136d(f)), the Agency hereby approves the requested amendment to terminate uses of Botran 5F Fungicide identified in Table 1 above.

Accordingly, the Agency orders that the product registration identified in Table 1 above is hereby amended to terminate the affected uses. Any distribution, sale, or use of existing stocks of the subject

product in a manner inconsistent with any of the provisions for disposition of existing stocks set forth below will be considered a violation of FIFRA.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. The review of this label entailed only the removal of the aforementioned uses. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. Existing stocks are those stocks of registered pesticide products which are currently in the United States and which were packaged, labeled, and released for shipment prior to the effective date of this action terminating uses. In accordance with the Order to Amend Pesticide Registrations to Terminate Uses (geraniums and hydrangeas) published in the Federal Register on October 29, 2021, you may distribute or sell this product under the previously approved labeling until April 29, 2023. Persons other than the registrant may continue to sell and/or use existing stocks of products bearing old labeling until such stocks are exhausted, provided that such use is consistent with the terms of the previously approved labeling on, or that accompanied, the product with the terminated use. After April 29, 2023, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Marcel Howard by phone at (703)305-6784, or via email at howard.marcel@epa.gov.



Nathan Mellor, Product Manager 21
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

Botran® 5F Fungicide

ACTIVE INGREDIENT: Dicloran	% BY. WT.
2, 6-dichloro-4-nitroaniline	46.7%
OTHER INGREDIENTS	53.3%
TOTAL	100.0%

Contains 5 lbs. dicloran per gallon


KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-478-0798 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist of vapor. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

NET CONTENTS _____ GALLONS



EPA Reg. No. 10163-226
EPA Est. No. XXXX

Gowan
The Go To Company

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ 86366-5569

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material.

All mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (except for pilots) made of barrier laminate, nitrile rubber ≥ 14mils, neoprene rubber ≥ 14mils, or viton ≥ 14mils

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instruction for washables exists, use detergent and hot water.

Keep and wash PPE separately from other laundry.

Early Entry Personal Protective Equipment

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls

USER SAFETY RECOMMENDATIONS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

ENVIRONMENTAL HAZARDS

This product is toxic to fish.

This product may contaminate water through runoff. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment, washwaters or rinsate.

This chemical can contaminate surface water through spray or dust applications. Under some conditions, it may also have a potential for runoff into surface water after application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

All other crops and use patterns: For all other crops and use patterns, do not enter or allow worker entry during the restricted entry interval of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, nitrile rubber ≥ 14mils, neoprene rubber ≥ 14mils, or viton ≥ 14mils
- made of any waterproof material
- Shoes plus socks

RESISTANCE MANAGEMENT

For Use as a Fungicide:

For resistance management, Botran 5F Fungicide contains a Group 14 fungicide. Any fungal/bacterial population may contain individuals naturally resistant to Botran 5F Fungicide and other Group 14 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of Botran 5F Fungicide or other Group 14 fungicides within a growing season sequence with different groups that control the same pathogens.

- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact a Gowan Company representative. You can also contact your pesticide distributor or university extension specialist to report resistance.

SPRAY DRIFT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product. Do not make any type of application into temperature inversions.

WIND SPEED: Do not apply at wind speeds greater than 10 mph at the application site.

DROPLET SIZE: Apply as a medium or coarser spray (ASAI standard 572).

RELEASE HEIGHT (GROUND APPLICATION): Apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

RELEASE HEIGHT (AERIAL APPLICATION): Do not release spray at a height greater than 10 feet above the ground or crop canopy.

Additional requirements for aerial applications:

- The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter
- Aerial applicators must consider flight speed and nozzle orientation in determining droplet size
- When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind

Additional requirements for airblast applications:

- Direct sprays into the canopy
- Turn off outward pointing nozzles at row ends and when spraying outer rows

CHEMIGATION STATEMENT

This product can be applied through the following types of chemigation systems: Sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move, or drip irrigation systems. Do not apply this product through any other type of irrigation system. Refer to section entitled CHEMIGATION USE INSTRUCTIONS for specific directions for use.

USE DIRECTIONS

Apply lower rates of BOTRAN 5F FUNGICIDE when disease pressure is anticipated to be light. Use higher rates for severe disease pressure.

USE RESTRICTION

- Do not apply through backpack mist blower.

COMPATIBILITY

BOTRAN 5F FUNGICIDE can normally be mixed with other fungicides, insecticides, or defoliants. It is suggested that such combinations be used on a small number of plants before treating large areas. However, combination sprays of BOTRAN 5F FUNGICIDE and miscible oil formulations of insecticides, particularly organic phosphorous compounds, may cause plant injury. Check for compatibility before large-scale treatment. Do not combine with any products unless previous use or small-scale treatment has proven mixture is physically compatible and safe to plants. When mixing BOTRAN 5F FUNGICIDE with any other product, add wettable powder or dry flowable formulations first, and then add liquid products.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DILUTION DIRECTIONS

The rate required for thorough, uniform coverage varies with plant growth at time of application. Except as specified for certain uses, the following rates are intended to cover a broad range of conditions.

DILUTE APPLICATION: Apply specified rate in 50-100 gals. of water per acre.

CONCENTRATE APPLICATION: Apply specified rate in not less than 10 gals. of water per acre.

AIR APPLICATION: Apply specified rate in a minimum of 5 gals. of water per acre.

PREHARVEST INTERVAL

Do not exceed the maximum rate of BOTRAN 5F FUNGICIDE per acre or the time limitations specified for the individual crops.

DIRECTIONS FOR USE ON ROW CROPS

CROP	DISEASE	DOSAGE QTS./ACRE	COMMENTS
CELERY	Pink rot <i>Sclerotinia sclerotiorum</i>	1.2* – 2 OR 3.2	Multiple applications: Begin application when disease is anticipated or approximately 10 weeks before harvest. Repeat at 7-day intervals in summer or 14-day intervals in fall and winter. May be applied up to 7 days before harvest. OR Single Application: If early disease is not likely, make a single application of 3.2 qts. of BOTRAN 5F FUNGICIDE per acre 7 days before harvest. Use a sprayer with a drop nozzle boom to direct spray to the base of plants and adjacent soil.
	Restrictions: <ul style="list-style-type: none">• Pre-harvest Interval: 7 days• *Do not apply less than 2 quarts (2.5lbs AI) per application in California.• Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per application.• Do not apply more than 4 quarts (5lbs AI) formulated product per acre per year (5lbAI/acre/year).		
FLORENCE FENNEL	Pink rot <i>Sclerotinia sclerotiorum</i>	1.2* – 2 OR 3.2	Multiple applications: Begin application when disease is anticipated or approximately 10 weeks before harvest. Repeat at 7-day intervals in summer or 14-day intervals in fall and winter. May be applied up to 7 days before harvest. OR Single Application: If early disease is not likely, make a single application of 3.2 qts. of BOTRAN 5F FUNGICIDE per acre 7 days before harvest. Use a sprayer with a drop nozzle boom to direct spray to the base of plants and adjacent soil.
	Restrictions: <ul style="list-style-type: none">• Pre-harvest Interval: 7 days• *Do not apply less than 2 quarts (2.5lbs AI) per application in California.• Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per application.• Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per year.		
ENDIVE (ESCAROLE)	Gray mold rot <i>Botrytis cinerea</i>	1.6	Apply 7 days following transplanting. Repeat application when plants are half-mature. Do not apply to wilted plants or seedlings. OR Apply at thinning and repeat application 7 days later. Use a sprayer with a drop nozzle boom to direct spray to the lower portion of plants and surrounding soil.
	Restrictions: <ul style="list-style-type: none">• Pre-harvest Interval: 14 days• Do not apply more than 1.6 quarts (2lbs AI) formulated product per acre per application.• Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per year.• Notes: Some leaf bronzing may be noted soon after application, but the effect is of a temporary nature.• Two applications of BOTRAN 5F FUNGICIDE per year may be applied.		
LETTUCE (LEAF AND HEAD)	Drop rot Watery soft rot <i>Sclerotinia sclerotiorum</i> <i>Sclerotinia minor</i> Gray mold rot <i>Botrytis cinerea</i>	0.6	At time of planting: Apply 0.6 qt. in a 4-6 inch band over seedlings after seeding or before transplanting. Transplants may be sprayed following planting.
		0.6 -1.8	Prethinning: Apply 0.6 - 1.8 qts. as a directed spray over unthinned seedlings.
		1.8 - 3.2	Post-thinning (direct seeded) and Established Transplants: Apply 1.8 - 3.2 qts. as a basal drench in 50-100 gals. of water as soon after thinning as possible. Following this treatment, soil surrounding the plants should not be disturbed. A second post-thinning treatment may be made if significant soil disturbance occurs or high soil moisture conditions exist.
	Restrictions: <ul style="list-style-type: none">• Pre-harvest Interval: 14 days• Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per application.• Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per year.• Note: One or more of the above treatments may be made to the same crop; however, do not apply more than 3.2 qts. (4lbs AI) of Botran 5F Fungicide per acre per crop or within 14 days of harvest.• Note: When applying in a band, do not reduce the per acre rate.• Certain environmental conditions or tank-mix combinations may cause minor leaf bronzing which should be temporary and quickly outgrown.		

ONIONS, GARLIC, SHALLOTS	White rot <i>Sclerotium cepivorum</i>	2 – 3.2	Direct seeded: For white rot control, apply a 5 inch band over seed row/bed and incorporate in to 1.5 inches of soil 1-2 weeks before onions are seeded. Use restricted to eastern Oregon, eastern Washington, and western Idaho. Planting sets or buds: For white rot control in furrow applications, use low-pressure spray equipment. Use 50 mesh or coarser screens. Center the front nozzle on the furrow. Spray a 5-inch band on 40-inch beds.
	Leaf blight <i>Botrytis</i> rot <i>Botrytis cinerea</i>	1 – 2	Apply when disease is anticipated. Repeat at 14-day intervals if necessary.
	Neck rot <i>Botrytis allii</i> (except CA)	1 -2	Apply when disease is anticipated. Repeat at 14-day intervals if necessary.
	Restrictions: <ul style="list-style-type: none"> • Pre-harvest Interval: 14 days • Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per application. • Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per year. 		
SNAP BEANS	White mold <i>Sclerotinia sclerotiorum</i> (<i>Sclerotini</i> ose)	1.3	Bush varieties: Begin applications when disease is anticipated and continue at 7-day intervals during periods favorable to the development of disease.
		1.6 (except CA)	Pole varieties: Begin applications when disease is anticipated and continue at 7-day intervals during periods favorable to the development of disease.
	Restrictions: <ul style="list-style-type: none"> • Pre-harvest Interval: 2 days • Do not apply more than 1.6 quarts (2lbs AI) formulated product per acre per application. • Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per year. • Do not feed treated foliage to livestock. 		

DIRECTIONS FOR USE ON SWEET POTATOES

CROP	DISEASE	COMMENTS
SWEET-POTATOES	Scurf <i>Monilochaetes infuscans</i> Sclerotial blight <i>Sclerotium rolfsii</i>	Seed dip: Mix 0.6 quarts of Botran 5F Fungicide in 7.5 gals. of water. Dip seed sweet potatoes 10 to 15 seconds in a well-agitated Botran 5F Fungicide suspension. Drain sweet potatoes and bed promptly. Prepare fresh Botran 5F Fungicide suspension daily. Plant bed spray: For a 42 inch plant bed, mix 5.73 oz of Botran 5F Fungicide in 14 gals. of water per 1000 linear ft. of plant bed. Spray or sprinkle Botran 5F Fungicide suspension over bedded sweet potatoes before covering them with soil.
	Plant bed spray Restrictions: <ul style="list-style-type: none"> • Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per application. • Do not apply more than 3.2 quarts (4lbs AI) formulated product per acre per year. • Restriction: Do not plant tomatoes as a follow-up crop. 	
	Postharvest decay <i>Rhizopus</i> rot (except CA)	Postharvest: Mix 0.6 quarts of Botran 5F Fungicide rate in 100 gals. of water. Spray using 8001 fan-shaped nozzle at 50 - 70 psi immediately after washing while the potatoes are conveyed on chain rollers or belts. Do not rinse after treatment. In recirculating systems, as volume decreases, make up to original volume with suspension equivalent to 0.6 qt. of Botran 5F Fungicide per 100 gals. of water. Where black rot is a problem, do not recirculate used spray suspension. Root Dip: Dip roots for 5 to 10 seconds in well-agitated suspension of 0.6 quarts of Botran 5F Fungicide in 100 gallons of water. Do not rinse after treatment. Add 0.3 qt. Botran 5F Fungicide to 100 gals. of treating suspension after 500 bushels are treated. After each 1000 bushels treated, drain and flush tank. Refill with fresh dip suspension.

CHEMIGATION USE INSTRUCTIONS

Apply this product only through sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move, and drip irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

CHEMIGATION SPRINKLER CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ), or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the

public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. In addition, all directions and requirements specified for SPRINKLER OR DRIP CHEMIGATION SYSTEMS must be followed.

SPRINKLER OR DRIP CHEMIGATION SYSTEMS

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

BOTRAN 5F FUNGICIDE should be injected at the end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until all BOTRAN 5F FUNGICIDE has been cleared from the last emitter.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep closed. Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. After cleaning, if recycling is not available, puncture and dispose of in a sanitary landfill or by incineration or if allowed by State and local authorities by burning. If burned stay out of smoke.

Mini-Bulk Containers: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.

For other product information contact Gowan Company or see Material Safety Data Sheet

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE FULLEST EXTENT PERMITTED BY LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

BOTRAN® 5F is a registered trademark of Gowan Company, LLC.

Chemtrec® is a registered trademark of American Chemistry Council, Inc.

EPA TEXT: BOTRAN 5F FUNGICIDE (TO EPA 7-6-20)