

ACCEPTED
SUPPLEMENTAL LABELING FOR TANK-MIX WITH EPTAM 7-E ON BEANS
 MAR 16 2005
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. 10163-254

Sandea®

Herbicide

Sandea is a selective herbicide for control of listed broadleaf weeds and nutsedge

ACTIVE INGREDIENT:	% BY WT.
* Halosulfuron-methyl	75.0%
OTHER INGREDIENTS	25.0%
	TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

A tank-mix combination of Sandea Herbicide plus EPTAM 7-E will give a broader spectrum of weed control than either product used separately. **Caution:** Read both the Sandea Herbicide and EPTAM 7-E labels carefully before using. Observe all cautions and limitations on labeling of both products.

CROP	RATE/ACRE	COMMENTS
DRY BEANS	Sandea @ 1/2 – 2/3 oz. PLUS EPTAM 7-E @ 3 1/2 – 4 1/2	Apply uniformly with ground equipment in a minimum of 15 gallons of water per acre. PREPLANT OR AT PLANTING • Incorporation: Apply and incorporate 1/2 to 2/3 ounce Sandea and 3-1/2 to 4-1/2 pints EPTAM 7-E per acre to a depth of approximately 2 inches just before planting. Use lower rate on lighter textured soils with low organic matter. Refer to EPTAM 7-E label for specific incorporation directions. Rotary hoe lightly during or shortly after emergence of the beans to break any crust which occurs.
		<ul style="list-style-type: none"> • Do not apply more than 2/3 ounce Sandea per acre per crop-cycle, not to exceed 2 ounces per acre per 12-month period (includes applications to the crop and to Row Middles/Furrows). • Do not use EPTAM 7-E on Adzuki beans, cowpeas (black-eyed peas, black-eyed beans), soybeans, lima beans, Mung beans, Garbanzo beans or other flat-podded beans except Romano. Under abnormal weather conditions, stunting may occur on Gratiot, Michilite, Sanilac, Seafarer, and Seaway varieties. Do not exceed 9 pints EPTAM 7-E per acre per crop. • Do not exceed 3-1/2 pints EPTAM 7-E per acre on small white beans or green beans grown on coarse textured soils. • Do not exceed 7 pints per acre per crop of Eptam in the Southwestern and Southeastern regions. Do not exceed 8 pints per acre per crop of Eptam in the Western Region. Do not exceed 9 pints per acre per crop of Eptam in the Pacific Northwestern Region. Do not exceed 9 3/4 pints of Eptam in the Northern Region. • Consult "Use Precautions" and "For Optimum Results" sections for important usage information.

**GOWAN'S SPECIAL CONDITIONS AND DISCLAIMER FOR USE OF SANDEA® HERBICIDE ON DRY BEAN CROPS
 USE OF SANDEA HERBICIDE ("PRODUCT") IN COMBINATION WITH EPTAM 7-E OR ANY OTHER PESTICIDE AND IN ACCORDANCE WITH
 THIS SUPPLEMENTAL LABEL MAY RESULT IN CROP INJURY, YIELD REDUCTION OR CROP LOSS. READ AND UNDERSTAND THESE
 CONDITIONS AND RISK OF USE BEFORE USING THE PRODUCT ON THE CROP.**

Gowan Company offers this Product only to end users and/or growers who expressly agree that the benefit and utility, in the sole opinion of the end user/grower, outweigh the extent of potential injury associated with the use of this Product. Due to the risk of crop damage, yield reduction or crop loss, all such use is at the end user/grower's risk.

By purchasing this Product for use, or using it, in the manner described in this Supplemental Label for Tank Mix with Eptam 7-E the user/grower expressly agrees and accepts that:

1. user/grower assumes all risks, including crop injury or damage, associated with using this Product as a mix with Eptam 7-E or any other mix partner;
2. Gowan Company does not make and does not authorize any agent or third party to make any representations, recommendations or warranties other than those expressly stated in this Supplemental Label and disclaims all warranties, express or implied, including any implied warranty of merchantability;
3. Gowan Company will not be held liable for any claims, damages, losses, expenses, or causes of action arising out of or related to the use of this Product; and
4. these Conditions supercede any contrary representations or recommendations made by Gowan Company or its agents or representatives, and any provisions in or any Product literature or labeling, including label attached to the Product container.

These terms and conditions are unacceptable, return Sandea at once unopened to the seller for a full refund.



SUPPLEMENTAL LABELING FOR TANK-MIX WITH EPTAM 7-E ON BEANS

Sandea® Herbicide

Sandea® is a selective herbicide for control of listed broadleaf weeds and nutsedge

ACTIVE INGREDIENT:

* Halosulfuron-methyl	75.0%
OTHER INGREDIENTS	25.0%
	TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

A tankmix combination of Sandea Herbicide plus EPTAM 7-E will give a broader spectrum of weed control than either product used separately.
Caution: Read both the Sandea Herbicide and EPTAM 7-E labels carefully before using. Observe all cautions and limitations on labeling of both products.

PREHARVEST INTERVAL

The required days between last application and harvest are given in () after crop name.

CROP	RATE/ACRE	COMMENTS
SUCCULENT SNAP BEANS (30)	Sandea @ 1/2 – 1 oz.	<p>Apply uniformly with ground equipment in a minimum of 15 gallons of water per acre. PREPLANT OR AT PLANTING</p> <ul style="list-style-type: none"> • Incorporation: Apply and incorporate 1/2 to 1 ounce Sandea and 3-1/2 to 4-1/2 pints EPTAM 7-E per acre to a depth of 2-3 inches just before planting. Use lower rate on lighter textured soils with low organic matter. Refer to EPTAM 7-E label for specific incorporation directions. Rotary hoe lightly during or shortly after emergence of the beans to break any crust which occurs.
	PLUS EPTAM 7-E @ 3 1/2 – 4 1/2 pts	
		<ul style="list-style-type: none"> • Do not apply more than 1 ounce Sandea per acre per crop-cycle, not to exceed 2 ounces per acre per 12-month period (includes applications to the crop and to Row Middles/Furrows). • Do not use EPTAM 7-E on flat-podded beans except Romano. Under abnormal weather conditions, stunting may occur on Gratiot, Michilite, Sanilac, Seafarer, and Seaway varieties. • Do not exceed 3-1/2 pints EPTAM 7-E per acre on small white beans or green beans grown on coarse textured soils. • Do not exceed 7 pints per acre per crop of Eptam in the Southwestern and Southeastern regions. Do not exceed 8 pints per acre per crop of Eptam in the Western Region. Do not exceed 9 pints per acre per crop of Eptam in the Pacific Northwestern Region. Do not exceed 9 3/4 pints of Eptam in the Northern Region. • Application of Sandea may cause significant, temporary stunting and delay maturity of snap beans resulting in delayed harvest. This product is available to the end-user/grower solely to the extent that the benefit and utility, in the sole opinion of the end user/grower, outweigh the extent of potential injury associated with the use of this product. Due to the risk of crop damage, all such use is at the end user/grower's risk. • Consult "Use Precautions" and "For Optimum Results" sections for important usage information.

**GOWAN'S SPECIAL CONDITIONS AND DISCLAIMER FOR USE OF SANDEA® HERBICIDE ON DRY BEAN CROPS
 USE OF SANDEA HERBICIDE ("PRODUCT") IN COMBINATION WITH EPTAM 7-E OR ANY OTHER PESTICIDE AND IN ACCORDANCE WITH
 THIS SUPPLEMENTAL LABEL MAY RESULT IN CROP INJURY, YIELD REDUCTION OR CROP LOSS. READ AND UNDERSTAND THESE
 CONDITIONS AND RISK OF USE BEFORE USING THE PRODUCT ON THE CROP.**

Gowan Company offers this Product only to end users and/or growers who expressly agree that the benefit and utility, in the sole opinion of the end user/grower, outweigh the extent of potential injury associated with the use of this Product. Due to the risk of crop damage, yield reduction or crop loss, all such use is at the end user/grower's risk.

By purchasing this Product for use, or using it, in the manner described in this Supplemental Label for Tank Mix with Eptam 7-E the user/grower expressly agrees and accepts that:

1. user/grower assumes all risks, including crop injury or damage, associated with using this Product as a mix with Eptam 7-E or any other mix partner;
2. Gowan Company does not make and does not authorize any agent or third party to make any representations, recommendations or warranties other than those expressly stated in this Supplemental Label and disclaims all warranties, express or implied, including any implied warranty of merchantability;
3. Gowan Company will not be held liable for any claims, damages, losses, expenses, or causes of action arising out of or related to the use of this Product; and
4. these Conditions supercede any contrary representations or recommendations made by Gowan Company or its agents or representatives, and any provisions in or any Product literature or labeling, including label attached to the Product container.

These terms and conditions are unacceptable, return Sandea at once unopened to the seller for a full refund.

