

10163-265

02/04/2010

1/9



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

FEB 04 2010

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms. Kyla Smith
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

Subject: Revised Storage and Disposal Statement

Dear Ms. Smith::

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated January 7, 2010 for:

EPA Registration 10163-265

Carzol SP

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, call me at 703 305-5409 or electronically at daniel.dani@epa.gov.

Sincerely,

Dani Daniel
Registration Division (7505P)
Insecticide/Rodenticide Branch

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060

Print Form



United States
Environmental Protection Agency
Washington, DC 20460

Registration
Amendment
Other **NOTIFICATION**
OPP Identifier Number
FEB 04 2010

Application for Pesticide - Section I

1. Company/Product Number 10163-265	2. EPA Product Manager George LaRocca	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Carzol SP	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) Gowan Company P.O. Box 5569 Yuma, AZ 85366 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Update Storage and Disposal

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted	If "Yes" Unit Packaging wgt. 2.5lb	No. per container 12	If "Yes" Package wgt	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
			No. per container	<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 30lbs		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Kyla S. Smith	Title Registration Specialist	Telephone No. (Include Area Code) 928-819-1531
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Specialist	
4. Typed Name Kyla S. Smith	5. Date 1-7-10	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethelene, polyvinyl chloride and viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Mixers, Loaders, Applicators, Flaggers and Other Handlers Must Wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Chemical-resistant gloves and apron for mixers and other handlers exposed to the concentrate
- Applicators using groundboom and air-blast equipment must wear a NIOSH-approved respirator with:
 - a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or
 - any N, R, P, or HE filter. N designation for respirator filters does not apply when application is made with oils.

See Engineering Controls for Additional Requirements.

Applicators Involved in Airblast Applications Must Wear:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical resistant gloves
- Protective eyewear
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear
- Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any R, P or HE filter).

User Safety Requirements: Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

Water-soluble packets when used correctly qualify as a closed mixing system under Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (5)]. Mixers and loaders using water-soluble packets must: wear the personal protective equipment required in the PPE section of this labeling for mixers and loaders, and be provided and must have immediately available for use in an emergency, such as a broken package, spill or equipment breakdown; chemical-resistant footwear and the respirator as specified in the PPE section of the label.

Pilots must use enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (6)].

USER SAFETY RECOMMENDATION

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical can contaminate surface water through spray applications. Under some conditions, it may also have a potential for runoff into surface water after application.

This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area. Do not allow animals to graze in treated orchard areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.


Do not apply this product in a way that will contact worker or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of:

- 9 days for citrus,
- 5 days for pome and stone fruits,
- 4 days for alfalfa grown for seed.



ORANGES, LEMONS, LIMES, TANGERINES, TANGELOS and GRAPEFRUIT (California Only)	Thrips	Apply 0.5 to 1.25 lbs. of CARZOL SP per acre as outside coverage spray when thrips appear.
	<ul style="list-style-type: none"> Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 9 days. Do not apply more than 1.25 lbs. CARZOL SP per acre in one crop season. Do not apply after fruit reach a diameter of one inch. If unharvested grapefruit and Valencia oranges are present from the previous crop, an application may be made to the new crop. However, a preharvest interval of 30 days must be observed for the unharvested crop. Note: If soft brown scale is present in the grove, a suitable scallicide should be applied after the CARZOL SP application. Aerial applications to orchard crops is prohibited. 	
APPLES	Tentiform Leafminer	Use 4 to 5 oz. CARZOL SP per 100 gallons of water at pink or petal fall. For full cover dilute spray, apply 400 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1 to 1.25 lbs. CARZOL SP per acre in at least 50 gallons of water.
	White Apple Leafhopper	Use 2 to 4 oz. CARZOL SP per 100 gallons of water when leafhoppers appear. For full cover dilute spray, apply 400 gallons of water per acre. For concentrate spray, apply 1/2 to 1 lb. CARZOL SP per acre in at least 50 gallons of water.
	Thrips	Use 4 to 5 oz. CARZOL SP per 100 gallons of water at bloom. For full cover dilute spray, apply 400 gallons of water per acre. For concentrate spray or dilute application of less than 400 gallons, apply 1 to 1.25 lbs. CARZOL SP per acre in a minimum of 50 gallons of water per acre. Apply at night or while bees are not working in blooms.
	European Red Mite, Twospotted Mite	Use 4 to 5 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 400 gallons of water per acre depending on the size of the trees and severity of the mite infestation. For concentrate spray, apply 1 to 1.25 lbs. CARZOL SP per acre in at least 50 gallons of water.
	<ul style="list-style-type: none"> Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 5 days. Do not apply more than 1.25 lbs. CARZOL SP per acre in one crop season. Late season applications to apples is prohibited; Only apply at petal fall. Aerial applications to orchard crops is prohibited. 	
APPLES (except California)	Campylomma	Use 4 oz. CARZOL SP per 100 gallons of water at pre-bloom/bloom. For full cover dilute spray, apply 400 gallons of water per acre. For concentrate spray or dilute application of less than 400 gallons, apply 1 lb. CARZOL SP per acre in a minimum of 50 gallons of water per acre. Apply at night or while bees are not working in blooms.
	<ul style="list-style-type: none"> Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 5 days. Do not apply more than 1.25 lbs. CARZOL SP per acre in one crop season. Late season applications to apples is prohibited; Only apply at petal fall. Aerial applications to orchard crops is prohibited. 	
PEARS	European Red Mite, Twospotted Mite	Use 4 to 5 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 400 gallons of water per acre depending on the size of the trees and severity of the mite infestation. For concentrate spray, apply 1 to 1.25 lbs. CARZOL SP per acre in at least 50 gallons of water.
	Pear Rust Mite	Use 4 oz. CARZOL SP per 100 gallons of water. For full cover dilute spray, apply 400 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1 to 1.25 lbs. CARZOL SP per acre in at least 50 gallons of water.
	<ul style="list-style-type: none"> Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 5 days. Do not apply more than 1.25 lbs. CARZOL SP per acre in one crop season. Do not apply this product after petal fall. A second, later season, application may be made to control stink bugs, lygus bugs, and pear rust mites in CA, OR, WA and ID, only after written approval by the State agency responsible for enforcement of FIFRA or authorized by that State agency. Aerial applications to orchard crops is prohibited. 	
PEACHES	European Red Mite, McDaniel Mite, Twospotted Mite	Use 4 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 400 to 500 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1 to 1.25 lbs. CARZOL SP per acre in at least 50 gallons of water.
	Lygus Bugs, Stink Bugs, Thrips	Use 4 oz. CARZOL SP per 100 gallons of water at first sign of infestation. For full cover dilute spray, apply 400 to 500 gallons of water per acre. For concentrate spray, apply 1 to 1.25 lbs. CARZOL SP per acre in at least 50 gallons of water.
	<ul style="list-style-type: none"> Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 5 days. Do not apply more than 1.25 lbs. CARZOL SP per acre in one crop season. Do not apply this product after petal fall. Aerial applications to orchard crops is prohibited. 	

PEACHES (except California)	Plum Curculio	Apply 1 lb. CARZOL SP as a concentrate spray of 100 gallons per acre, or less.
	<ul style="list-style-type: none"> Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 5 days. Do not apply more than 1.25 lbs. CARZOL SP per acre in one crop season. Do not apply this product after petal fall. Aerial applications to orchard crops is prohibited. 	
NECTARINES	Thrips	Use 4 oz. CARZOL SP per 100 gallons of water when thrips infest blossoms through petal fall. For full cover dilute spray, apply 400 gallons of water per acre depending on the size of the trees. For concentrate, apply 1 to 1.25 lbs. CARZOL SP per acre in at least 50 gallons of water. Apply at night or while bees are not working in blooms.
	European Red Mite, McDaniel Mite, Twospotted Mite	Use 4 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 400 to 500 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1 to 1.25 lbs. CARZOL SP per acre in at least 50 gallons of water.
	<ul style="list-style-type: none"> Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 5 days. Do not apply more than 1.25 lbs. CARZOL SP per acre in one crop season. Do not apply this product after petal fall. Aerial applications to orchard crops is prohibited. 	
ALFALFA (seed crop only in California, Oregon, Washington, Idaho and Nevada. Use pre-bloom only on alfalfa in all areas except California)	Lygus Bugs	Use ½ to 1 lb. CARZOL SP per acre. Apply when lygus bugs reach damaging numbers.
	Conspere Stink Bugs	Use 1 lb. CARZOL SP per acre for knockdown of a heavy nymphal population. A lygus control program will control stink bugs. When applied by air use a minimum of 5 gallons per acre. When applied by ground use a minimum of 20 gallons per acre.
	Twospotted Mite	Use ½ to 1 lb. CARZOL SP per acre. Apply when mites appear. When applied by air use a minimum of 10 gallons of spray per acre. When applied by ground use a minimum of 20 gallons per acre.
	<ul style="list-style-type: none"> Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 days. Hand harvesting prohibited. Use this product only on pure stands of alfalfa to be harvested for seed. Do not allow spray to drift onto adjacent crops. Do not spray while bees are visiting the treatment area. Do not allow livestock to graze treated fields. Do not use treated crops for livestock feed. Crop residues are not to be fed to livestock or baled for such use. Fields treated with CARZOL SP must be cleared after harvest by burning or by disking into the soil. CARZOL SP does not control aphids. If aphids appear, add an aphicide to the spray. Do not apply more than 1 lb. CARZOL SP per acre in one crop season. Allow 21 days before harvest. Observe the following rotational crop intervals following the last application of Carzol SP to alfalfa: <ol style="list-style-type: none"> 30 days for leafy vegetables 120 days for root crops 12 months for small grains and all other crops (except leafy vegetables and root crops) 	

AERIAL SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses, or to applications using dry formulations.

- The distance of the outer-most nozzles on the boom must not exceed 75% the length of the wingspan or rotor.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
- Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.
- Do not apply when gusts or sustained winds exceed 15 mph.

Where States have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

AERIAL DRIFT REDUCTION ADVISORY

This section is advisory in nature and does not supersede the mandatory label requirements.

Information on Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see Wind, Temperature, and Humidity, and Temperature Inversions).

Controlling Droplet Size:

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher-rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

9/9

Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length: For some use patterns, reducing the effective boom length to less than 75% of the wingspan or rotor length may further reduce drift without reducing swath width.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 3 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the wind is blowing away from adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store product in original container only, away from other pesticides, fertilizers, food or feed.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Open outer pouch and place soluble bag into application equipment. Then dispose of empty outer pouch in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DO NOT REUSE EMPTY CONTAINERS.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.

For other product information, contact GOWAN COMPANY or see Material Safety Data Sheet.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our recommendations for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label and is reasonably fit for the intended purpose referred to on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Carzol® is a registered trademark of Gowan Company, LLC.

Chemtreco® is a registered trademark of American Chemistry Council, Inc.

EPA Text: Carzol SP (sent notf 1-8-10)

