

10163-273

3/7/2004

ACCEPTED

MAR 7 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 10163-273

**SUPPLEMENTAL LABELING
FOR REVISED RATE ON EASTERN GRAPES**

Rubigan® E.C.

% By Wt.

ACTIVE INGREDIENT		
Fenarimol: a-(2-chlorophenyl)-a-(4-chlorophenyl)-5-pyrimidinemethanol	12.0%	
OTHER INGREDIENTS	88.0%	
	TOTAL	100.0%

Contains 1 pound active ingredient per gallon.
Contains petroleum distillate.

**KEEP OUT OF REACH OF CHILDREN
WARNING – AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the EPA registered label are to be followed

**Grapes: Disease Control Rates for Eastern United States (East of the Rocky Mountain Continental Divide)
Rubigan E.C.¹**

Crop Disease Growth Stage	Use rate per acre (fl oz)	Maximum spray interval (days)	Use Directions and Restrictions
Grapes			<p>Application timing: Rubigan E.C. may be applied alone or in tank mix combinations with recommended rates of a protectant fungicide, insecticide and/or miticide. Begin applications as recommended by your State Extension Specialist or Agricultural Experiment Station, or Certified Pest Control Advisor. Use rates per acre as specified by the crop phenology.</p> <p>Restrictions: Do not apply more than 6 fl oz of Rubigan E.C. per acre per application and 19 fl oz per acre per season.</p> <p>Preharvest interval: Do not apply within 30 days of harvest.</p>
Powdery Mildew			
Pre-bloom sprays	3-4	14-18	
Post bloom	5-6	14-18	
Summer sprays	5-6	14-18	
<p>Note: Using higher label rates of Rubigan E.C. during the "pre-bloom" stage of growth may result in foliar effects consisting of leaf cupping and/or darker leaf color.</p>			

¹Use a surfactant when Rubigan E.C. is applied alone.

Use of Surfactants

When used alone, tank mix Rubigan E.C. with a surfactant. Surfactants compatible with Rubigan E.C. include R-56 Spreader Sticker, Wilbur Ellis Co.; Bio-88 Spreader Activator, Kalo Laboratories; Regulaid, Kalo Laboratories; Leafact 80A Activator, Puregro Co.; Induce, Helena Chemical Co.; LI-700, Loveland Industries; Triton B-1956, Rohm and Haas, Latron B-1956, Rohm and Haas, or others where prior use has shown compatibility. Certain states require registration of surfactants. Where applicable, verify that the surfactant is registered for agricultural use in your state.

EPA Reg. No. 10163-273
EPA Est. No. 67545-AZ-1



Gowan Company
P. O. Box 5569
Yuma, AZ 85366-5569