

SUPPLEMENTAL LABELING For use on Pecans Rubigan® E.C.

ACTIVE INGREDIENT

% By Wt.

Fenarimol: <i>a</i> -(2-chlorophenyl)- <i>a</i> -(4-chlorophenyl)-5-pyrimidinemethanol	12.0%
OTHER INGREDIENTS	88.0%
TOTAL 100.0%	

Contains 1 pound active ingredient per gallon.
Contains petroleum distillate.

KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the EPA registered label are to be followed

**Pecans: Disease Control recommendations
Rubigan E.C.**

CROP	USE DIRECTIONS AND RESTRICTIONS
Pecans	<p>Use Rates: Do not apply less than 9 fl. ozs. or more than 12 fl. ozs. of Rubigan E.C. per acre. Use the higher range of the Rubigan E.C. rates during conditions suitable for heavy disease infection (i.e., for scab, several hours of leaf wetting at temperatures above 50°).</p> <p>Spray Volume Per Acre: Dilute Spray (200 or more gallons per acre): Apply Rubigan E.C. at the rate of 3 or more fl. ozs. per 100 gallons of water in a uniform spray applied to the point of run-off with conventional ground spray equipment. Do not exceed 12 fl. ozs. per acre.</p> <p>Concentrate Spray: (less than 200 gallons per acre), use ground application equipment only: Apply Rubigan E.C. at 9 to 12 fl. ozs. per acre.</p> <p>Application Schedule: For the control of scab and powdery mildew, apply Rubigan E.C. at 7 to 10 day schedule starting at bud break. Applications should continue up to and including the point where nuts are up to one-half inch (1/2") long. Thereafter, Rubigan E.C. should be used in tank-mix combination with a protectant fungicide at its recommended use rate. The Rubigan E.C. application prior to switching to a program utilizing a protectant-only application should include a protectant fungicide (a nonsterol-inhibiting fungicide with limited curative activity) as a tank-mix combination.</p> <p>Do not use Rubigan E.C. on the Wichita variety.</p> <p>Do not apply Rubigan E.C. within 30 days of harvest.</p> <p>Do not apply more than 84 fl. ozs. of Rubigan E.C. per acre per season on Pecans</p>

EPA Reg. No. 10163-273
EPA Est. No. 67545-AZ-1

ACCEPTED

SEP 20 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 10163-273

Gowan Company
P. O. Box 5569
Yuma, AZ 85366-5569

Submitted to EPA 08/31/04