

10163-277

2/6/2013

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

FEB 6 2013

Ms Kyla S. Smith
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Subject: Label Amendment – New Uses on Alfalfa and Timothy

EPA Registration No. 10163-277
Onager® Miticide
D#: 456988

Dear Ms Smith:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable. At your next label printing, or within eighteen (18) months, whichever comes first, you must incorporate this supplemental labeling into the main product labeling. Two (2) copies of the final printed label must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

If you have questions regarding this action, please contact Olga Odiott at (703) 308-9369.

Sincerely,

A handwritten signature in black ink, appearing to read "Mark Suarez".

Mark Suarez
Product Manager 13
Insecticide Branch
Registration Division (7505P)

Enclosure

GROUP ACARICIDE

Onager[®]

Miticide

EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY

ACTIVE INGREDIENT: Hexythiazox	% By Wt.
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide	11.8%
OTHER INGREDIENTS	88.2%
	TOTAL 100.00%

Onager contains 1.0 lb. active ingredient per gallon
Contains Petroleum Distillate

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

ACCEPTED

FEB 6 2013

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10163-277

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

PREHARVEST INTERVAL

The required days between the last application and harvest are given in () after each crop name.

APPLICATION DIRECTIONS

CROP	PEST	RATE Oz./acre	COMMENTS
Alfalfa (West of the Rockies)	Twospotted spider mite	10 - 24	Apply at the first sign of mites before populations begin to build. Ground Application: Apply the recommended rate of Onager by ground equipment at 15 - 20 GPA. Air Application: Apply the recommended rate of Onager by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems on Section 3 label.
			<ul style="list-style-type: none"> • Do not apply within 14 days of harvest. • Do not apply more than a total of 24 oz. formulated product per acre per year. • Do not make more than one application per year.
Timothy (West of the Rockies)	McDaniel spider mite, Twospotted spider mite, Banks grass mite	10 - 24	Apply at the first sign of mites before populations begin to build. Ground Application: Apply the recommended rate of Onager by ground equipment at 15 - 20 GPA. Air Application: Apply the recommended rate of Onager by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems on Section 3 label.
			<ul style="list-style-type: none"> • Do not apply within 14 days of harvest for hay. • Animals may graze treated fields on the day of application. • Do not apply more than a total of 24 oz. formulated product per acre per year. • Do not make more than one application per year.

RESTRICTIONS ON ALL CROPS

- Do not graze or feed livestock on cover crops growing in treated areas.
- Onager is an emulsifiable concentrate to be diluted with water for application in commercial plantings only.
- Apply Onager prior to adult mite build up. Onager will not control adult spider mites. Use higher rates on moderate to high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium to high populations, better results may be obtained using Onager in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager.
- Do not make more than one application of Onager or any other Hexythiazox product to the same crop per year.
- Do not plant rotational crops other than those on this label within 120 days of this application.

NET CONTENTS _____



EPA Reg. No. 10163-277
EPA Est. No.

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366