



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 2, 2020

Ms. Kyla S. Smith
Registration Specialist
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

Subject: Label Amendment – Agency Initiated Amendment to Correct Pounds AI per Gallon and Associated Use Rates, Remove Aerial Application Restriction, and Other Updates
Product Name: GWN-1708 MITICIDE/INSECTICIDE
EPA Registration Number: 10163-297
Application Date: January 23, 2020
Decision Number: 559376

Dear Ms. Smith:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is attached for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

SEE NEXT PAGE

Page 2 of 2
EPA Reg. No. 10163-297
Decision No. 559376

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Debra Rate by phone at (703) 306-0309, or via email at rate.debra@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'Michael Walsh', written in a cursive style.

Michael Walsh
Product Manager 11
Invertebrate & Vertebrate Branch 2
Registration Division
Office of Pesticide Programs

Attachment

FENAZAQUIN	GROUP	21A	ACARICIDE
FENAZAQUIN	GROUP	39	FUNGICIDE

GWN-1708 MITICIDE/INSECTICIDE

[MAGUS MITICIDE]

ACTIVE INGREDIENT: Fenazaquin
4-tert-butylphenethyl quinazolin-4-yl ether (IUPAC).....18.79%
OTHER INGREDIENTS:.....81.21%
TOTAL 100.0%

Contains 1.7 pounds active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN


WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have the person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT CALL 1-888-478-0798.	
NOTE TO PHYSICIAN	
If ingested, the product may cause irritation of the GI tract, vomiting and diarrhea. Treatment should be symptomatic and supportive. No antidote is recommended.	

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS WARNING/AVISO

WARNING: May be fatal if swallowed. Harmful if inhaled. Causes moderate eye irritation. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Avoid breathing dust. Remove and wash contaminated clothing before reuse. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.



NET CONTENTS ___ QUARTS



PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants (Coveralls for mixing/loading and applying with high-pressure handwand)
- Chemical-resistant gloves
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or residues on blooming crops or weeds. **Do not apply this product or allow it to drift to blooming crops, ornamentals, or weeds if bees or other pollinating insects are visiting the treatment area.**

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu* or directly to EPA at: beekill@epa.gov‡

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls

- Chemical-resistant gloves
- Shoes plus socks

Additional Requirements for Early Entry into Greenhouse and Treated Indoor Areas

Do not enter a treated greenhouse or a treated indoor area without protective equipment for 12 hours unless one of the following items is completed:

- 10 air exchanges
- 2 hours of system ventilation
- 4 hours of ventilation using vents, windows or other passive ventilation

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas or vicinity where there may be drift. Do not enter treated areas without protective equipment until sprays have dried.

USE RESTRICTIONS:

Use Pattern

- Apply this product only on ornamentals, Christmas tree plantations, and non-bearing tree fruits and nuts to control mites and whiteflies as described in the mixing and application below. Apply only for the sites, pests, and application methods described on this label.

Prohibited Methods of Application

- **Chemigation:** Do not apply this product through any type of irrigation system.

Protection of Workers or Other Persons

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Only commercial applicators may apply product to residential areas.

State or Tribe Requirements

- For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Repackaging and Reformulation

- For end use only. Do not repack or reformulate without manufacturer's written approval.

Notice of Conditions of Sale

- Follow all applicable directions, restrictions, precautions and Notice of Conditions of Sale and Warranty and Liability Limitations.

Label in User's Possession

- This labeling must be in the user's possession during application.

Bee Restriction

- Do not apply this product or allow it to drift to blooming crops, ornamentals, or weeds if bees or other pollinating insects are visiting the treatment area.

Water Body Buffer

- Do not apply within 100 feet of water bodies (including, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

USE PRECAUTIONS:

Stressed Plants

- Do not spray plants under stress.

Phytotoxicity

- GWN-1708 has been used successfully on many ornamental plants species. Do not treat new species or varieties without first testing a few plants on a small scale.
- Not all Rose varieties have been tested. Phytotoxicity is likely to occur on some varieties of roses (Particularly long stem and cut varieties). Avoid direct spray of blooms and test on a small scale first to ensure crop safety. Applications made to roses are at grower risk.

PRODUCT INFORMATION:

GWN-1708 is a suspension concentrate formulation that acts by contact to provide rapid knockdown of mites and whiteflies.

MODE OF ACTION AND RESISTANCE

GWN-1708 is a METI (mitochondrial electron transport inhibitor) classified as Group 21. To prevent or delay resistance development, only apply a single application of **GWN-1708** per crop season. If additional chemical control is required, rotate to a product with a different mode of action grouping.

RESISTANCE MANAGEMENT

For Use as an Acaricide:

For resistance management, GWN-1708 contains a Group 21 acaricide. Any insect/mite population may contain individuals naturally resistant to GWN-1708 and other Group 21 acaricides. The resistant individuals may dominate the insect/mite population if this group of acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed

To delay acaricide resistance, take the following steps:

- Rotate the use of GWN-1708 or other Group 21 acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):

- Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
- Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
- When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact a Gowan Company representative.

For Use as a Fungicide:

For resistance management, GWN-1708 contains a Group 39 fungicide. Any fungal/bacterial population may contain individuals naturally resistant to GWN-1708 and other Group 39 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of GWN-1708 or other Group 39 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact a Gowan Company representative. You can also contact your pesticide distributor or university extension specialist to report resistance.

MIXING DIRECTIONS

Dilution Rate

Dilute **GWN-1708** in sufficient water to ensure application thoroughly covers foliage. Maintain agitation during mixing.

Additives

In general, no additives are necessary for effective use of **GWN-1708**.

However, in situations where local conditions such as hard water are a problem, use adjuvants or wetting agents to achieve thorough spray coverage.

GENERAL TANK MIXING INFORMATION

Phytotoxicity

The phytotoxic potential of **GWN-1708** has been assessed on a wide variety of common ornamental plants with no phytotoxicity observed. However, all plant species and their varieties and cultivars have not been tested with possible tank mix combinations, sequential pesticide treatments, and adjuvants and surfactants. Local conditions can also influence crop tolerance and may not match those under which testing has been conducted. Therefore, before using GWN-1708, test the product on a sample of the crop to be treated to ensure that a phytotoxic response will not occur as a result of applications.

Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the **Mixing Order** (see below) using 2 teaspoons for each pound or 1 teaspoon for each 16ozs of labeled rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order

- 1) **Water:** Begin by agitating a thoroughly clean sprayer tank half full of clean water.
- 2) **Water-dispersible products:** (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
- 3) **Emulsifiable concentrates**
- 4) **Water-soluble Products**
- 5) **Water-soluble additives**
- 6) **Remaining quantity water**

Maintain constant agitation during application.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

APPLICATION DIRECTIONS**Application Equipment**

Application can be made with the following types of low or high volume spray equipment: machine ground spray application (open cab groundboom), ground spray application (open cab airblast sprayer), high pressure hand wand sprayer, backpack sprayer and aerial.

Constant Agitation Before and During Application

Shake well before using and frequently while applying. If agitation is stopped for an extended period of time, the spray solution must be thoroughly re-mixed prior to further use.

Importance of Coverage

Thorough coverage is required for optimum control.

Application Rates

Apply GWN-1708 at rates in Table 1. Do not apply during conditions when product will drift to all other crops and non-target areas.

Table 1. Application Rates Per Cropping and Year for Sites and Pests Controlled by GWN-1708 Miticide/Insecticide

SITES	PESTS	APPLICATION RATES
Ornamental plants (greenhouse, shade houses)	MITES, WHITEFLIES, POWDERY MILDEW	<p>Mites = 12 - 24 fl oz per 100 gallons of water 12 - 24 fl oz /acre per cropping (0.28 - 0.55 fl oz/1000 sq ft/per cropping) (0.16 - 0.32 lb ai/acre per cropping)</p> <p>Whiteflies = 18 - 24 fl oz per 100 gallons of water 18-24 fl oz/ acre per cropping (0.41 - 0.55 fl oz/1000 sq ft/per cropping) (0.24- 0.32 lb ai/acre per cropping)</p>
<p>Restrictions:</p> <ul style="list-style-type: none"> Do not make more than one application per cropping. Do not exceed 24 ozs (0.32 lb ai) per acre per year for outdoor applications Do not exceed 48 ozs (0.64 lb ai) per acre per year for indoor applications 		
Christmas Tree Plantations and Ornamental plants (field grown, outdoor)	MITES, WHITEFLIES, POWDERY MILDEW	<p>Mites = 12 - 24 fl oz per 100 gallons of water 12 - 24 fl oz /acre per cropping (0.28 - 0.55 fl oz/1000 sq ft/per cropping) (0.16 - 0.32 lb ai/acre per cropping)</p> <p>Whiteflies = 18 - 24 fl oz per 100 gallons of water 18-24 fl oz/ acre per cropping (0.41 - 0.55 fl oz/1000 sq ft/per cropping) (0.24 - 0.32 lb ai/acre per cropping)</p> <p>Air Application: Apply the labeled rate of GWN-1708 by air in a minimum of 10 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.</p>
<p>Restrictions:</p> <ul style="list-style-type: none"> Do not make more than one application per cropping. Do not exceed 24 ozs (0.32 lb ai) per acre per year for outdoor applications 		
Non-bearing tree fruits and Nuts (field grown and nursery) <i>Non-bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application</i>	MITES, WHITEFLIES, POWDERY MILDEW	<p>Mites = 12 - 24 fl oz per 100 gallons of water 12 - 24 fl oz /acre per cropping (0.28 - 0.55 fl oz/1000 sq ft/per cropping) (0.16 - 0.32 lb ai/acre per cropping)</p> <p>Whiteflies = 18 - 24 fl oz per 100 gallons of water 18-24 fl oz/ acre per cropping (0.41 - 0.55 fl oz/1000 sq ft/per cropping) (0.24 - 0.32 lb ai/acre per cropping)</p>
<p>Restrictions:</p> <ul style="list-style-type: none"> Do not make more than one application per cropping. Do not exceed 24 ozs (0.32 lb ai) per acre per year for outdoor applications Do not exceed 48 ozs (0.64 lb ai) per acre per year for indoor applications 		
Established Ornamental Landscape Plantings , Including those in: interiorscapes, residences, public areas, commercial areas, institutional areas, industrial areas, rights of way and other easements, and recreational sites such as campgrounds, golf courses, parks and athletic fields	MITES, WHITEFLIES, POWDERY MILDEW	<p>Mites = 12 - 24 fl oz per 100 gallons of water 12 - 24 fl oz /acre per cropping (0.28 - 0.55 fl oz/1000 sq ft/per cropping) (0.16 - 0.32 lb ai/acre per cropping)</p> <p>Whiteflies = 18 - 24 fl oz per 100 gallons of water 18-24 fl oz/ acre per cropping (0.41 - 0.55 fl oz/1000 sq ft/per cropping) (0.24- 0.32 lb ai/acre per cropping)</p>
<p>Restrictions:</p> <ul style="list-style-type: none"> Do not make more than one application per cropping. Do not exceed 24 ozs (0.32 lb ai) per acre per year for outdoor applications Do not exceed 48 ozs (0.64 lb ai) per acre per year for indoor applications 		

Amount of GWN-1708 Miticide/Insecticide and Water to Apply

Use Table 2 to calculate the amount of GWN-1708 Miticide/Insecticide and water to apply.

Table 2. Calculation of Amount of GWN-1708 Miticide/Insecticide and Water to Apply to Treat 1000 sq ft and 1 Acre at Rates in Table 1.¹

To treat ...	at this rate ...	use this amount of:	
		Water (gallons)	GWN-1708 Miticide/Insecticide
1000 sq ft	0.28 fl oz/1000 sq ft (0.16 lb ai/acre)	2.3	0.28 fl oz or 1.7 teaspoons
	0.41 fl oz/1000 sq ft (0.24 lb ai/acre)	2.3	0.41 fl oz or 2.5 teaspoons
	0.55 fl oz/1000 sq ft (0.32 lb ai/acre)	2.3	0.55 fl oz or 3.3 teaspoons
1 Acre	12 fl oz/acre (0.16 lb ai/acre)	100	12 fl oz
	18 fl oz/acre (0.24 lb ai/acre)	100	18 fl oz
	24 fl oz/acre (0.32 lb ai/acre)	100	24 fl oz

¹To calculate the total amount of product needed, multiply the amount of product by 1) number of acres or 2) number of 1000 sq ft areas (total sq ft/1000).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place away from heat or open flame.**Pesticide Disposal:** Pesticide wastes are acutely hazardous. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law.

If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in sanitary landfill, or incineration, or other procedures allowed by state and local authorities.**FOR 24 HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE). CALL CHEMTREC® (800) 424-9300****NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS****Important:** Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. All such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Chemtrec® is a registered trademark of American Chemistry Council, Inc.

EPA Text Pending: GWN-1708 (To EPA 1-23-20, 8-6-20, 8-20-20, 9-1-20)