



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

May 25, 2017

Ms. Kyla Smith  
Registration Specialist  
Gowan Company  
P.O. Box 5569  
Yuma, AZ 85366-5569

Subject: PRIA Label Amendment - Adding Hops & Tree Nuts Group 14-12 (Includes Almonds)  
Product Name: Magister SC Miticide  
EPA Registration Number: 10163-322  
Application Dates: December 5, 2015; January 5, 2016  
Decision Numbers: 512803; 512016

Dear Ms. Smith:

The application referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable under FIFRA Section 3(c)(7)(B), subject to the following conditions:

1. You must submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.
2. As per the conditions of registration of EPA Reg. No. 10163-322 under 3(c)(7)(B) of FIFRA for uses on almonds and cherries (Agency letter dated April 30, 2015), the agency required several studies to be submitted. These studies included: 1) OECD TG 237: Honey bee larvae acute oral toxicity; 2) Honey bee adult chronic oral toxicity; and 3) Translocation study in plants. The deadline for you to submit these required data to the Agency is November 1, 2017. Your failure to provide these data in a timely or adequate manner may result in initiation of a cancellation action against your registration.
3. The EPA National Pesticide Standards Repository has a fenazaquin analytical standard with an expiration date of November 30, 2019; however, analytical standards for Metabolite M29 and Metabolite I are not available. You must submit analytical grade standards for Metabolite M29 and Metabolite I to the standard Repository.

A stamped copy of your labeling is attached for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR

**SEE NEXT PAGE**

152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). If you have any questions, please contact Debra Rate by phone at (703) 306-0309, or via email at [rate.debra@epa.gov](mailto:rate.debra@epa.gov).

Sincerely,



Mike Walsh, Product Manager 11  
Invertebrate & Vertebrate Branch 2  
Registration Division  
Office of Pesticide Programs

Attachment

# MAGISTER SC MITICIDE

**ACTIVE INGREDIENT:** Fenazaquin  
*4-tert-butylphenethyl quinazolin-4-yl ether (IUPAC)*.....18.79%  
**OTHER INGREDIENTS:**.....81.21%  
**TOTAL** 100.0%

Contains 1.6 pounds active ingredient per gallon.

## KEEP OUT OF REACH OF CHILDREN **WARNING/AVISO**


Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have the person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b><u>HOTLINE NUMBER</u></b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. <b>FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT CALL 1-888-478-0798.</b>	
<b><u>NOTE TO PHYSICIAN</u></b>	
If ingested, the product may cause irritation of the GI tract, vomiting and diarrhea. Treatment should be symptomatic and supportive. No antidote is recommended.	

### PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS **WARNING/AVISO**

WARNING: May be fatal if swallowed. Harmful if inhaled. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Avoid breathing dust. Remove and wash contaminated clothing before reuse. Avoid contact with eyes, skin or clothing.

NET CONTENTS \_\_\_ QUARTS



## PERSONAL PROTECTIVE EQUIPMENT (PPE)

### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants (Coveralls for mixing/loading and applying with high-pressure handwand)
- Chemical-resistant gloves
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls Statement:** When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fenazaquin from runoff water and sediment.

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.

### PROTECTION OF POLLINATORS



**APPLICATION RESTRICTIONS** EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

#### **This product can kill bees and other insect pollinators.**

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: [www.aapco.org/officials.html](http://www.aapco.org/officials.html). Pesticide incidents should also be reported to the National Pesticide Information Center at: [www.npic.orst.edu](http://www.npic.orst.edu)-or directly to EPA at: [beekill@epa.gov](mailto:beekill@epa.gov)

### PRODUCT INFORMATION

**MAGISTER SC** is a suspension concentrate formulation that acts by contact to provide rapid knockdown of mites. **MAGISTER SC** is a foliar miticide belonging chemical class quinazoline. **MAGISTER SC** provides control of eggs by contact and controls immature and adult mites by both contact and ingestion. **MAGISTER SC** has proven fungicidal activity. **MAGISTER SC** is a METI (mitochondrial electron transport inhibitor) classified as Group 21.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply until all varieties in the orchard to be treated are through petal fall. Aerial applications are prohibited. Do not apply by any type of irrigation system.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

### RESISTANCE MANAGEMENT

To prevent or delay resistance development, only apply a single application of **MAGISTER SC** per year. If additional chemical control is required, rotate to a product with a different mode of action grouping.

### APPLICATION INSTRUCTIONS

Apply this product to the crops listed on this label. Apply as described in the mixing and application section below. Apply only for the sites, pests, and application methods described on this label.

#### FOLIAR SPRAY APPLICATIONS

Foliar applications may be made using properly calibrated ground sprayers. Thorough and uniform coverage of plants, with direct contact of the spray mixture to the target pests is required for satisfactory control.

Apply **MAGISTER SC** in sufficient water to ensure thorough coverage of fruit and foliage. Applications can be made with low or high volume spray equipment; however, thorough coverage is required for optimum control.

Do not apply where thorough coverage of plant is not possible. Applications made with less than thorough coverage may result in slower activity and/or less overall control from a single application than an application made with higher gallonages.

See application volume directions by crop in the Application Directions table.

#### SPRAY DRIFT MANAGEMENT

Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

Do not apply within 25 ft. of lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, or commercial fish farm ponds.

For ground boom applications, apply with nozzle height no more than 4 feet above the ground or crop canopy and when wind speed is 10 mph or less at the application site as measured by an anemometer. Use medium or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles.

For orchard/vineyard airblast applications, do not direct spray above trees/vines and turn off outward pointing nozzles at row ends and outer rows. Apply only when wind speed is 3 -10 mph at the application site as measured by an anemometer outside of the orchard/vineyard on the upwind side. The applicator also must use all other measures necessary to control drift.

#### MIXING INSTRUCTIONS

Shake well before using and agitate frequently while applying. If agitation is stopped for an extended period of time, the spray solution must be thoroughly re-mixed prior to further use. Maintain agitation during mixing.

In general, no additives are necessary for effective use of **MAGISTER SC**. **MAGISTER SC** may be tank mixed with non-ionic surfactants.

#### Phytotoxicity

The phytotoxic potential of **MAGISTER SC** has been assessed on a wide variety of crops with no phytotoxicity observed. However, all plant species and their varieties and cultivars have not been tested with possible tank mix combinations, sequential pesticide treatments, and adjuvants and surfactants. Local conditions can also influence crop tolerance and may not match those under which testing has been conducted. Therefore, before using **MAGISTER SC**, test the product on a sample of the crop to be treated to ensure that a phytotoxic response will not occur as a result of applications.

#### Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test.

For 50 gallons per acre spray volume, use 8.25 cups (2000 ml) of water. For other spray volumes, adjust rates accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the **Mixing Order** (see below) using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

**Mixing Order**

- 1) **Water:** Begin by agitating a thoroughly clean sprayer tank half full of clean water.
- 2) **Water-dispersible products:** (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
- 3) **Emulsifiable concentrates**
- 4) **Water Soluble Products**
- 5) **Water-soluble additives**
- 6) **Remaining quantity water**

Maintain constant agitation during application.

Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Always follow the most restrictive label use directions.


**Pests Controlled by MAGISTER SC**


MITES	
<b>TETRANYCHIDAE</b>	
Brown mite	<i>Bryobia rubrioculus</i>
European red mite	<i>Panonychus ulmi</i>
McDaniel spider mite	<i>Tetranychus mcdanieli</i>
Pacific spider mite	<i>Tetranychus pacificus</i>
Strawberry spider mite	<i>Tetranychus turkestanii</i>
Spruce spider mite	<i>Oligonychus ununguis</i>
Twospotted spider mite	<i>Tetranychus urticae</i>
Willamette spider mite	<i>Eotetranychus willamettei</i>
<b>ERIOPHYIDAE</b>	
Plum nursery mite	<i>Aculus fockeui</i>
<b>DISEASES</b>	
Powdery Mildew	<i>Podosphaera clandestina</i>

**PREHARVEST INTERVAL**

The required days between the last application and harvest are given in ( ) after each crop name.

**APPLICATION DIRECTIONS**

CROP	RATE FL OZ/ACRE	LB A.I./ACRE	COMMENTS
<b>CHERRIES (3)</b> 	24 - 36	0.3 – 0.45	For best results, use a minimum of 50 gpa
			<ul style="list-style-type: none"> <li>• Do not make more than one application per year.</li> <li>• Do not exceed application rate of 36 fl oz (0.45 lb a.i.).</li> <li>• Do not apply until all varieties in the orchard to be treated are through petal fall.</li> </ul>
<b>CHRISTMAS TREE PLANTATIONS</b> (field grown, outdoor) <b>(0)</b>	12 - 24	0.15 - 0.3	
			<ul style="list-style-type: none"> <li>• Do not make more than one application per cropping.</li> <li>• Do not exceed 24 fl oz (0.3 lb a.i.) per acre per year.</li> </ul>

<b>HOPS (7)</b>	24 - 36	0.3 – 0.45	For best results, use a minimum of 100 gpa.
	<ul style="list-style-type: none"> <li>Do not make more than one application per year.</li> <li>Do not exceed application rate of 36 fl oz (0.45 lb a.i.).</li> </ul>		
<b>NON-BEARING TREE FRUITS AND NUTS</b> (field grown and nursery) <i>Non-bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application (0)</i>	12 - 24	0.15 - 0.3	
	<ul style="list-style-type: none"> <li>Do not make more than one application per cropping.</li> <li>Do not exceed 24 fl oz (0.3 lb a.i.) per acre per year for outdoor applications.</li> <li>Do not exceed 48 fl oz per acre per year for indoor applications.</li> </ul>		
 <b>TREE NUT CROP GROUP 14-12 (7)</b> African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these	24 - 36	0.3 – 0.45	For best results, use a minimum of 100 gpa.
	<ul style="list-style-type: none"> <li>Do not make more than one application per year.</li> <li>Do not exceed application rate of 36 fl oz (0.45 lb a.i.).</li> <li>Do not apply until all varieties in the orchard to be treated are through petal fall.</li> </ul>		

**RESTRICTIONS ON ALL CROPS:**

- Aerial applications are prohibited.
- Do not apply by any type of irrigation system.
- Only ground sprayer and airblast applications are permitted.
- Do not apply during conditions when product will drift to other crops and non-target areas.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place away from heat or open flame.

**Pesticide Disposal:** Pesticide wastes are acutely hazardous. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

**FOR 24 HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE). CALL CHEMTREC® (800) 424-9300**

**NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS**

**Important:** Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. All such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Chemtrece<sup>®</sup> is a registered trademark of American Chemistry Council, Inc.

EPA Text Pending: MAGISTER SC (TO EPA 5-8-17)



# SUPPLEMENTAL LABELING

GROUP	21	ACARICIDE
-------	----	-----------

# MAGISTER SC

MITICIDE

EPA REG. No. 10163-322

**For Control of twospotted spider mite and powdery mildew in hops.**  
**This supplemental label expires on May 31, 2020 and must not be used of distributed after this date.**

**ACTIVE INGREDIENT:** Fenazaquin  
*4-tert-butylphenethyl quinazolin-4-yl ether (IUPAC)*.....18.79%  
**OTHER INGREDIENTS:**.....81.21%  
 TOTAL 100.0%

Contains 1.6 pounds active ingredient per gallon.

## KEEP OUT OF REACH OF CHILDREN WARNING/AVISO


### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Magister SC before applying. Carefully following all precautionary statements and applicable use directions
- Use of Magister SC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Magister SC.

### DIRECTIONS FOR USE PREHARVEST INTERVAL


The required days between the last application and harvest are given in ( ) after each crop name.

#### APPLICATION DIRECTIONS

CROP	PEST	RATE FL OZ/ACRE	LB A.I./ACRE	COMMENTS
HOPS (7) 	Twospotted spider mite, powdery mildew	24 - 36	0.3 – 0.45	For best results, use a minimum of 100 gpa.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Do not make more than one application per year.</li> <li>• Do not exceed application rate of 36 fl oz (0.45 lb a.i.).</li> <li>• Aerial applications are prohibited.</li> <li>• Do not apply by any type of irrigation system.</li> <li>• Only ground sprayer and airblast applications are permitted.</li> <li>• Do not apply during conditions when product will drift to other crops and non-target areas.</li> </ul>				



\*See EPA Section 3 Label for "Protection of Pollinators" Information



Produced For:  
 Gowan Company  
 P.O. Box 5569  
 Yuma, AZ 85366-5569  
 EPA Text Pending: GWN-1708 F (EPA 5-8-16)



# SUPPLEMENTAL LABELING

GROUP	21	ACARICIDE
-------	----	-----------

# MAGISTER SC

MITICIDE

EPA REG. No. 10163-322

For control of mites in tree nut crop group 14-12

This supplemental label expires on May 31, 2020 and must not be used or distributed after this date.

**ACTIVE INGREDIENT:** Fenazaquin  
*4-tert-butylphenethyl quinazolin-4-yl ether (IUPAC)*.....18.79%  
**OTHER INGREDIENTS:**.....81.21%  
 TOTAL 100.0%

Contains 1.6 pounds active ingredient per gallon.

## KEEP OUT OF REACH OF CHILDREN WARNING/AVISO


### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Magister SC before applying. Carefully following all precautionary statements and applicable use directions
- Use of Magister SC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Magister SC.

### DIRECTIONS FOR USE PREHARVEST INTERVAL


The required days between the last application and harvest are given in ( ) after each crop name.

#### APPLICATION DIRECTIONS

CROP	PEST	RATE FL OZ/ACRE	LB A.I./ACRE	COMMENTS
 <b>TREE NUT CROP GROUP 14-12 (7)</b> African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these	McDaniel spider mite, Pacific spider mite, Twospotted spider mite, Willamette spider mite	24 - 36	0.3 – 0.45	For best results, use a minimum of 100 gpa.
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>Do not make more than one application per year.</li> <li>Do not exceed application rate of 36 fl oz (0.45 lb a.i.).</li> <li>Do not apply until all varieties in the orchard to be treated are through petal fall.</li> <li>Aerial applications are prohibited.</li> <li>Do not apply by any type of irrigation system.</li> <li>Only ground sprayer and airblast applications are permitted.</li> <li>Do not apply during conditions when product will drift to other crops and non-target areas.</li> </ul>				



\*See EPA Section 3 Label for "Protection of Pollinators" Information



Produced For:  
Gowan Company  
P.O. Box 5569  
Yuma, AZ 85366-5569

EPA Text Pending: GWN-1708 F (EPA 5-8-17)