

#### U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs

Biopesticides and Pollution Prevention Division (7511P) 1200 Pennsylvania Ave., N.W.

Washington, D.C. 20460

#### NOTICE OF PESTICIDE:

X Registration
Reregistration
(under FIFRA, as amended)

<b>EPA</b>	Reg.	Num	ber:
------------	------	-----	------

Date of Issuance:

10163-336

7/28/2016

Term of Issuance:

Unconditional

Name of Pesticide Product:

Captiva® Prime

Name and Address of Registrant (include ZIP Code):

Gowan Company, LLC PO Box 5569 Yuma, AZ 85366-5569

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

- 1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
- 2. Make the following labeling change before you release this product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 10163-336."

Signature of Approving Official:

Onclose Approving Official:

7/28/2016

Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)
Office of Pesticide Programs

Page 2 of 2 EPA Reg. No. 10163-336 OPP Decision No. 505825

3. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

Basic CSF dated 06/05/2015

Any CSFs other than those listed above are superseded.

If you have any questions, please contact Menyon Adams of my branch by phone at (703) 347-8496 or via email at adams.menyon@epa.gov.

Sincerely,

Andrew Bryceland, Team Leader Biochemical Pesticides Branch Biopesticides and Pollution Prevention Division (7511P) Office of Pesticide Programs

andrew . Buycelow

Enclosure

#### **MASTER LABEL**

## CAPTIVA® PRIME

Alternate Brand Name: **GWN-10285 INSECT REPELLENT / INSECTICIDE** 

### ACCEPTED

07/28/2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 40400 0000

10163-336

Active Ingredient	% By Wt.
Capsicum oleoresin extract*	7.60%
Garlic Oil	23.40%
Canola Oil	55.00%
Other Ingredients	14.00%
*contains 0.26% nominal concentration of Capsaicin and related capsaicinoids	Total: 100.00%

# KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID						
If on skin or clothing	Take off contaminated clothing					
	Rinse skin immediately with plenty of water for 15-20 minutes					
	<ul> <li>Call a poison control center or doctor for treatment advice.</li> </ul>					
	HOT LINE NUMBER					
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may						
also contact 1-888-478-07	also contact 1-888-478-0798 for emergency medical treatment information.					

# PRECAUTIONARY STATEMENT HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, and chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear chemical-resistant gloves. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

NET CONTENTS \_\_\_\_

EPA Reg No. 10163-XXX EPA Est No. (batch code XXX)





Produced For: Gowan Company, LLC. PO Box 5569 Yuma, AZ 85366-5569

Sublabel A: Agricultural Uses Sublabel B: Non-Agricultural Uses

### CAPTIVA® PRIME

### Alternate Brand Name: **GWN-10285 INSECT REPELLENT / INSECTICIDE**

Active Ingredient	% By Wt.
Capsicum oleoresin extract*	7.60%
Garlic Oil	23.40%
Canola Oil	55.00%
Other Ingredients	14.00%
*contains 0.26% nominal concentration of Capsaicin and related capsaicinoids	Total: 100.00%

# KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID						
If on skin or clothing	Take off contaminated clothing					
	Rinse skin immediately with plenty of water for 15-20 minutes					
	Call a poison control center or doctor for treatment advice.					
	HOT LINE NUMBER					
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-						
	y medical treatment information.					

# PRECAUTIONARY STATEMENT HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, and chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear chemical-resistant gloves. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

NET CONTENTS \_\_\_\_\_

EPA Reg No. 10163-XXX EPA Est No. (batch code XXX)





Produced For: Gowan Company, LLC. PO Box 5569 Yuma, AZ 85366-5569

PERSONAL PROTECTIVE EQUIPMENT (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

Users should

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash
  thoroughly and change into clean clothing.

#### **ENVIRONMENTAL HAZARDS**

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to industrial area below the mean high-water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all directions on the EPA approved label and any supplemental labeling

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read all Directions for Use carefully before applying.

#### **Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

#### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

#### **USE INFORMATION**

CAPTIVA PRIME is a repellent and insecticide formulated using concentrated plant extracts that can be used alone as a repellent or in tank mixes—with other biological or conventional insecticides or miticides to improve the level of control.

The repellent effect of CAPTIVA PRIME protects the crops from pests population establishment by its antifeedant and anti–egg laying properties. The irritant effect of CAPTIVA PRIME increases pest mobility, exposure and vulnerability to insecticides or miticides that act on contact. CAPTIVA PRIME weakens the cuticle of some immature stages of insect and mite pests.

Read all directions before using this product. Apply CAPTIVA PRIME directly to any food or non-food crop growing in field and greenhouses up to and including the day of harvest, at a maximum rate of 4 pints per acre. Refer to the **Insecticidal and Miticidal Uses Section** for a complete listing of crops.

#### MIXING AND APPLICATION

GROUND APPLICATIONS: CAPTIVA PRIME has a repellent and irritant effect on adult and immature stages of soft bodied insects and mites

- · Apply with most types of ground spray equipment.
- This product must come into contact with the targeted pests to be effective. For best results coverage typically requires a minimum of 10 gallons of total spray volume per acre. Higher volumes will be required for larger perennial crops.
- · For best preventive results, this product must be applied at least once a week on a regular and continuous basis.
- When pest pressure is heavy or plant canopy is dense, use higher rates and increase spray frequency. Repeat application if
  measurable rain occurs within two to three hours of spraying.

#### **AERIAL APPLICATIONS**

Apply CAPTIVA PRIME uniformly with properly calibrated equipment in 2 to 15 gallons of water per acre. Thoroughly clean equipment prior to mixing spray solution. Avoid streaking, skips, overlaps and spray drift during applications. Apply product only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g. when wind is blowing away from the sensitive areas). Repeat application if rain occurs before the product is dry.

#### MIXING:

- Shake well before using. Add required amount of CAPTIVA PRIME to a clean spray tank with at least one-half of the water to be sprayed. Agitate the mixture thoroughly, add other products and then fill the tank with the remaining water and continue agitation.
- · Always use this product promptly after mixing with water and do not let tank mix sit for more than 6 hours.
- Use water for dilution with a pH less than 8.
- CAPTIVA PRIME has been found to be compatible with most commonly used pesticides. To avoid problems, conduct a compatibility
  test before using this product in a tank mix with other pesticides. To test for compatibility, mix a small amount of each product, in
  the appropriate proportions, in a small jar.

PHYTOTOXICITY: CAPTIVA PRIME has been evaluated for phytotoxicity on a wide range of crops and ornamentals. However, since testing on all varieties of all crops and ornamentals is not feasible, testing a small portion of the area to be treated for phytotoxicity is recommended before treating the entire area. Further, all possible combinations or sequences of pesticide sprays, including other fertilizers, surfactants, adjuvants and other pesticides, have not been tested, thus testing for phytotoxicity of spray mixtures is recommended. It is further recommended that spray equipment used to apply CAPTIVA PRIME be thoroughly cleaned before use. The addition of spray adjuvants enhances control in some crops under ideal conditions. Addition of certain adjuvants may cause phytotoxicity therefore, the addition of crop oils and other adjuvants should be thoroughly tested before using. Highly alkaline products can cause unacceptable phytotoxicity and/or reduced effectiveness on target pests. Avoid tank mix combinations of CAPTIVA PRIME with compounds known to be incompatible with oil-based formulation to prevent phytotoxicity. Waxy bloom" on certain crops and ornamental plants may be reduced after a CAPTIVA PRIME application.

#### Insecticidal and Miticidal Uses

Use CAPTIVA PRIME as a foliar spray when pest pressure is low or use as a preventative management tool or integrated pest management system. When pest pressure is high use the product alone at its medium to high rates or in tank mix with other contact insecticides or miticides.

PREHARVEST INTERVAL: CAPTIVA PRIME has a zero day preharvest interval and can be applied up to and including the day of harvest.

TARGETED PESTS: Use CAPTIVA PRIME in a foliar spray to repel or suppress soft bodied pests.

- Mites (such as Tetranychus spp., Oligonychus spp., Panonychus spp., Polyphagotarsonemus latus, Phyllocoptruta oleivora, Brevipalpus phoenicis, Steneotarsonemus spinki)
- Thrips (such as Thrips spp., Frankliniella spp., Heliothrips spp., Scirtothrips citri.)
- Psyllids (such as Bactericera spp., Diaphorina citri, Cacopsylla spp.)
- Leafhoppers (such as Empoasca spp., Macrosteles spp., and Graphocephala spp.)
- Lepidoptera larvae (such as Spodoptera spp., Helicoverpa spp., Tuta absoluta, Thecla spp, Strymon basilides)
- White flies (such as Dialeurodes spp., Trialeurodes spp. Bemisia spp.)
- Other pests such as borers (Hypothenemus hampei), mealybugs (Planococcus spp., Pseudococcus spp., Dysmicoccus brevipes) and plant bugs (Leptoglossus spp., Nezara spp)

CROP GROUPS	USE SITES
BERRIES	Blackberry, Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Raspberry (black and red) (Note: For Strawberries – See Other.)
BULB VEGETABLES	Garlic, Leek, Onion (dry bulb, green and Welch), Shallot
CEREAL GRAINS AND COMMODITIES	Barley, Buckwheat, Corn, Millet (pearl and Proso), Oats, Popcorn, Rice, Rye, Sorghum (milo), Teosinte, Triticale, Wheat, Wild rice
CHRISTMAS TREES AND CHRISTMAS TREE PLANTATIONS	
CITRUS FRUIT GROUP	Calamondin, Citrus citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sour and sweet), Pummelo, Satsuma mandarin, White Sapote, Uniq Fruit
COTTON	

CUCURBIT VEGETABLES	Chayote, Chinese waxgourd, Citron melon, Cucumber, Gherkin, Gourd (edible), Muskmelon,
	Pumpkin, Squash (summer and winter), Watermelon
NONGRASS ANIMAL FEED	Alfalfa, Clover, Trefoil or Vetch.
FRUITING VEGETABLES	Eggplant, Groundcherry, Pepino, Pepper (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato
HERBS AND SPICES GROUP	Allspice, Angelica, Anise (anise seed and star), Annatto (seed), Balm (lemon balm), Basil, Borage, Burnet, Chamomile, Caper buds, Caraway, Caraway (black), Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese Chive, Cinnamon, Clary, Clove buds, Coriander (cilantro or Chinese parsley – leaf), Coriander (cilantroseed), Costmary, Cilantro (leaf and seed), Cumin, Curry (leaf), Dill (Dillweed and seed), Fennel (common, Florence), Fenugreek, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood
LEGUME VEGETABLES	Bean, Broad Bean, Chickpea, Guar, Jackbean, Lablab bean, Lentil, Pea, Pigeon Pea, Soybean, Sword bean
BRASSICA (COLE) LEAFY VEGETABLES	Amaranth, Arugula, Broccoli, Broccoli raab (rapini), Brussels Sprouts, Cabbage, Cauliflower, Cardoon, Cavalo broccoli, Celery, Chinese Broccoli (gai lon), Chinese Cabbage (bok choy, Napa), Chinese mustard Cabbage (gai choy), Chinese Celery, Celtuce, Chervil, Chrysanthemum (edible-leaved, Garland), Collards, Corn salad, Cress (garden, upland), Dandelion, Dock (sorrel), Endive (escarole), Fennel (florence), Kale, Kohlrabi, Lettuce (head and leaf), Mizuna, Mustard Greens, Mustard Spinach, Orach, Parsley, Purslane (garden, winter), Radicchio (red chicory), Rape Greens, Rhubarb, Spinach, Spinach (New Zealand, vine), Swiss Chard, Turnip Greens
OTHER	Asparagus, Avocado, Banana, Coffee, Cocoa, Cranberry, Fig, Globe artichoke, Grape, Hops, Kiwifruit, Mango, Mushroom, Okra, Olives, Papaya, Pawpaw, Peanut, Persimmon, Pineapple. Pomegranate, Strawberry, Tea, Water chestnut, Watercress, and all other food
POME FRUITS	Apple, Crabapple, Loquat, Mayhaw, Quince, Oriental Pear, or Pear
ROOT AND TUBER VEGETABLE	Arracacha, Arrowroot, Artichoke (Jerusalem, Chinese), Beet (garden, sugar), Burdock (edible), Canna (edible), Carrot, Cassava (bitter and sweet), Celeriac (celery root), Chayote (root), Chervil, (turnip-rooted), Chicory, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Oriental Radish (daikon), Parsley (turnip-rooted), Parsnip, Potato, Radish, Rutabaga, Salsify (oyster plant, black, Spanish), Skirret, Sweet potato, Tanier, Turmeric, Turnip, Yam bean (jicama, manoic pea), Yam (true)
STONE FRUIT	Apricot, Cherry (sweet and tart), Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot, Prune
TREE AND NUT	Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut, Hickory nut, Macadamia nut (bush nut), Pecan, Walnut (black and English), Pistachios
TROPICAL FRUITS	Papaya, Black Sapote, Canistel, Mamey Sapote, Mango, Sapodilla, Star Apple, Guava, Feijoa, Jaboticaba, Wax Jambu, Star Fruit, Passion Fruit, Acerola, Lychee, Longan, Spanish Lime, Rambutan, Pulasan, Sugar Apple, Atemoya, Custard Apple, Cherimoya, Ilama, Soursop, and Biriba.

#### GREENHOUSE

Use Sites	Comment
	Tomatoes, cucumbers, leaf and Bibb lettuces, bell peppers, eggplant, herbs, spices, and other vegetables cited above grown via greenhouse vegetable production.
	Brassicas such as broccoli and brussels sprouts, cucumbers, eggplants, melons, onion, peppers, squash are some of the most commonly grown vegetables for field transplant.

#### ORNAMENTAL (Herbaceous and Woody) Growing Outdoors, in Nurseries or in Greenhouses

Use Sites	Representative Commodities
ORNAMENTAL SHRUBS AND PLANTS	Amaranthus, Aster, Azalea, Ferns, Fuchsias, Caladium, Carnation, Chrysanthemum, Dahlia, Daisy, Lilies, Ivy, Ficus, Gardenia, Impatiens, Iris, Jasmine, Lilac, Marigold, Philodendron, Poinsettia, Rose, Zinnia
ORNAMENTAL TREES	Ash, Birch, Cedar, Cyprus, Dogwood, Fir, Elm, Juniper, Maple, Oak, Pine, Spruce

#### AGRICULTURAL USE RATES

Rates here indicated are for uses in agricultural and ornamental crops grown in the field or in greenhouses.

#### Stand-Alone:

- Dilute 1-2 pints of CAPTIVA PRIME per acre and apply by spraying every 4-7 days, when used as a preventative management tool or in an integrated pest management system.
- For hard to wet crops, consider using a spreader/sticker or adjuvant to enhance coverage of CAPTIVA PRIME to the crop.
- When tank mixing, read and carefully follow all applicable use directions, precautions, and limitations on the respective product labels. In interpreting all labels for the tank mix, the most restrictive situations must apply.
- · When pest pressure is heavier or canopy is dense, use higher rates and consider tank mixes with other insecticides or miticides.

#### Tank Mix:

- When tank mixed with other insecticides or miticides CAPTIVA PRIME will provide improved levels of control by increasing pest
  exposure to those products due to its irritant effect.
- Mix CAPTIVA PRIME at a rate of ½ to 1.5 pints per acre in water with other chemical or biological insecticides or miticides. Follow
  the use directions of all products used in the tank mix including use directions and additive/adjuvant requirements or constraints.

### **GREENHOUSE AND ORNAMENTAL USE RATES Stand Alone:**

- Dilute 1-2 pints of CAPTIVA PRIME in water and apply by spraying every 4 7 days when used as a preventative management tool or in an integrated pest management system.
- For hard to wet crops, consider using a spreader/sticker or adjuvant to enhance coverage of CAPTIVA PRIME to the crop.
- When tank mixing, read and carefully follow all applicable use directions, precautions and limitations on the respective product labels. In interpreting all labels for the tank mix, the most restrictive situations must apply.
- When pest pressure is heavier, use higher rates and consider tank mixes with other insecticides or miticides.

#### Tank Mix:

- When tank mixed with other insecticides or miticides CAPTIVA PRIME will provide improved levels of control increasing pest
  exposure to those products due to its irritant effect.
- Mix CAPTIVA PRIME at a rate of ½ to 1.5 pints in water with other chemical or biological insecticides or miticides
- When pest pressure is heavier, use higher rates.

#### STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container and keep closed. Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by State and local authorities by burning. If burned stay out of smoke. **Mini-Bulk Containers:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300. For other product information contact Gowan Company or see Material Safety Data Sheet

#### NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILTY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off

target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE FULLEST EXTENT PERMITTED BY LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

CAPTIVA® is a registered trademark of Gowan Company, L.L.C. Chemtrec® is a registered trademark of American Chemistry Council, Inc.

SUMBIT TO EPA (07-27-2016)

### CAPTIVA® PRIME

Alternate Brand Name: **GWN-10285 INSECT REPELLENT / INSECTICIDE** 

Active Ingredient	% By Wt.
Capsicum oleoresin extract*	7.60%
Garlic Oil	23.40%
Canola Oil	55.00%
Other Ingredients	14.00%
*contains 0.26% nominal concentration of Capsaicin and related capsaicinoids	Total: 100 00%

# KEEP OUT OF REACH OF CHILDREN CAUTION

	FIRST AID
If on skin or clothing	Take off contaminated clothing
	Rinse skin immediately with plenty of water for 15-20 minutes
	<ul> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
	HOT LINE NUMBER
	or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-

# PRECAUTIONARY STATEMENT HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, and chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear chemical-resistant gloves. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Ν	EΤ	C	O	N	T	ΕI	V	T	S				

EPA Reg No. 10163-XXX EPA Est No. (batch code XXX)





Produced For: Gowan Company, LLC. PO Box 5569 Yuma, AZ 85366-5569

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

#### Users should

- · Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **ENVIRONMENTAL HAZARDS**

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to industrial area below the mean high-water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all directions on the EPA approved label and any supplemental labeling

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read all Directions for Use carefully before applying.

#### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

#### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

#### **USE INFORMATION**

CAPTIVA PRIME is a repellent and insecticide formulated using concentrated plant extracts that can be used alone as a repellent or in tank mixes with other biological or conventional insecticides or miticides to improve the level of control.

The repellent effect of CAPTIVA PRIME protects the crops from pests population establishment by its antifeedant and anti–egg laying properties. The irritant effect of CAPTIVA PRIME increases pest mobility, exposure and vulnerability to insecticides or miticides that act on contact. CAPTIVA PRIME weakens the cuticle of some immature stages of insect and mite pests.

Read all directions before using this product. Apply CAPTIVA PRIME directly to any food or non-food crop growing in field and greenhouses up to and including the day of harvest, at a maximum rate of 4 pints per acre. Refer to the **Insecticidal and Miticidal Uses Section** for a complete listing of crops.

Read and follow these directions when using:

Do not allow spray drift from application site.

Use only with pressurized hand-held sprayers or spray trigger bottles.

Do not allow spray mixture to stand overnight for prolonged periods.

For best results use a spreader/sticker (adjuvant) to increase the efficacy.

CAPTIVA PRIME has been found to be compatible with most commonly used pesticides. To test for compatibility, mix a small amount of each product, in the appropriate proportions, in a small jar.

PHYTOTOXICITY: CAPTIVA PRIME has been evaluated for phytotoxicity on a wide range of crops and ornamentals. However, since testing on all varieties of all crops and ornamentals is not feasible, testing a small portion of the area to be treated for phytotoxicity is recommended before treating the entire area. Further, all possible combinations or sequences of pesticide sprays, including other fertilizers, surfactants, adjuvants and other pesticides, have not been tested, thus testing for phytotoxicity of spray mixtures is recommended. It is further recommended that spray equipment used to apply CAPTIVA PRIME be thoroughly cleaned before use. The addition of spray adjuvants enhances control in some crops under ideal conditions. Addition of certain adjuvants may cause phytotoxicity therefore, the addition of crop oils and other adjuvants should be thoroughly tested before using. The addition of crop oils to spray mixtures is NOT recommended on ornamental crops. Avoid tank mix combinations of CAPTIVA PRIME with compounds known to be incompatible with oil-based formulation to prevent phytotoxicity. 'Waxy bloom' on certain crops and ornamental plants may be reduced after a CAPTIVA PRIME application.

#### Insecticidal and Miticidal Uses

Use CAPTIVA PRIME as a foliar spray when pest pressure is low or use as a preventative management tool or integrated pest management system. When pest pressure is high use the product alone at its medium to high rates or in tank mix with other contact insecticides or miticides.

PREHARVEST INTERVAL: CAPTIVA PRIME has a zero day preharvest interval and can be applied up to and including the day of harvest.

TARGETED PESTS: Use CAPTIVA PRIME in a foliar spray to repel or suppress soft bodied pests.

- Mites (such as Tetranychus spp., Oligonychus spp., Panonychus spp., Polyphagotarsonemus latus, Phyllocoptruta oleivora, Brevipalpus phoenicis, Steneotarsonemus spinki)
- Thrips (such as Thrips spp., Frankliniella spp., Heliothrips spp., Scirtothrips citri.)
- Psyllids (such as Bactericera spp., Diaphorina citri, Cacopsylla spp.)
- Leafhoppers (such as Empoasca spp., Macrosteles spp., and Graphocephala spp.)
- Lepidoptera larvae (such as Spodoptera spp., Helicoverpa spp., Tuta absoluta, Thecla spp, Strymon basilides)
- White flies (such as Dialeurodes spp., Trialeurodes spp. Bemisia spp.)
- Other pests such as borers (Hypothenemus hampei), mealybugs (Planococcus spp., Pseudococcus spp., Dysmicoccus brevipes) and plant bugs (Leptoglossus spp., Nezara spp)

DIRECTIONS FOR USE ON INDOOR AND OUTDOOR VEGETABLES, FRUIT AND TREE NUTS, ORNAMENTAL FLOWERS, TREES, SHRUBS AND PLANTS GROWING IN CONTAINERS, INTERIOSCAPES, HOME AND GARDEN USE, LANDSCAPES, PARKS, AND RECREATIONAL AREAS

#### WHEN TO USE

For best results, apply CAPTIVA PRIME before populations reach damaging levels or when egg deposition is observed.

#### **BEFORE YOU USE**

Read and follow these directions when using: Do not allow spray to drift from application site. Use only with pressurized hand-held sprayers or spray trigger bottles. Do not allow spray mixture to stand overnight or for prolonged periods. For best results use a spreader/sticker (adjuvant) to increase the efficiency of CAPTIVA PRIME.

CAPTIVA PRIME has been found to be compatible with most commonly used pesticides. To test for compatibility, mix a small amount of each product, in the appropriate proportions, in a small jar.

Some pesticides can cause phytotoxic effects ranging from slight burning or browning of leaves to distorted leaves, fruit, flowers or stems. Damage symptoms can vary with the type of plant that has been treated. It is impossible to test all plant species for phytotoxicity. To assure that the plants to be treated are not sensitive to the treatment, apply a small amount of the product to a few leaves or the above ground portion of the plant and check back in 2-4 days for signs of phytotoxicity.

#### HOW TO USE FOR HAND-HELD SPRAYERS AND SPRAY TRIGGER BOTTLES

Fill sprayer or bottle with appropriate amount of water and concentrate. Mix the spray solution thoroughly. Keep the spray solution agitated during application.

#### HOW TO USE FOR HOSE-END SPRAYERS

Follow hose-end sprayer instructions to determine how to fill, set dial, clean and disconnect from hose. Set dial on sprayer to deliver rate as directed below.

#### HOW MUCH TO USE FOR ALL APPLICATIONS

1/2 - 1.44 teaspoons of CAPTIVA PRIME per gallon of water, 1 gallon of mixture will cover approximately 350 sq. ft.

#### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in a cool dry place inaccessible to children.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency (800)858-7378 (National Pesticide Information Center) for disposal instructions. Never place unused product down an indoor or outdoor drain.

### FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300. For other product information contact Gowan Company or see Material Safety Data Sheet

#### NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILTY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE FULLEST EXTENT PERMITTED BY LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

CAPTIVA® is a registered trademark of Gowan Company, L.L.C. Chemtrec® is a registered trademark of American Chemistry Council, Inc.