



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 31, 2017

Ms. Kyla S. Smith
Registration Specialist
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Subject: PRIA Label Amendment – Amendment of Existing Tolerance in/on Hop, Dried Cones; Associated with PP #6F8489
Product Name: Onager® EW Miticide
EPA Registration Number: 10163-337
Application Date: June 16, 2016
Decision Numbers: 519000 & 519001

Dear Ms. Smith:

The application referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable under FIFRA Section 3(c)(7), subject to the following conditions:

1. You must submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.
2. You are required to comply with the data requirements described in the DCI identified below:
 - a. Hexythiazox GDCI-128849-879

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:

<http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Be aware that proposed data requirements have been identified in a Work Plan. For more information on these proposed data requirements, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:
http://www.epa.gov/oppsrrd1/contacts_prd.htm

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). If you have any questions, please contact Mr. Carmen J. Rodia, Jr. by phone at (703) 306-0327, or via email at Rodia.Carmen@epa.gov.

Sincerely,



Richard Gebken
Product Manager 10
Invertebrate & Vertebrate Branch 2
Registration Division (7505P)
Office of Pesticide Programs

*Enclosures: Stamped "Accepted" Master & Supplemental Labels, dated October 30, 2017
E-Copy of Published Hexythiazox FR for Tolerances, dated October 30, 2017
EFED Risk Assessment, dated May 25, 2017
HED Risk Assessment, dated July 5, 2017*

ONAGER EW MITICIDE

[Alternate Brand Name: ONAGER OPTEK MITICIDE]

EMULSION FOR AGRICULTURAL USE ONLY

| | |
|---|-----------------|
| ACTIVE INGREDIENT: Hexythiazox | % By Wt. |
| <i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide..... | 11.93% |
| OTHER INGREDIENTS | 88.07% |
| TOTAL | 100.00% |


Onager EW EW contains 1.0 lb. active ingredient per gallon
Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

| FIRST AID | |
|--|---|
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person. |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-478-0798, 24 hours per day, 7 days per week. | |
| NOTE TO PHYSICIAN | |
| Contains petroleum distillate. May pose an aspiration pneumonia hazard. | |

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash clothing before reuse.



NET CONTENTS _____

Gowan®
The Go To Company

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

EPA Reg. No. 10163-337
EPA Est. No. 67545-AZ-01

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, nitrile rubber ≥ 14mils, neoprene rubber ≥ 14mils, or viton ≥ 14mils
- Shoes plus socks

Remove and wash contaminated clothing before reuse. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

PRODUCT INFORMATION

Onager EW Miticide is recommended for use as a foliar application in labeled crops. Onager EW Miticide provides effective control of Tetranychid spider mite species. It is an emulsion to be mixed with water and applied as a spray.

Onager EW controls mites through activity on eggs and immature stages. Control is achieved from either direct contact with the spray or from contact with treated plant surfaces. Through its ovicidal activity, Onager EW controls newly deposited mite eggs and eggs which are laid after application. Onager EW is also highly effective in controlling immature motile stages of target mite species that are sprayed or move onto treated surfaces. Adult mites are not directly affected. However, eggs produced by females in contact with treated surfaces will be rendered nonviable. Onager EW provides residual control of pest mite species. The degree and duration of control is dependent on the rate used, growth stage of the mite, species of mite, and climatic conditions under which the material is applied.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all directions on the EPA approved label and any supplemental labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves made of barrier laminate, nitrile rubber ≥ 14mils, neoprene rubber ≥ 14mils, or viton ≥ 14mils
- Shoes plus socks

Managing Mite Resistance: Repeated use of miticides with similar modes of action may lead to buildup of resistant strains of mites. For resistance management, apply Onager EW only once per year. To reduce the potential for developing mite resistance, do not apply a miticide with a similar mode of action more than once per year (Onager EW and Apollo® should not be applied to the same crop in any one year). If additional miticide applications are needed, use a product with a different mode of action, such as a contact adulticide. Read and follow product label before applying any miticide.

Mixing Directions:

- Turn on spray tank agitation. Pour recommended amount of this material on the surface of water in a nearly filled spray tank.
- Add balance of water to fill the tank.
- Keep agitator running during filling and spraying operations. Do not allow mixture to stand.

Compatibility: Onager EW is compatible with most agricultural chemicals. However, all possible combinations have not been evaluated. Before full-scale mixing of Onager EW with other products, test mix small proportionate quantities of each to ensure compatibility. Do not mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

Phytotoxicity: As is common with most emulsion formulations, adverse effects such as spotting or discoloration of the treated surfaces can occur. Some conditions known to contribute to phytotoxicity include, but are not limited to, high temperatures, poor spray drying conditions, excessive spray deposit or run-off, certain spray mixtures, stage of crop development or tank mixes with other pesticides.

Application Information: Use sufficient spray volume to obtain thorough, uniform coverage of all plant surfaces. Apply using ground equipment, chemigation or air unless otherwise specified under the crop comments or on supplemental labeling supplied by Gowan Company.

Important: **Onager EW Miticide is a 1 pound active ingredient per gallon** emulsifiable concentrate formulation to be diluted with water for application in commercial plantings only. Do not use in home plantings.

Use Recommendations: For best results, apply Onager EW Miticide at first sign of mites before adult mite build up. Onager EW Miticide will not control adult spider mites. Use higher rates on moderate-to-high mite infestations, large trees with dense foliage, or for extended residual control. The lowest rate of Onager EW Miticide may be used in conjunction with other IPM strategies, or for shorter residual control later in the season. The use of less than label rates is not recommended since it will result in poor performance and contribute to resistance development. If adult mites are present in medium-to-high populations, better results may be obtained using Onager EW Miticide in combination with a registered contact adulticide. Onager EW Miticide may be tank mixed with horticultural grade oil. Refer to all precautions regarding phytotoxicity on the oil label.

Do not make more than one application of Onager EW Miticide per year.

Chemigation Systems: Onager EW may be applied through irrigation systems (chemigation) to any crop on the Onager EW label unless otherwise specified under the crop comments or on supplemental labeling supplied by Gowan Company. Do not allow chemigation to run off field.

Types of Irrigation Systems: Apply Onager EW only through sprinkler, including center pivot, lateral move, Low Energy Precision Applications (LEPA), end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply Onager EW through any other type of irrigation system.

GENERAL DIRECTIONS FOR ALL RECOMMENDED TYPES OF IRRIGATION SYSTEMS

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. For best results apply at 100% input/travel speed, for center pivots or 0.10 inch (2.716 gallons) up to 0.15 inch (4,073 gallons) of water/A, for other systems. Higher labeled rates of Onager EW may be necessary for chemigation applications.

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift: Do not apply when wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump; such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems: Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regular serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Cleaning the Chemical Injection System: In order to accurately apply pesticides, the chemical injection system must be kept clean; free of chemical or fertilizer residues and sediments. Refer to your owner's manual or ask your equipment supplier for the cleaning procedure for your injection system.

Flushing the Irrigation System: At the end of the application period, allow time for all lines to flush the pesticide through all nozzles or emitters before turning off irrigation water. To ensure the lines are flushed and free of pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

Equipment Area Contamination Prevention

It is recommended that nozzles in the immediate area of control panels, chemical supply tanks, pumps and system safety devices be plugged to prevent chemical contamination of these areas.

Center-Pivot and Automatic-Move Linear Systems: Inject the specified dosage per acre continuously for one complete revolution or move of the system. DO NOT USE END GUNS. The system should be run at maximum speed.

Solid Set and Manually Controlled Linear Systems: Injection should be during the last 30 to 60 minutes of regular irrigation period or as a separate 30 to 60 minute application not associated with a regular irrigation. DO NOT USE END GUNS.

PREHARVEST INTERVAL

The required days between the last application and harvest are given in () after each crop name.

APPLICATION DIRECTIONS

| CROP | PEST | RATE OZS./ACRE | COMMENTS |
|--|---|-------------------|--|
| ALFALFA (14) (WESTERN US <u>ONLY</u> *) | Two-spotted spider mite | 10 - 24 | Ground Application: Apply the recommended rate of Onager EW by ground equipment at 15 – 20 GPA. Air Application: Apply the recommended rate of Onager EW by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not apply more than a total of 24 ozs. formulated product per acre per year. Do not make more than one application per calendar year. *May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281 | | |
| TIMOTHY (14) (West of the Rockies) | McDaniel spider mite, Two-spotted spider mite, Banks grass mite | 10 - 24 | Ground Application: Apply the recommended rate of Onager EW by ground equipment at 15 – 20 GPA. Air Application: Apply the recommended rate of Onager EW by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not apply within 14 days of harvest for hay. Animals may graze treated fields on the day of application. Do not apply more than a total of 24 ozs. formulated product per acre per calendar year. Do not make more than one application per calendar year. | | |
| BEANS, DRY AND SUCCULENT (14) (WESTERN US <u>ONLY</u> *) | Two-spotted spider mite, Pacific spider mite, Strawberry spider mite | 10 - 24 | Ground Application: Apply a minimum of 20 gallons finished spray per acre. Air application: apply a minimum of 10 gallons finished spray per acre. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not apply more than a total of 24 ozs. of formulated product per acre per calendar year. Do not make more than one application per calendar year. Do not harvest or graze bean vine forage or hay. *May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281 | | |
| BERMUDAGRASS (7) (West of the Rockies) | Banks grass mite, Two-spotted spider mite | 12 – 24 | Apply at the first sign of mites before populations begin to build. Ground Application: Apply the recommended rate of Onager EW by ground equipment at 15 – 20 GPA. Air Application: Apply the recommended rate of Onager EW by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems on Section 3 label. |
| | <ul style="list-style-type: none"> Do not apply more than a total of 24 ozs. formulated product per acre per calendar year. Do not make more than one application per calendar year. | | |

| | | | |
|--|--|---------|---|
| CITRUS (7) (CA, AZ AND TX ONLY) | Tetranychid Mites (such as European Red Mite and Two-spotted Spider mite) | 12 - 24 | Application by ground equipment only. Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre. |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. Do not apply more than 24 ozs. per acre per calendar year. | | |
| COTTON (35) | Two-spotted spider mite, Pacific spider mite, Strawberry spider mite, Carmine spider mite | 12 - 20 | Apply Onager EW before boll opening and prior to adult mite buildup. Complete coverage of foliage is essential to optimal performance. Ground Application: Apply at recommended rates Air application: Use recommended per acre rate and apply a minimum of 20 gallons of finished spray per acre. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not apply more than a total of 20 ozs. formulated product per acre per calendar year. Do not make more than one application per calendar year. | | |
| CORN, FIELD (30) | Banks grass mite, Two-spotted spider mite, Pacific spider mite, Strawberry spider mite | 6 - 24 | Ground Application: Apply the recommended rate of Onager EW with conventional spray boom equipment at 15 – 20 GPA. Thorough and uniform coverage of plants is required for optimal results. Air Application: Apply the recommended rate of Onager EW by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not apply more than a total of 24 ozs. formulated product per acre per calendar year. Do not make more than one application per calendar year. Do not harvest field corn forage, stover, and grain for 30 days after application. | | |
| CORN, SWEET (including ears and forage) (28) (WESTERN US ONLY*) | Banks grass mite, Two-spotted spider mite, Pacific Spider Mite, Strawberry spider mite | 10 - 24 | Ground Application: Apply the recommended rate of Onager EW with conventional spray boom equipment at a minimum of 20 gallons finished spray per acre. Thorough and uniform coverage of plants is required for optimal results. Air Application: Apply the recommended rate of Onager EW by air in a minimum of 10 gallons finished spray per acre. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not apply more than a total of 24 ozs. formulated product per acre per calendar year. Do not make more than one application per calendar year. Do not apply within 28 days of harvest of sweet corn or forage. <p>*May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281</p> | | |
| CHRISTMAS TREE, CHRISTMAS TREE PLANTATIONS | Pacific Spider Mite, Spruce Spider Mite, Two-spotted Spider Mite | 12 - 24 | Ground Application: Apply at recommended rates Air application: Apply in 2-10 gallons finished spray per acre. For best coverage, apply in two opposing passes of 5 gallons each. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not make more than one application per year. | | |
| SMALL FRUIT VINE CLIMBING SUBGROUP, EXCEPT FUZZY KIWIFRUIT 13-07F (7) Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these | Tetranychid Mites (such as European Red Mite and Two-spotted Spider Mite), Pacific spider mite, Willamette mite | 12 - 24 | Ground Application: Apply at recommended rates. Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre. Air Application: apply a minimum of 10 gallons finished spray per acre. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. Do not apply more than 24 ozs. per acre per calendar year. | | |
| GREENHOUSE TOMATOES (1) | Tetranychid Mites (such as European Red Mite and Two-spotted Spider Mite) | 12 - 24 | Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre. |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. Do not apply more than 24 ozs. per acre per calendar year. | | |

| | | | |
|--|---|---------|---|
| HOPS (30) | Tetranychid Mites (such as Two-spotted spider mite) | 12 - 24 | Ground Application: Apply a minimum of 20 gallons of finished spray per acre. Air Application: Use sufficient volume of water to obtain complete coverage but not less than 10 gallons per acre. |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. Do not apply more than 24 ozs. per acre per calendar year. | | |
| NON-BEARING TREE FRUITS and NUTS, CITRUS, and VINES (FIELD GROWN, NURSERY, GREENHOUSE, SHADEHOUSE) <i>Non-bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application</i> | Citrus Red Mite, European Red Mite, Two-spotted Spider Mite, McDaniel Spider Mite, Pacific Spider Mite, Sixspotted Mite, Texas Citrus Mite | 12 - 24 | Ground Application: Apply at recommended rates Air Application: apply a minimum of 5 gallons finished spray per acre. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. | | |
| PEPPER/EGGPLANT SUBGROUP 8-10B (1) African eggplant; bell pepper; eggplant; martynia; nonbell pepper; okra; pea eggplant; pepino; roselle; scarlet eggplant; cultivars, varieties, and/or hybrids of these | Tetranychid Mites (such as Two-spotted spider mite), Pacific spider mite, Strawberry spider mite | 12 - 24 | Ground Application: Apply a minimum of 20 gallons of finished spray per acre. Air Application: apply a minimum of 20 gallons finished spray per acre. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. Do not apply more than 24 ozs. per acre per calendar year. | | |
| POME FRUIT GROUP 11-10 (28) Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these | European red mite, Two-spotted spider mite, McDaniel spider mite, Pacific spider mite, Willamette mite | 12 - 24 | Ground Application: Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre. Air Application: Use sufficient volume of water to obtain complete coverage but not less than 10 gallons per acre. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. For D'Anjou or Bosc pear varieties, use higher labeled rates (16-24 ozs./acre). | | |
| POTATOES (21) (WESTERN US <u>ONLY</u> *) | Two-spotted Spider Mites; Potato Psyllid | 12 - 24 | Ground Application: Apply at recommended rates Air Application: Apply a minimum of 5 gallons finished spray per acre. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. Do not apply more than a total of 24 ozs. formulated product per acre per calendar year. <p>*May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281</p> | | |
| SORGHUM (30) (Western US only, including Oklahoma and Texas*) | Banks grass mite, Two-spotted spider mite | 10 – 24 | Ground Application: Apply the recommended rate of Onager EW by ground equipment at 15 – 20 GPA. Air Application: Apply the recommended rate of Onager EW by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems |
| | <ul style="list-style-type: none"> Do not apply more than a total of 24 ozs. formulated product per acre per calendar year. Do not make more than one application per calendar year. <p>*May only be applied in the Western US - west of the line defined as follows: KS, NE, SD, ND: W of Rt. 281 Entire States of OK and TX included for use</p> | | |
| STONE FRUITS GROUP (7) (including Plums) | European red mite, Two-spotted spider mite, McDaniel spider mite, Strawberry spider mite, Pacific spider mite, Pecan leaf scorch mite, Willamette mite | 12 - 24 | Ground Application: Apply a minimum of 20 gallons finished spray per acre. Air Application: Apply a minimum of 10 gallons finished spray per acre. Chemigation: see details under Chemigation Systems. |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. | | |

| | | | |
|---|---|---------|---|
| SUGAR BEETS (45) (WESTERN US <u>ONLY</u> *) | Banks grass mite, Two-spotted spider mite | 12 – 24 | <p>Apply at the first sign of mites before populations begin to build.</p> <p>Ground Application: Apply the recommended rate of Onager EW by ground equipment at 15 – 20 GPA.</p> <p>Air Application: Apply the recommended rate of Onager EW by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.</p> <p>Chemigation: see details under Chemigation Systems on Section 3 label.</p> |
| | <ul style="list-style-type: none"> Do not apply more than a total of 24 ozs. formulated product per acre per calendar year. Do not make more than one application per calendar year. <p>*May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281</p> | | |
| TREE NUTS GROUP AND PISTACHIOS (7) | European red mite, Two-spotted spider mite, McDaniel spider mite, Strawberry spider mite, Pacific spider mite, Pecan leaf scorch mite, Willamette mite, Citrus Red Mite | 12 - 24 | <p>Ground Application: Apply a minimum of 20 gallons finished spray per acre.</p> <p>Air Application: Apply a minimum of 5 gallons finished spray per acre.</p> <p>Chemigation: see details under Chemigation Systems.</p> |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. | | |
| WHEAT, including emmer wheat and triticale (14) (West of the Rockies) | Banks grass mite | 10 - 24 | <p>Apply at the first sign of mites before populations begin to build.</p> <p>Ground Application: Apply the recommended rate of Onager EW by ground equipment at 15 – 20 GPA.</p> <p>Air Application: Apply the recommended rate of Onager EW by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.</p> <p>Chemigation: see details under Chemigation Systems.</p> |
| | <ul style="list-style-type: none"> Do not make more than one application per calendar year. Do not apply after the boot stage of growth. Do not harvest hay or forage for at least 14 days after application. Do not apply more than a total of 24 ozs. formulated product per acre per calendar year. | | |

RESTRICTIONS ON ALL CROPS

- Do not graze or feed livestock on cover crops growing in treated areas.
- Onager EW is an emulsion to be diluted with water for application in commercial plantings only.
- Apply Onager EW prior to adult mite build up. Onager EW will not control adult spider mites. Use higher rates on moderate-to-high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium-to-high populations, better results may be obtained using Onager EW in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager EW.
- Do not make more than one application of Onager EW or any other Hexythiazox product to the same crop per year.
- Do not plant rotational crops other than those on this label within 120 days of this application.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container. Do not use or store this product near heat or open flame. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, and away from other pesticides, fertilizers, food, or feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

FOR BULK AND MINI-BULK CONTAINERS

CONTAINER DISPOSAL: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

CONTAINER PRECAUTIONS: Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices. REFILL ONLY WITH ONAGER EW. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than Onager EW will result in contamination and may weaken container. After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.

For other product information, contact Gowan Company or see Material Safety Data Sheet.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our recommendations for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. All such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Chemtrec® is a registered trademark of American Chemistry Council Inc.

Onager EW® and the donkey logo are registered trademarks of Gowan Company, L.L.C.

Apollo® is a registered trademark of Irvita Plant Protection N.V.

EPA Text Pending: Onager EW (to EPA 10-26-17)

| | | |
|-------|-----|-----------|
| GROUP | 10A | ACARICIDE |
|-------|-----|-----------|

This supplemental label expires on October 30, 2020, and must not be used or distributed after this date.

ONAGER EW MITICIDE

(Alternate Brand Name: Onager Optek Miticide)

EMULSION FOR AGRICULTURAL USE ONLY

| | |
|---|-----------------|
| ACTIVE INGREDIENT: Hexythiazox | % By Wt. |
| <i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide..... | 11.93% |
| OTHER INGREDIENTS | 88.07% |
| TOTAL | 100.00% |

Onager EW contains 1.0 lb. active ingredient per gallon
Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE


- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

APPLICATION DIRECTIONS

| CROP | PEST | RATE Oz./acre | COMMENTS |
|------|--|------------------|---|
| HOPS | Two-spotted spider mite | 12 - 24 | For application by ground equipment only. Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre. |
| | <ul style="list-style-type: none"> • Do not make more than one application per year. • Do not apply more than 24 ounces of formulated product per acre per calendar year. • PHI = 30 days | | |

RESTRICTIONS ON ALL CROPS

- Do not graze or feed livestock on cover crops growing in treated areas.
- Onager EW is an emulsifiable concentrate to be diluted with water for application in commercial plantings only.
- Apply Onager EW prior to adult mite build up. Onager EW will not control adult spider mites. Use higher rates on moderate-to-high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium-to-high populations, better results may be obtained using Onager EW in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager EW.
- Do not make more than one application of Onager EW or any other Hexythiazox product to the same crop per year.
- Do not plant rotational crops other than those on this label within 120 days of this application.



NET CONTENTS ____

Gowan®
The Go To Company

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

EPA Reg. No. 10163-337
EPA Est. No.

EPA Text: Onager EW (To EPA 10-26-17)