



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 9, 2022

Kyla Smith
Registration Specialist
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

Subject: Registration Review Label Mitigation for Benfluralin and Trifluralin
Product Name: TEAM 2G
EPA Registration Number: 10163-340
Application Dates: 3/7/22 and 6/11/19
Decision Numbers: 582361 and 582360

Dear Ms. Smith:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Benfluralin and Trifluralin Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2
EPA Reg. No. 10163-340
Decision No. 582361 and 582360

If you have any questions about this letter, please contact DeMariah Koger by phone at (202)-566-2288, or via email at koger.demariah@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington".

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Program

Enclosure

Team[®] 2G

Specialty Herbicide

A selective preemergence herbicide for control of crabgrass and most other annual weed grasses on turfgrass and golf courses in established:

Bahiagrass

Bermudagrass

Bluegrass (perennial)

Centipedegrass

Fescue

Ryegrass (perennial)

St. Augustinegrass

Zoysiagrass

Bentgrass fairways

and other high cut

bentgrass areas

and/or

Controls [tough] weeds [such as annual bluegrass] [smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [prostrate spurge] [spotted spurge] [creeping spurge] [creeping woodsorrel] [yellow woodsorrel]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds [like spurge] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a spurge preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [spurge]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees and shrubs

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

ACCEPTED

Mar 09, 2022

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 10163-340

10163-340

NET CONTENTS _____

EPA Reg. No. 10163-340
EPA Est. No.
85366

Gowan
The Go To Company

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ

Active Ingredients:

Benfluralin: <i>N</i> -butyl- <i>N</i> - ethyl- α,α,α -trifluoro-2, 6-dinitro- <i>p</i> -toluidine	1.33%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine	0.67%
Other Ingredients	98.00%
Total	100.00%

Contains 1 lb active ingredient per 50 lb bag.

Keep Out of Reach of Children CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-478-0798 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Harmful If Swallowed Or Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reaction In Some Individuals
 Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
 - Shoes plus socks
 - Chemical-resistant gloves made of any waterproof material
- Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
 - Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Non-Target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. This product must be watered in as soon as possible after application.

Entry Restrictions: Do not enter or allow other people (or pets) to enter the treated area until dusts have settled. Except for personnel involved in the watering-in, do not enter or allow other people or pets to enter the treated area until the watering-in is complete and the surface is dry.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WEED RESISTANCE MANAGEMENT

For resistance management, please note that Team 2G is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to Team 2G and Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

Best Management Practices

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Team 2G or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - (2) a spreading patch of non-controlled plants of a particular weed species;
 - (3) surviving plants mixed with controlled individuals of the same species.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Gowan Company representative.

GENERAL INFORMATION

Team® 2G specialty herbicide is a preemergence herbicide for application to established turfgrass for control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass), and green and yellow foxtail. Team 2G also provides partial control of prostrate spurge (*Euphorbia supina*), spotted spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*), creeping woodsorrel (*Oxalis corniculata*), and yellow woodsorrel (*Oxalis stricta*). Team 2G controls susceptible annual weeds by killing their seeds as they germinate. Apply Team 2G prior to germination of target weeds since it does not control established weeds. Team 2G will not harm most nearby established ornamental trees and shrubs.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

- Do not apply this product to plants that will bear fruits, berries, or nuts within 12 months of application.
- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not make more than 2 applications of Team 2G per year.
- Do not apply Team 2G in the spring to turfgrass planted the previous fall.
- Team 2G may thin established annual bluegrass (*Poa annua*) turfgrass and fine-leaved fescues at rates above 100 lb per acre (2 lb active ingredient).
- In Bermudagrass areas that have been overseeded with winter grasses, a spring application of Team 2G may thin the overseeded grasses.
- Do not aerially apply this product.
- **Maximum Single Application Rate:** Cool season turfgrass – 120 lb of Team 2G per acre; warm season turfgrass – 150 lb of Team 2G per acre.
- **Maximum Yearly Rate:** Cool season turfgrass – 240 lb of Team 2G per acre; warm season turfgrass – 300 lb of Team 2G per acre.
- **Minimum Re-Treatment Interval:** Cool season turfgrass – 56 to 70 days; warm season turfgrass – 70 to 84 days.
- This product must be watered in as soon as possible after application.
- This product must be soil-incorporated within 24 hours of application either by mechanical incorporation or at least 0.5 acre inch of water (rainfall, irrigation). If a second incorporation is needed it should be delayed 3 to 5 days after the first incorporation and be completed prior to planting.

Reseeding

Do not reseed treated areas for at least 8 weeks after a single application of 75 to 100 lb of product (1.5 to 2 lb active ingredient) per acre unless injury can be tolerated, or 12 to 16 weeks after a single application of 100 to 150 lb of product (2 to 3 lb active ingredient) per acre unless injury can be tolerated. When reseeding, follow proper cultural practices such as soil cultivation, irrigation, and fertilization. For satisfactory reseeding results after using Team 2G, increase the amount of seed used and use equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder).

APPLICATION DIRECTIONS

Apply Team 2G using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

Application Techniques for Applying Team 2G

- When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treated smaller areas, a small rotary applicator is recommended. To insure proper coverage and avoid missed areas, early morning treatment when wheel tracks are evident in the dew and the wind is calm is recommended.
- When using tractor mounted or pull-type rotary applicators, early morning applications when the wind is calm and dew is on the ground is recommended. Wheel tracks are evident in the dew and granules can be spread uniformly.

Spreader Settings as a Guide for Calibration

Note: These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by seller or the spreader manufacturer.

Speed (mph)	Settings for Spreader		
	Recommended Rate of Team 2G		
	75 lb/acre	100 lb/acre	150 lb/acre
XX	XX	XX	XX

[Editor's Note: When Team 2G is formulated on limestone granules rather than clay granules, the above table should be replaced with the following Spreader Setting table.]

Spreader Type	Settings for Spreader		
	Recommended Rate of Team 2G		
	1.75 lb/1000 sq ft	2.25 lb/1000 sq ft	3.5 lb/1000 sq ft
XX	XX	XX	XX

Rates, Frequency, and Timing of Application

Application Rate Reference Chart

Application Rate of Team 2G	
(lb/acre)	(lb/1000 sq ft)
75	1 3/4
100	2 1/4
150	3 1/2

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply Team 2G approximately 1 to 2 weeks prior to the germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

Spring Germinating Summer Grasses (Crabgrasses, Goosegrasses, Foxtails, Barnyardgrass)

- **Northern Cool Season Turfgrass:** For preemergence control, apply 75 to 120 lb of Team 2G per acre. Apply in the spring, 1 to 2 weeks prior to the onset of conditions favorable for annual grass weed seed germination. Where heavy goosegrass pressure exists, use the rate of 120 lb of Team 2G per acre. If split applications are desired, apply a cumulative rate of 240 lb of Team 2G per acre and make the second application 8 to 10 weeks after the first.
- **Southern Warm Season Turfgrass:** For preemergence control, apply 100 to 150 lb of Team 2G per acre. Apply in the late winter or early spring, 1 to 2 weeks prior to the onset of conditions favorable for annual grass weed seed germination. In areas where germination of the annual weed grass species continues for extended periods of time, make a second application of Team 2G at the recommended rate 10 to 12 weeks after the first application for continued control.

Poa annua (Annual Bluegrass)

Eliminating annual bluegrass in areas of heavy infestation will temporarily result in thin turfgrass areas. Proper fertilization, irrigation, and soil incorporated reseeding will encourage existing desirable turfgrass and newly planted seedling turfgrass to fill in thin areas previously occupied by annual bluegrass.

- **Northern Cool Season Turfgrass:** For preemergence control, apply 75 to 100 lb of Team 2G per acre in late summer or early fall and again in late winter or spring, 1 to 2 weeks prior to the expected germination of annual bluegrass.
- **Southern Warm Season Turfgrass:** For preemergence control, apply 150 lb of Team 2G per acre in the late summer or early fall, 1 to 2 weeks prior to the expected germination of annual bluegrass. In southern areas where germination of annual bluegrass continues for extended periods of time, a second application of Team 2G at 150 lb per acre can be made 10 to 12 weeks after the initial treatment.

(Storage and Disposal for rigid containers 5 gal or less)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into a mix tank and drain for 10 seconds. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into a mix tank and continue to drain for 10 seconds. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrigid containers any size)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for refillable rigid containers greater than 5 gal)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into a mix tank and continue to drain for 10 seconds. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.
For other product information, contact Gowan Company or see Material Safety Data Sheet.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. All such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Team® is a registered trademark of Gowan Company LLC.
All other brands are registered trademarks of their respective owners.

Team 2G (to EPA 6-12-19, 2-21-22)