



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 25, 2018

Kyla S. Smith
Registration Specialist
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

Subject: Label Amendment – Label updates including PRN 2017-1 and 2017-2
Product Name: Turf Fertilizer Contains Gallery Plus Team
EPA Registration Number: 10163-344
Application Date: November 20, 2017
Decision Number: 537055

Dear Ms. Smith:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 10163-344
Decision No. 537055

with FIFRA section 6. If you have any questions, please contact Emily Schmid at 703-347-0189 or by email at schmid.emily@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long, sweeping horizontal line extending from the end of the name.

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

Turf Fertilizer - Contains Gallery® plus Team®

EPA Reg. No. 10163-344


The master label for this product consists of separate, self-contained sub labels for professional use and residential use, respectively.

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

Turf Fertilizer - Contains Gallery® plus Team®

EPA Reg. No. 10163-344

[Sub Label A: Professional Uses]



(Base label for rigid containers 5 gal or less):

Turf Fertilizer - Contains Gallery® plus Team®

Herbicide

A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established turfgrass including golf courses, parks, cemeteries, and non-residential and residential turf.

This product is intended for use by persons licensed by the state for herbicide application only.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristlegass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

- and/or
- For use on all turfgrass types [and/or] including bentgrass**
- and/or
- For use on cool season and warm season turfgrass**
- and/or
- Just one application needed**
- and/or
- More than a crabgrass preventer [also a barnyardgrass preventer]**
- and/or
- Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**
- and/or
- Labeled for over 40 different [grassy] [and] [broadleaf] weeds as listed**

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	0.29%
trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N, N</i> -dipropyl- <i>p</i> -toluidine	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α, α, α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.53%
Other Ingredients	98.91%
Total	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor’s note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N)	XX%
Ammoniacal Nitrogen	XX%
Urea Nitrogen	XX%
Available Phosphoric Acid (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%
Calcium	XX%
Magnesium (Mg)	XX%
Copper (Cu)	XX%
Iron (Fe)	XX%
Manganese (Mn)	XX%
Zinc (Zn)	XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/Personal Protective Equipment (PPE) immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. Contact 1-888-478-0798 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

GroundWater Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-888-478-0798.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 10163-344

EPA Est. _____

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

**Produced for
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569**

Net Weight _____

(Base label for nonrigid containers any size):

Turf Fertilizer - Contains Gallery[®] plus Team[®]

Herbicide

A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established turfgrass including golf courses, parks, cemeteries, and non-residential and residential turf.

This product is intended for use by persons licensed by the state for herbicide application only.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

Labeled for over 40 different [grassy] [and] [broadleaf] weeds as listed

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]- 2,6-dimethoxybenzamide and isomers	0.29%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.53%
Other Ingredients	98.91%
Total	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N)	XX%
Ammoniacal Nitrogen	XX%
Urea Nitrogen	XX%
Available Phosphoric Acid (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%
Calcium	XX%
Magnesium (Mg)	XX%
Copper (Cu)	XX%
Iron (Fe)	XX%
Manganese (Mn)	XX%
Zinc (Zn)	XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/ Personal Protective Equipment (PPE) immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. Contact 1-888-478-0798 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

GroundWater Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-888-478-0798.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 10163-344

EPA Est. _____

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

Produced for
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Net Weight _____

(Base label for refillable rigid containers greater than 5 gal):

Turf Fertilizer - Contains Gallery[®] plus Team[®]

Herbicide

A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established turfgrass including golf courses, parks, cemeteries, and non-residential and residential turf.

This product is intended for use by persons licensed by the state for herbicide application only.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

Labeled for over 40 different [grassy] [and] [broadleaf] weeds as listed

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	0.29%
trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N, N</i> -dipropyl- <i>p</i> -toluidine	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α, α, α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.53%
Other Ingredients	98.91%
Total	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N)	XX%
Ammoniacal Nitrogen	XX%
Urea Nitrogen	XX%
Available Phosphoric Acid (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%
Calcium	XX%
Magnesium (Mg)	XX%
Copper (Cu)	XX%
Iron (Fe)	XX%
Manganese (Mn)	XX%
Zinc (Zn)	XX%

Keep Out of Reach of Children

CAUTION**Precautionary Statements****Hazards to Humans and Domestic Animals**

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/ Personal Protective Equipment (PPE) immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. Contact 1-888-478-0798 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

GroundWater Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-888-478-0798.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 10163-344

EPA Est. _____

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

Produced for
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Net Weight _____

(Base label for nonrefillable rigid containers larger than 5 gal):

Turf Fertilizer - Contains Gallery[®] plus Team[®]

Herbicide

A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established turfgrass including golf courses, parks, cemeteries, and non-residential and residential turf.

This product is intended for use by persons licensed by the state for herbicide application only.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

Labeled for over 40 different [grassy] [and] [broadleaf] weeds as listed

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	0.29%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.53%
Other Ingredients	98.91%
Total	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N)	XX%
Ammoniacal Nitrogen	XX%
Urea Nitrogen	XX%
Available Phosphoric Acid (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%
Calcium	XX%
Magnesium (Mg)	XX%
Copper (Cu)	XX%
Iron (Fe)	XX%
Manganese (Mn)	XX%
Zinc (Zn)	XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/ Personal Protective Equipment (PPE) immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. Contact 1-888-478-0798 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

GroundWater Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-888-478-0798.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 10163-344

EPA Est. _____

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

**Produced for
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569**

Net Weight _____

(Label booklet cover):

ISOXABEN	GROUP	21	HERBICIDE
TRIFLURALIN	GROUP	3	HERBICIDE
BENEFIN	GROUP	3	HERBICIDE

Turf Fertilizer - Contains Gallery® plus Team®

Herbicide

A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established turfgrass including golf courses, parks, cemeteries, and non-residential and residential turf.

This product is intended for use by persons licensed by the state for herbicide application only.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristlegass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass] and/or

Labeled for over 40 different [grassy] [and] [broadleaf] weeds as listed

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	0.29%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine.....	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine.....	0.53%
Other Ingredients.....	98.91%
Total	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor’s note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N).....	XX%
Ammoniacal Nitrogen	XX%
Urea Nitrogen	XX%
Available Phosphoric Acid (P ₂ O ₅).....	XX%
Soluble Potash (K ₂ O)	XX%
Calcium	XX%
Magnesium (Mg)	XX%
Copper (Cu).....	XX%
Iron (Fe).....	XX%
Manganese (Mn)	XX%
Zinc (Zn).....	XX%

Keep Out of Reach of Children

CAUTION

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-888-478-0798.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 10163-344

EPA Est. _____

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

**Produced for
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569**

Net Weight _____

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/ Personal Protective Equipment (PPE) immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. Contact 1-888-478-0798 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

GroundWater Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons (or pets), either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow other people (or pets) to enter treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable nonrigid containers:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refillable rigid containers larger than 5 gal:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate

vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable rigid containers larger than 5 gal:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Product Information

This product is a granular fertilizer containing Gallery® and Team®, preemergence herbicides for application to established turfgrass including golf courses, parks, cemeteries, and non-residential and residential turf for control of certain broadleaf weeds and annual grasses. This product will not control existing weeds. Existing weeds should be controlled with a foliarly applied postemergence herbicide, either before or after application of this product.

Weed Resistance Management

For resistance management, please note that Turf Fertilizer Contains Gallery® Plus Team® contains both Group (21) Isoxaben and Group (3) Trifluralin and Benefin herbicides. Any weed population may contain plants naturally resistant to Group (21) and/or Group (3) herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same fields. Appropriate resistant-management strategies should be followed.

Best Management Practices

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Turf Fertilizer Contains Gallery® Plus Team® or other Group (21) or Group (3) herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - (2) a spreading patch of non-controlled plants of a particular weed species;
 - (3) surviving plants mixed with controlled individuals of the same species.

- If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Gowan Company representative.

Spray Drift Management

Spray equipment and weather affect spray drift. Consider all factors when making application decisions. Where states have more stringent regulations, they must be observed. Avoiding spray drift is the responsibility of the applicator or grower. To reduce the potential for drift, the application equipment must be set to apply medium to coarse droplets (i.e., ASAE Standard 572) with corresponding spray pressure. Use high flow rate nozzles to apply the highest practical spray volume. With most nozzle types, narrower spray angles produce larger droplets. Follow the nozzle manufacturer's directions on pressure, orientation, spray volume, etc. in order to minimize drift and optimize coverage and control.

Wind: Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and non-target plants are growing. Do not spray near sensitive plants if wind is gusty, below 2 mph, or in excess of 10 mph and moving in the direction of adjacent areas of sensitive areas. Local terrain may influence wind patterns; the applicator must be familiar with local conditions and understand how they may impact spray drift.

Sensitive Areas: Sensitive areas to this product are defined as bodies of water (ponds, lakes, rivers, streams, and ditches), wetlands, habitats of endangered species, and non-labeled agricultural crop areas. Applicators must take all precautions necessary to keep spray drift from reaching those areas.

Temperature Inversion: A surface temperature inversion (i.e., increasing temperature with increasing altitude) greatly increases the potential for drift. Presence of ground fog is a good indicator of a surface temperature inversion. Do not apply during temperature inversions. Always make applications when there is some air movement to determine the direction and distance of possible spray drift.

Boom Height: Set the boom and make applications at the lowest height that safely permits uniform coverage of the soil and minimizes droplet evaporation. Boom or nozzle shielding can reduce the effects of wind or air currents on drift. Verify that the shields do not interfere with uniform deposition of product prior to application.

Use Restrictions

- Do not apply to dichondra.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not make more than 2 applications of this product per year.
- Apply this product **only** to established turf. Apply to newly seeded turf (including overseeded turf) only after seedlings are established (three leaf stage and tillering) and well rooted (generally after 3 mowings).
- **Reseeding:** Do not reseed treated areas for at least 8 weeks after a single application of 275 lb of product per acre, or 12 to 16 weeks after a single application of 375 lb of product per acre.
- Do not apply to non-turf areas such as flower beds or gardens. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. The herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply this product in the spring and summer to tall fescue turfgrass seeded the previous fall.

- Do not apply in the spring to Bermudagrass that has been overseeded with winter annual grasses since thinning may occur.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.
- Apply this product only to the grass species listed on this label. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on turf species not listed on this label.**
- **Maximum Single Application Rate:** Cool season turf – 275 lb of Turf Fertilizer – Contains Gallery plus Team per acre; warm season turf – 375 lb of Turf Fertilizer – Contains Gallery plus Team per acre.
- **Maximum Yearly Rate:** Cool season turf – 550 lb of Turf Fertilizer – Contains Gallery plus Team per acre; warm season turf – 750 lb of Turf Fertilizer – Contains Gallery plus Team per acre.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- This product must be watered in as soon as possible after application.
- Not for sale, distribution or use in New York State.
- Do not apply by air.
- Do not apply Turf Fertilizer - Contains Gallery Plus Team to turfgrass grown for seed.

This product is for use on the following established turfgrass species:

Established Cool Season Turfgrass

Common Name	Scientific Name
bentgrass, colonial	<i>Agrostis tenuis</i>
bentgrass, creeping	<i>Agrostis stolonifera</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewings	<i>Festuca rubra var. commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheep	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

Established Warm Season Turfgrass

Common Name	Scientific Name
bahiagrass	<i>Paspalum notatum var. Saurae parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
fescue, tall	<i>Festuca arundinaceae</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

Weeds Controlled

Weeds controlled when applied at 250 to 375 lb per acre (5.75 to 8.6 lb/1000 sq ft):

Common Name	Scientific Name
annual bluegrass	<i>Poa annua</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
burweed, lawn	<i>Soliva pterosperma</i>
buttercup	<i>Ranunculus</i> spp.
carpetweed	<i>Mollugo verticillata</i>
celery, wild	<i>Apium leptophyllum</i>
chickweed, common	<i>Stellaria media</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
crabgrass	<i>Digitaria</i> spp.
large crabgrass	
smooth crabgrass	

field sandbur (field sandspur)	<i>Cenchrus incertus</i>
foolsparsley	<i>Aethusa cynapium</i>
foxtails	<i>Setaria</i> spp.
bottlegrass	
bristlegrass	
giant foxtail	
green foxtail	
pigeongrass	
robust foxtail	
geranium, Carolina	<i>Geranium carolinianum</i>
goosegrass	<i>Eleusine indica</i>
henbit	<i>Lamium amplexicaule</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
mallow, little	<i>Malva parviflora</i>
parsley - piert	<i>Alchemilla arvensis</i>
pennywort, lawn	<i>Hydrocotyle sibthorpioides</i>
pineappleweed	<i>Matricaria matricarioides</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
spurge, spotted	<i>Euphorbia maculata</i>
spurry, corn	<i>Spergula arvensis</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Weeds controlled by a repeat application 8 weeks after the first application when applied at 250 to 375 lb per acre (5.75 to 8.6 lb/1000 sq ft):

Common Name	Scientific Name
clover spp. ¹	<i>Trifolium</i> spp.
clover, white ¹	<i>Trifolium repens</i>
dandelion ¹	<i>Taraxacum officinale</i>
groundsel, common	<i>Senecio vulgaris</i>
medic, black	<i>Medicago lupulina</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
plantain, slender	<i>Plantago elongata</i>

¹From seed only

Application Directions

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/acre)				
	250	XXX	XXX	XXX	375
XXXXXXXX	XX	XX	XX	XX	XX

Rates and Timing of Application

Broadcast Application Rates

	Cool Season Turfgrass	Warm Season Turfgrass
summer annual grass weeds (spring application)	250 lb/acre (5.75 lb/1000 sq ft)	250 to 375 lb/acre (5.75 to 8.6 lb/1000 sq ft)
winter annual grass weeds (fall application)	275 lb/acre (6.3 lb/1000 sq ft)	250 to 375 lb/acre (5.75 to 8.6 lb/1000 sq ft)

Apply this product at a rate of 250 to 375 lb per acre. A minimum of 250 lb per acre is recommended for optimum results. Apply in late summer or early fall, approximately 3 weeks prior to germination of winter annual weeds, or in early spring, 3 weeks prior to germination of weeds which emerge in spring or early summer. Use the higher rate in the rate range where heavy weed infestation is anticipated.

**FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.
For other product information, contact Gowan Company or see Material Safety Data Sheet.**

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

EPA Text Pending: (to EPA 09-04-18)

[Sub Label B: Residential Uses]

(Base label):

Turf Fertilizer - Contains Gallery[®] plus Team[®]**Herbicide**

A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established home lawns.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

Labeled for over 40 different [grassy] [and] [broadleaf] weeds as listed

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	0.29%
trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N, N</i> -dipropyl- <i>p</i> -toluidine	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α, α, α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.53%
Other Ingredients	98.91%
Total	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N)	XX%
Ammoniacal Nitrogen	XX%
Urea Nitrogen	XX%
Available Phosphoric Acid (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%
Calcium	XX%
Magnesium (Mg)	XX%
Copper (Cu)	XX%
Iron (Fe)	XX%
Manganese (Mn)	XX%
Zinc (Zn)	XX%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/Personal Protective Equipment (PPE) immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. Contact 1-888-478-0798 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

GroundWater Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-888-478-0798.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 10163-344

EPA Est. _____

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

Produced for

Gowan Company

P.O. Box 5569

Yuma, AZ 85366-5569

Net Weight _____

(Label booklet cover):

Turf Fertilizer - Contains Gallery® plus Team®

Herbicide

A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established home lawns.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

Labeled for over 40 different [grassy] [and] [broadleaf] weeds as listed

Active Ingredients:

isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-5-isoxazoly]-
2,6-dimethoxybenzamide and isomers 0.29%

trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine.....	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α, α, α -trifluoro-2,6-dinitro- <i>p</i> -toluidine.....	0.53%
Other Ingredients.....	98.91%
Total	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis: XX-XX-XX

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N).....	XX%
Ammoniacal Nitrogen	XX%
Urea Nitrogen	XX%
Available Phosphoric Acid (P ₂ O ₅).....	XX%
Soluble Potash (K ₂ O)	XX%
Calcium	XX%
Magnesium (Mg)	XX%
Copper (Cu).....	XX%
Iron (Fe).....	XX%
Manganese (Mn)	XX%
Zinc (Zn).....	XX%

Keep Out of Reach of Children

CAUTION

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-888-478-0798.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 10163-344

EPA Est. _____

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

**Produced for
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569**

Net Weight _____

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

User Safety Recommendations

Users should:

- Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/Personal Protective Equipment (PPE) immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. Contact 1-888-478-0798 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

GroundWater Advisory: This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not allow people or pets to enter the treated areas until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

How This Product Works

This product is a granular fertilizer containing Gallery® and Team®, preemergence herbicides for application to established home lawns for control of certain broadleaf weeds and annual grasses. This product also feeds the lawn with a fertilizer designed to provide prolonged, even greening without excess top growth.

This product will not control existing weeds. Existing weeds should be controlled with a foliarly applied postemergence herbicide either before or after application of this product.

Spray Drift Management

Spray equipment and weather affect spray drift. Consider all factors when making application decisions. Where states have more stringent regulations, they must be observed. Avoiding spray drift is the responsibility of the applicator or grower. To reduce the potential for drift, the application equipment must be set to apply medium to coarse droplets (i.e., ASAE Standard 572) with corresponding spray pressure. Use high flow rate nozzles to apply the highest practical spray volume. With most nozzle types, narrower spray angles produce larger droplets. Follow the nozzle manufacturer's directions on pressure, orientation, spray volume, etc. in order to minimize drift and optimize coverage and control.

Wind: Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and non-target plants are growing. Do not spray near sensitive plants if wind is gusty, below 2 mph, or in excess of 10 mph and moving in the direction of adjacent areas of sensitive areas. Local terrain may influence wind patterns; the applicator must be familiar with local conditions and understand how they may impact spray drift.

Sensitive Areas: Sensitive areas to this product are defined as bodies of water (ponds, lakes, rivers, streams, and ditches), wetlands, habitats of endangered species, and non-labeled agricultural crop areas. Applicators must take all precautions necessary to keep spray drift from reaching those areas.

Temperature Inversion: A surface temperature inversion (i.e., increasing temperature with increasing altitude) greatly increases the potential for drift. Presence of ground fog is a good indicator of a surface temperature inversion. Do not apply during temperature inversions. Always make applications when there is some air movement to determine the direction and distance of possible spray drift.

Boom Height: Set the boom and make applications at the lowest height that safely permits uniform coverage of the soil and minimizes droplet evaporation. Boom or nozzle shielding can reduce the effects of wind or air currents on drift. Verify that the shields do not interfere with uniform deposition of product prior to application.

Use Restrictions

- Do not apply to dichondra.

- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not make more than 2 applications of this product per year.
- Apply this product **only** to established turf. Apply to newly seeded turf (including overseeded turf) only after seedlings are established and well rooted (generally after 3 mowings).
- **Reseeding:** Do not reseed treated areas for at least 8 weeks following application of this product.
- Do not apply to non-turf areas such as flower beds or gardens. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. The herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply this product in the spring and summer to tall fescue turfgrass seeded the previous fall.
- Do not apply in the spring to Bermudagrass that has been overseeded with winter annual grasses since thinning may occur.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.
- Apply this product only to the grass species listed on this label. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on turf species not listed on this label.**
- **Maximum Single Application Rate:** 7.2 lb of Turf Fertilizer – Contains Gallery plus Team per 1000 sq ft.
- **Maximum Yearly Rate:** 14.4 lb of Turf Fertilizer – Contains Gallery plus Team per 1000 sq ft.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- Not for sale, distribution or use in New York State.
- Do not apply by air.
- Do not apply Turf Fertilizer - Contains Gallery Plus Team to turfgrass grown for seed.

This product is for use on the following established turfgrass species:

Established Cool Season Turfgrass

bentgrass, colonial
bentgrass, creeping
bluegrass, Kentucky
fescue, chewings
fescue, creeping red
fescue, sheep
fescue, tall
ryegrass, perennial

Established Warm Season Turfgrass

bahiagrass
Bermudagrass
centipedegrass
fescue, tall
St. Augustinegrass
zoysiagrass

Weeds Controlled

Weeds controlled when applied prior to weed germination:

annual bluegrass	geranium, Carolina
barnyardgrass	goosegrass
burweed, lawn	henbit
buttercup	knotweed, prostrate
carpetweed	mallow, little
celery, wild	parsley - piert
chickweed, common	pennywort, lawn
chickweed, mouseear	pineappleweed
crabgrass	speedwell, thymeleaf
large crabgrass	spurge, prostrate
smooth crabgrass	spurge, spotted
field sandbur (field sandspur)	spurry, corn

foolsparsley	woodsorrel, creeping
foxtails	woodsorrel, yellow
bottlegrass	
bristlegrass	
giant foxtail	
green foxtail	
pigeongrass	
robust foxtail	

Weeds controlled by a repeat application 8 weeks after the first application when applied prior to weed germination:

clover spp. ¹	medic, black
clover, white ¹	plantain, broadleaf
dandelion ¹	plantain, buckhorn
groundsel, common	plantain, slender

¹From seed only

How to Apply

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

Spreader Settings as a Guide for Calibration

Brand and Model of Spreader	Calibration Settings at Specified Rate (lb/1000 sq ft)				
	5.75	XX	XX	XX	8.6
XXXXXXXX	XX	XX	XX	XX	XX

How Much to Apply

Apply this product to cool season turfgrass at a rate of 5.75 lb per 1000 sq ft and to warm season turfgrass at a rate of 7.2 lb per 1000 sq ft.

When to Apply

This product can be applied at any time of the spring or fall. The type of weeds controlled will vary depending upon time of application so other weed control products may be needed to provide a total annual weed control program.

Early spring (before forsythia blooms): If applied in early spring, this product will prevent crabgrass and other annual grasses (including *Poa annua*) and broadleaf weeds, such as dandelion, clover, plantain, and spurge, that normally germinate in the spring and summer. When used in this manner, this product may be substituted for a multi-step weed and feed program. For your area, consult the local extension service for optimum timing for control of crabgrass.

Mid-late spring (after forsythia blooms, before Memorial Day): This product will prevent annual grasses (including *Poa annua*) and broadleaf weeds, such as dandelion, clover, plantain, and spurge, that have yet to germinate in the spring and summer, ensuring season-long weed control.

Early fall (before average night temperatures fall below 60°F): This product will prevent establishment of winter annual grasses (including *Poa annua*) and broadleaf weeds, such as chickweed, henbit, and speedwell, that normally germinate in the fall and winter and broadleaf weeds that germinate the following spring season. A supplemental application of a crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control..

Late fall: This product will prevent establishment of broadleaf weeds, such as chickweed, henbit, and speedwell, that may germinate over winter and broadleaf weeds that germinate the following spring season. A supplemental application of a crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control.

**FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.
For other product information, contact Gowan Company or see Material Safety Data Sheet.**

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Team® is a registered trademark of Gowan Company, L.L.C.
All other brands are trademarks of their respective owners.

EPA Text Pending: (to EPA 9-20-17)