



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 23, 2022

Kyla S. Smith
Registration Specialist
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

Subject: Registration Review Label Mitigation for Benfluralin and Trifluralin
Product Name: TURF FERTILIZER - CONTAINS TEAM PRO 0.86%
HERBICIDE
EPA Registration Number: 10163-346
Application Dates: 6/11/19, 3/7/22
Decision Numbers: 582479,582480

Dear Ms. Smith:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Benfluralin and Trifluralin Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2
EPA Reg. No. 10163-346
Decision No. 582479,582480

If you have any questions about this letter, please contact DeMariah Koger by phone at (202)-566-2288, or via email at koger.demariah@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a long horizontal flourish extending to the right.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure

TRIFLURALIN / BENFLURALIN

GROUP

3

HERBICIDE

Turf Fertilizer - Contains Team[®] Pro 0.86%

The master label consists of separate, self-contained sub labels for commercial use and residential use, respectively.

A C C E P T E D

Mar 23, 2022

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 10163-346

[Sub Label A: Commercial Uses]

Turf Fertilizer - Contains Team[®] Pro 0.86% Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established turfgrass and landscape ornamentals

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

and/or

Controls [tough] weeds [such as annual bluegrass] [large, smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [common chickweed] [prostrate knotweed] [common lambsquarters] [pigweed] [common purslane]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees and shrubs

and/or

Never see crabgrass again**Refer to label booklet for Directions for Use.**

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

NET CONTENTS _____

EPA Reg. No. 10163-346
EPA Est. No.

Gowan
The Go To Company

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

Active Ingredients:

| | |
|---|---------|
| benfluralin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine | 0.43% |
| trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine | 0.43% |
| Other Ingredients | 99.14% |
| Total | 100.00% |

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

| Guaranteed Fertilizer Analysis | |
|--|------|
| Total Nitrogen (N) | X.X% |
| X.X% Nitrate Nitrogen | |
| X.X% Water Soluble Nitrogen | |
| X.X% Water Insoluble Nitrogen | |
| Available Phosphoric Acid (P ₂ O ₅) | X.X% |
| Soluble Potash (K ₂ O) | X.X% |

Keep Out of Reach of Children WARNING

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

| FIRST AID | |
|---|---|
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-478-0798 for emergency medical treatment information. | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In some Individuals
Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- Goggles, faceshield or safety glasses

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Non-Target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons or pets, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow people or pets to enter the treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

WEED RESISTANCE MANAGEMENT

For resistance management, please note that Turf Fertilizer Contains Team 0.92% is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to Turf Fertilizer Contains Team 0.92% and Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

Best Management Practices

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Turf Fertilizer Contains Team 0.92% or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - (2) a spreading patch of non-controlled plants of a particular weed species;
 - (3) surviving plants mixed with controlled individuals of the same species.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Gowan Company representative.

GENERAL INFORMATION

This product is a granular fertilizer containing Team®, a preemergence herbicide for application to established turfgrass and landscape ornamentals (trees, shrubs, groundcovers, and flowers) for control of annual grasses and broadleaf weeds.

General Use Precautions and Restrictions

- Do not apply this product to plants that will bear fruits, berries, or nuts within 12 months of application.
- Do not aerially apply this product
- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- This product must be soil-incorporated within 24 hours of application either by mechanical incorporation or at least 0.5 acre inch of water (rainfall, irrigation). If a second incorporation is needed it should be delayed 3 to 5 days after the first incorporation and be completed prior to planting.

Reseeding: Do not reseed treated areas for at least 8 weeks after a single application of 174 lb of product per acre, or 12 to 16 weeks after a single application of 349 lb of product per acre unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 174 lb of product per acre rate is applied, or 10 to 12 weeks after the last 233 lb of product per acre rate is applied unless injury can be tolerated.

This product is recommended for use on the following turfgrass species:

Established Cool Season Turfgrass

bentgrass fairways
fescues
Kentucky bluegrass
perennial ryegrass

Established Warm Season Turfgrass

bahiagrass
Bermudagrass
centipedegrass
fescue, tall
St. Augustinegrass
zoysiagrass

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees

Common Name

almond
apple, crabapple
apricot
ash, white
bald cypress
birch, European white
blackgum
cherry
chestnut, Chinese
cottonwood
dogwood, flowering
dogwood, kousa
douglas fir
fir, balsam
hemlock, Canada
honeylocust
larch, Japanese
locust, black
maple, Norway
maple, red
maple, silver
maple, sugar
oak, pin
oak, scarlet
peach
pine, Austrian
pine, eastern white
pine, Japanese black
pine, loblolly
pine, red
pine, scotch
planetree, London
plum
redbud, eastern
spruce, Colorado
spruce, Norway
spruce, white
sweetgum
sycamore
tuliptree
walnut, black

Scientific Name

Prunus dulcis
Malus spp.
Prunus armeniaca
Fraxinus americana
Taxodium distichum
Betula pendula
Nyssa sylvatica
Prunus spp.
Castanea mollissima
Populus deltoides
Cornus florida
Cornus kousa
Pseudotsuga menziesii
Abies balsamea
Tsuga canadensis
Gleditsia triacanthos
Larix kaempferi
Robinia pseudoacacia
Acer platanoides
Acer rubrum
Acer saccharinum
Acer saccharum
Quercus palustris
Quercus coccinea
Prunus persica
Pinus nigra
Pinus strobus
Pinus thunbergiana
Pinus taeda
Pinus resinosa
Pinus sylvestris
Platanus acerifolia
Prunus spp.
Cercis canadensis
Picea pungens
Picea abies
Picea glauca
Liquidambar styraciflua
Platanus occidentalis
Liriodendron tulipifera
Juglans nigra

Ornamental Shrubs

Common Name

andromeda, Japanese
arborvitae, American
azalea
barberry, Japanese
barberry, mento
boxwood, common
boxwood, littleleaf
camellia, Japanese

Scientific Name

Pieris japonica
Thuja occidentalis
Rhododendron spp.
Berberis thunbergii
Berberis mentorensis
Buxus sempervirens
Buxus microphylla
Camellia japonica

| | |
|-------------------------|--------------------------------|
| camellia, sasanqua | <i>Camellia sasanqua</i> |
| cherrylaurel, American | <i>Prunus caroliniana</i> |
| cinquefoil | <i>Potentilla</i> spp. |
| cleyera, Japanese | <i>Cleyera japonica</i> |
| cotoneaster, cranberry | <i>Cotoneaster apiculata</i> |
| cotoneaster, zabel | <i>Cotoneaster zabelii</i> |
| deutzia | <i>Deutzia</i> spp. |
| elaegnus, silverberry | <i>Elaeagnus pungens</i> |
| euonymus, spreading | <i>Euonymus kiautschovica</i> |
| euonymus, winged | <i>Euonymus alatus</i> |
| euonymus, wintercreeper | <i>Euonymus fortunei</i> |
| firethorn | <i>Pyracantha</i> spp. |
| forsythia | <i>Forsythia</i> spp. |
| guava, pineapple | <i>Feijoa sellowiana</i> |
| holly | <i>Ilex</i> spp. |
| honeysuckle | <i>Lonicera</i> spp. |
| India hawthorn | <i>Raphiolepis indica</i> |
| juniper | <i>Juniperus</i> spp. |
| laurel, mountain | <i>Kalmia latifolia</i> |
| lilac, common | <i>Syringa vulgaris</i> |
| mockorange | <i>Philadelphus</i> spp. |
| pittosporum, Japanese | <i>Pittosporum tobira</i> |
| privet | <i>Ligustrum</i> spp. |
| redcedar, eastern | <i>Juniperus virginiana</i> |
| rhododendron | <i>Rhododendron</i> spp. |
| spiraea, vanhoutte | <i>Spiraea vanhouttei</i> |
| viburnum | <i>Viburnum</i> spp. |
| weigela | <i>Weigela</i> spp. |
| willow | <i>Salix</i> spp. |
| yew, anglojap | <i>Taxus media</i> |
| yew, Japanese | <i>Taxus cuspidata</i> |
| yewpine | <i>Podocarpus macrophyllus</i> |

Ornamental Groundcover Plantings

| Common Name | Scientific Name |
|------------------------|----------------------------------|
| aaronsbeard | <i>Hypericum calycinum</i> |
| bellflower, adriatic | <i>Campanula elatines</i> |
| bellflower, poscharsky | <i>Campanula poscharskyana</i> |
| ceanothus | <i>Ceanothus</i> spp. |
| coreopsis | <i>Coreopsis</i> spp. |
| cotoneaster | <i>Cotoneaster</i> spp. |
| coyote brush | <i>Baccharis pilularis</i> |
| crown vetch | <i>Coronilla vana</i> |
| daisy trailing African | <i>Osteospermum fruticosum</i> |
| fern, asparagus | <i>Asparagus densiflorus</i> |
| gazania | <i>Gazania</i> spp. |
| germander | <i>Teucrium chamaedrys</i> |
| ice plant, largeleaf | <i>Carpobrotus edulis</i> |
| ivy, Algerian | <i>Hedera canariensis</i> |
| ivy, English | <i>Hedera helix</i> |
| lily-of-the-nile | <i>Agapanthus</i> spp. |
| lilyturf, bigblue | <i>Liriope muscari</i> |
| marigold | <i>Tagetes</i> spp. |
| myoporum | <i>Myoporum laetum</i> |
| plumbago, dwarf | <i>Ceratostigma umbaginoides</i> |
| rockrose | <i>Cistus</i> spp. |
| rosemary | <i>Rosmarinus officinalis</i> |
| rupturewort | <i>Herniaria glabra</i> |
| snow-in-summer | <i>Cerastium tomentosum</i> |
| speedwell | <i>Veronica</i> spp. |
| St. Johnswort | <i>Hypericum coris</i> |
| stonecrop (sedum) | <i>Sedum</i> spp. |
| strawberry, beach | <i>Fragaria chiloensis</i> |
| thrift | <i>Armeria maritima</i> |
| verbena | <i>Verbena</i> spp. |
| wirevine, creeping | <i>Muehlenbeckia axillaris</i> |
| yarrow, wooly | <i>Achillea tomentosa</i> |

Roses and Other Established Flowers

| | | | |
|-------------------|---------------|-------------------|----------------------|
| African daisy | dianthus | marigold | shasta daisy |
| aster (perennial) | dusty miller | marigold, cape | snapdragon |
| balsam | floss flower | morningglory | snow-on-the-mountain |
| blackeyed susan | forget-me-not | nasturtium | stock |
| calendula | four o'clock | nicotiana | sunflower |
| carnation | gaillardia | petunia | sweet alyssum |
| centaurea, velvet | gladiolus | phlox | sweet pea |
| chrysanthemum | golden glow | pincushion flower | sweet sultan |
| coreopsis | impatiens | poppy, California | sweet william |
| cornflower | ixora | portulaca | vinca |
| cosmos | lobelia | rose | yarrow |
| dahlia | lupine | salvia | zinnia |

Weeds Controlled**Weeds controlled when applied prior to weed germination:**

| Common Name | Scientific Name |
|---|-------------------------------|
| Annual Grasses | |
| annual bluegrass | <i>Poa annua</i> |
| barnyardgrass (watergrass) | <i>Echinochloa crus-galli</i> |
| crabgrass, large or hairy | <i>Digitaria sanguinalis</i> |
| crabgrass, smooth | <i>Digitaria ischaemum</i> |
| goosegrass (silver crabgrass or crowfoot) | <i>Eleusine indica</i> |
| green foxtail | <i>Setaria viridis</i> |
| yellow foxtail | <i>Setaria glauca</i> |

| | |
|------------------------|----------------------------|
| Broadleaf Weeds | |
| chickweed, common | <i>Stellaria media</i> |
| knotweed, prostrate | <i>Polygonum aviculare</i> |
| lambsquarters, common | <i>Chenopodium album</i> |
| pigweed | <i>Amaranthus</i> spp. |
| purslane, common | <i>Portulaca oleracea</i> |

Weeds Suppressed**Weeds suppressed when applied prior to weed germination:**

| Common Name | Scientific Name |
|---------------------|-----------------------------|
| creeping spurge | <i>Euphorbia serpens</i> |
| creeping woodsorrel | <i>Oxalis corniculata</i> |
| panicum, Texas | <i>Panicum texanum</i> |
| prostrate spurge | <i>Euphorbia humistrata</i> |
| pusley, Florida | <i>Richardia scabra</i> |
| spotted spurge | <i>Euphorbia maculata</i> |
| woodsorrel | <i>Oxalis</i> spp. |

APPLICATION DIRECTIONS

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Spreader Settings as a Guide for Calibration

| Brand and Model of Spreader | Calibration Settings at Specified Rate (lb/acre) | | | | |
|-----------------------------|--|-----|-----|-----|-----|
| | XXX | XXX | XXX | XXX | XXX |
| XXXXXXXX | XX | XX | XX | XX | XX |

USES

Established Turfgrass

Application Timing

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

In areas where germination of the annual weed grass species continues for extended period of time, make a second application at the recommended rate 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

Broadcast Application Rates

| | Cool Season Turfgrass | Warm Season Turfgrass |
|---|---|----------------------------------|
| summer annual grass weeds (spring application) | 174 to 349lb/acre 4 to 8 lb/1000 sq ft) | 349 lb/acre (8 lb/1000 sq ft) |
| winter annual grass weeds (fall application) | 174 to 349 lb/acre 4 to 8 lb/1000 sq ft) | 349 lb/acre (8 lb/1000 sq ft) |

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 174 lb plus 174 lb per acre for cool season turfgrass and in the northern half of the warm season turfgrass area, or 233 lb plus 233 lb per acre in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 174 lb per acre rate and 10 to 12 weeks following the 233 lb per acre rate.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** 349 lb of Turf Fertilizer – Contains Team Pro 0.86% per acre.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- This product must be watered in as soon as possible after application.

[Editor's note: Supplemental distributor will add one of the following statements as appropriate if their product is registered in the state of Florida.]

- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.
- Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, 'Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant'" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>.
- Florida Applications: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf.

Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

Application Timing

Apply 1 to 2 weeks prior to the germination of weed seeds or clean cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

Broadcast Application Rates

Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours after application.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
 - **Maximum Single Application Rate:** 8 lb of T&O Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.
 - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
 - **Maximum Single Application Rate:** 8 lb of T&O Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.

(Storage and Disposal for rigid containers 5 gal or less)**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrigid containers any size)**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for refillable rigid containers greater than 5 gal)**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.
For other product information, contact Gowan Company or see Material Safety Data Sheet.**

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. All such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Team® is a registered trademark of Gowan Company LLC.
All other brands are registered trademarks of their respective owners.

[Sub Label B: Residential Uses]

Turf Fertilizer - Contains Team[®] Pro 0.86%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual grasses and broadleaf weeds in established home lawns and landscape ornamentals.

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

Covers up to XX,000 sq ft (XXX' x XX')

and/or

This bag covers XX,000 sq ft

and/or

ONE STEP – extended grassy weed control

and/or

Controls [tough] weeds [such as annual bluegrass] [large, smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [common chickweed] [prostrate knotweed] [common lambsquarters] [pigweed] [common purslane]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees and shrubs

and/or

Never see crabgrass again

NET CONTENTS _____

EPA Reg. No. 10163-346

EPA Est. No.

Gowan
The Go To Company

Produced For:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Active Ingredients:

| | |
|---|---------|
| benfluralin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine | 0.43% |
| trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine | 0.43% |
| Other Ingredients | 99.14% |
| Total | 100.00% |

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

| Guaranteed Fertilizer Analysis | |
|--|------|
| Total Nitrogen (N) | X.X% |
| X.X% Nitrate Nitrogen | |
| X.X% Water Soluble Nitrogen | |
| X.X% Water Insoluble Nitrogen | |
| Available Phosphoric Acid (P ₂ O ₅) | X.X% |
| Soluble Potash (K ₂ O) | X.X% |

Keep Out of Reach of Children WARNING

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

| FIRST AID | |
|---|---|
| If in eyes | <ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-478-0798 for emergency medical treatment information. | |

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals
Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt and long pants, shoes plus socks and goggles or safety glasses.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Non-Target Organism Advisory Statement: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not allow people or pets to enter the treated area until dusts have settled. If watering-in is required after application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

WEED RESISTANCE MANAGEMENT

For resistance management, please note that Turf Fertilizer Contains Team 0.92% is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to Turf Fertilizer Contains Team 0.92% and Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

Best Management Practices

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Turf Fertilizer Contains Team 0.92% or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - (2) a spreading patch of non-controlled plants of a particular weed species;
 - (3) surviving plants mixed with controlled individuals of the same species.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Gowan Company representative.

HOW THIS PRODUCT WORKS

This product is a granular fertilizer containing Team[®], a preemergence herbicide for application to established home lawns and landscape ornamentals (trees, shrubs, groundcovers and flowers) for control of annual grasses and broadleaf weeds.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

- Do not apply this product to plants that will bear fruits, berries, or nuts within 12 months of application.
- Do not aerially apply this product
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- This product must be soil-incorporated within 24 hours of application either by mechanical incorporation or at least 0.5 acre inch of water (rainfall, irrigation). If a second incorporation is needed it should be delayed 3 to 5 days after the first incorporation and be completed prior to planting.

Reseeding: Do not reseed treated areas for at least 8 weeks after a single application of 4 lb of product per 1000 sq ft, or 12 to 16 weeks after a single application of 8 lb of product per 1000 sq ft unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 4 lb of product per 1000 sq ft rate is applied, or 10 to 12 weeks after the last 5.34 lb of product per 1000 sq ft rate is applied unless injury can be tolerated.

This product is recommended for use on the following turfgrass species:

Established Cool Season Turfgrass

bentgrass fairways
fescues
Kentucky bluegrass
perennial ryegrass

Established Warm Season Turfgrass

bahiagrass
Bermudagrass
centipedegrass
fescue, tall
St. Augustinegrass
zoysiagrass

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees

| | | |
|-----------------------|----------------------|-------------------|
| almond | hemlock, Canada | pine, loblolly |
| apple, crabapple | honeylocust | pine, red |
| apricot | larch, Japanese | pine, scotch |
| ash, white | locust, black | planetree, London |
| bald cypress | maple, Norway | plum |
| birch, European white | maple, red | redbud, eastern |
| blackgum | maple, silver | spruce, Colorado |
| cherry | maple, sugar | spruce, Norway |
| chestnut, Chinese | oak, pin | spruce, white |
| cottonwood | oak, scarlet | sweetgum |
| dogwood, flowering | peach | sycamore |
| dogwood, kousa | pine, Austrian | tuliptree |
| douglas fir | pine, eastern white | walnut, black |
| fir, balsam | pine, Japanese black | |

Ornamental Shrubs

| | |
|-------------------------|-----------------------|
| andromeda, Japanese | forsythia |
| arborvitae, American | guava, pineapple |
| azalea | holly |
| barberry, Japanese | honeysuckle |
| barberry, mento | India hawthorn |
| boxwood, common | juniper |
| boxwood, littleleaf | laurel, mountain |
| camellia, Japanese | lilac, common |
| camellia, sasanqua | mockorange |
| cherrylaurel, American | pittosporum, Japanese |
| cinquefoil | privet |
| cleyera, Japanese | redcedar, eastern |
| cotoneaster, cranberry | rhododendron |
| cotoneaster, zabel | spiraea, vanhoutte |
| deutzia | viburnum |
| elaeanus, silverberry | weigela |
| euonymus, spreading | willow |
| euonymus, winged | yew, anglojap |
| euonymus, wintercreeper | yew, Japanese |
| firethorn | yewpine |

Ornamental Groundcover Plantings

| | |
|------------------------|--------------------|
| aaronsbeard | lilyturf, bigblue |
| bellflower, adriatic | marigold |
| bellflower, poscharsky | myoporum |
| ceanothus | plumbago, dwarf |
| coreopsis | rockrose |
| cotoneaster | rosemary |
| coyote brush | rupturewort |
| crown vetch | snow-in-summer |
| daisy trailing African | speedwell |
| fern, asparagus | St. Johnswort |
| gazania | stonecrop (sedum) |
| germander | strawberry, beach |
| ice plant, largeleaf | thrift |
| ivy, Algerian | verbena |
| ivy, English | wirevine, creeping |
| lily-of-the-nile | yarrow, wooly |

Roses and Other Established Flowers

| | | | |
|-------------------|---------------|----------------|----------------------|
| African daisy | dianthus | marigold | shasta daisy |
| aster (perennial) | dusty miller | marigold, cape | snapdragon |
| balsam | floss flower | morningglory | snow-on-the-mountain |
| blackeyed susan | forget-me-not | nasturtium | stock |
| calendula | four o'clock | nicotiana | sunflower |

| | | | |
|-------------------|-------------|-------------------|---------------|
| carnation | gaillardia | petunia | sweet alyssum |
| centaurea, velvet | gladiolus | phlox | sweet pea |
| chrysanthemum | golden glow | pincushion flower | sweet sultan |
| coreopsis | impatiens | poppy, California | sweet william |
| cornflower | ixora | portulaca | vinca |
| cosmos | lobelia | rose | yarrow |
| dahlia | lupine | salvia | zinnia |

Weeds Controlled

Weeds controlled when applied prior to weed germination:

Annual Grasses

annual bluegrass
barnyardgrass (watergrass)
crabgrass, large or hairy
crabgrass, smooth
goosegrass (silver crabgrass or
crowfoot)
green foxtail
yellow foxtail

Broadleaf Weeds

chickweed, common
knotweed, prostrate
lambsquarters, common
pigweed
purslane, common

Weeds Suppressed

Weeds suppressed when applied prior to weed germination:

creeping spurge
creeping woodsorrel
panicum, Texas
prostrate spurge
pusley, Florida
spotted spurge
woodsorrel

HOW TO APPLY

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate the application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours. A single rainfall or spinkler irrigation of 0.5 inch or more is required to activate this product.

Spreader Settings as a Guide for Calibration

| Brand and Model of Spreader | Calibration Settings at Specified Rate (lb/1000 sq ft) | | | | |
|-----------------------------|--|----|----|----|----|
| | XX | XX | XX | XX | XX |
| XXXXXXXX | XX | XX | XX | XX | XX |

USES

Established Home Lawns

When to Apply

Apply this product 1 to 2 weeks prior to germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

How Much to Apply

| | Cool Season Turfgrass | Warm Season Turfgrass |
|---|-----------------------|-----------------------|
| summer annual grass weeds (spring application) | 4 to 8 lb/1000 sq ft | 8 lb/1000 sq ft |
| winter annual grass weeds (fall application) | 4 to 8 lb/1000 sq ft | 8 lb/1000 sq ft |

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 4 lb plus 4 lb per 1000 sq ft for cool season turfgrass and in the northern half of the warm season turfgrass area, or 5.34

lb plus 5.34 lb per 1000 sq ft in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 4 lb per 1000 sq ft rate and 10 to 12 weeks following the 5.34 lb per 1000 sq ft rate.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on turfgrass.
- **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.

[Editor's note: Supplemental distributor will add one of the following statements as appropriate if their product is registered in the state of Florida.]

- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.

Established Ornamental Plants in Landscape Settings

Apply to ornamental trees, shrubs, groundcovers and flowers, and to non-bearing fruit and nut trees and non-bearing vineyards, which are defined as plants that will not bear fruit for at least 1 year after treatment.

When to Apply

Apply 1 to 2 weeks prior to the germination of weed seeds or clean cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds.

How to Apply

Apply this product at a rate of 4 to 8 lb per 1000 sq ft as a spring application or 4 to 8 lb per 1000 sq ft as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours after application.

Specific Use Restrictions

- Do not make more than 2 applications of this product per year on all ornamental plants.
- **Non-bearing fruit and nut trees, non-bearing vineyards:**
 - **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.
 - Applications to plants that will bear fruits, berries, or nuts within 12 months are prohibited.
- **Landscape ornamentals:**
 - **Maximum Single Application Rate:** 8 lb of Turf Fertilizer – Contains Team Pro 0.86% per 1000 sq ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.

For other product information, contact Gowan Company or see Material Safety Data Sheet.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. All such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN

COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Team® is a registered trademark of Gowan Company LLC.
All other brands are registered trademarks of their respective owners.

Turf Fert Ctns Team Pro 0.86 (to EPA 6-12-19, 2-22-22)