



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

October 17, 2025

Ana Sanchez
Registration Specialist
Gowan Company
370 S. Main Street, P.O. Box 5569
Yuma, AZ 85366-5569

Subject: Label Amendment – Correct of Typographical Error in the Retreatment Interval Specified in Crop Use Directions
Product Name: Trigard® Insecticide
EPA Registration Number: 10163-411
Application Date: June 23, 2025
Action Case Number: 663938

Dear Ana Sanchez:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find

or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Carmen Rodia via e-mail at Rodia.Carmen@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Tamica L. Cain".

Tamica L. Cain, Product Manager 10
Invertebrate & Vertebrate Branch 2
Registration Division (7505T)

Enclosure: *Stamped "Accepted" Master Label, dated October 17, 2025*

{NOTE TO REVIEWER: Text in braces { } is for informational purposes and will not appear on the end-use label.}
{NOTE TO REVIEWER: Text in brackets [] is optional, and may or may not appear on the end-use label}
{NOTE TO REVIEWER: Text in parentheses () will appear on the end-use label.}

CYROMAZINE	GROUP	17	INSECTICIDE
------------	-------	----	-------------

Trigard[®] Insecticide

[ALTERNATE BRAND NAME: Trigard[®] 75W]

For control of leafminers in Brassica, leafy greens, (subgroup 4-16B); Bulb vegetables (subgroups 3-07A and 3-07B); Celtnce; Cucurbits; Dried beans (except cowpeas and soybeans); Florence fennel; Leaf petiole (subgroup 22B); Kohlrabi; Leafy Greens (subgroup 4-16A); Pepper/Eggplant (subgroup 8-10B); Succulent Beans and Peas; Tomato (subgroup 8-10A); Vegetable, brassica, head and stem (group 5-16), except broccoli; Broccoli; and control of Colorado potato beetle and leafminer in potatoes and Vegetable, tuberous and corm (subgroup 1C).

ACTIVE INGREDIENT: Cyromazine (CAS No. 66215-27-8)	% BY WT.
N-cyclopropyl-1,3,5-triazine-2,4,6-triamine.....	75.0%
OTHER INGREDIENTS:	25.0%
TOTAL:	100.0%

Trigard Insecticide is a wettable powder.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center, or doctor, or going for treatment. For non-emergency, general information on this pesticide product (including health concerns or pesticide incidents), you may call 1-888-478- 0798, 24 hours per day, 7 days per week.	

[See [inside] [side] [back panel] [booklet] [attached] [to] [pouch] [bag] [sealed pack] for [complete] [additional] [First Aid Statements,] [Precautionary Statements,] [Directions For Use,] [Storage and Disposal,] and [Conditions of Sale and Warranty.]]


[This outer protective container contains Trigard Insecticide in 6 small water-soluble packets. These packets and their contents dissolve in water. After opening outer container, immediately dump the required number of unopened packets into the partially filled sprayer or mix tank. **DO NOT** handle the soluble packets or expose them to moisture, since this may cause rupturing.]

Net Contents:

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

EPA Reg. No. 10163-411
EPA Est. No.



Produced For:
Gowan Company, LLC
P. O. Box 5569
Yuma, AZ 85366-5569

Personal Protective Equipment

Applicators and other handlers must wear:

- Coveralls worn over long-sleeved shirt and long pants;
- Chemical-resistant gloves such as barrier laminate or butyl rubber ≥14 mils or nitrile rubber ≥14 or neoprene rubber ≥14 mils or natural rubber ≥14 or polyethylene or polyvinyl chloride (PVC) ≥14 mils or Viton > 14 mils; and
- Chemical-resistant footwear plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

Water soluble packets, when used correctly, qualify as a closed mixing/loading system under the Worker Protection Standard [40 CFR 170.607(d)]. Mixers and loaders handling this product while it is enclosed in intact water soluble packets may elect to wear reduced PPE of long-sleeved shirt, long pants, shoes, socks, a chemical-resistant apron, and chemical-resistant gloves. When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water. **DO NOT** apply when weather conditions favor drift from target area.

Ground Water Advisory

A metabolite of Trigard Insecticide has been found in ground water, possibly as a result of agricultural use. Use of this product in areas where soils are permeable and water tables are shallow could result in leaching to ground water. Irrigation could increase the risk of contamination.

PROTECTION OF POLLINATORS

"Cyromazine acts as a growth regulator, affecting the molting process of target pests, and therefore, may have an effect on bee larvae, but not on adult bees. Exposure to adult bees, either through overspray or contact with residues may lead to effects to bee larvae."

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product, Applicators Should Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS FOR ALL CROPS

- Trigard Insecticide must be used only as directed in accordance with recommendations on this label or in separately published Gowan Company, LLC supplemental labeling recommendations for this product.
- **DO NOT** apply this product through any type of irrigation (chemigation) system.
- **DO NOT** use this product in greenhouses.
- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.
- **DO NOT** let spray mixture stand overnight in the spray tank. Rinse spray tank at the end of the day.
- To avoid spray drift, **DO NOT** apply under windy conditions. Avoid spray overlap, because crop injury may result.
- Use aerial applications only when conditions prevent application with ground equipment.
- When using ground application equipment, direct the spray toward crop foliage.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls worn over long-sleeved shirt and long pants.
- Chemical-resistant gloves such as barrier laminate or butyl rubber ≥14 mils or nitrile rubber ≥14 or neoprene rubber ≥14 mils or natural rubber ≥14 or polyethylene or polyvinyl chloride (PVC) ≥14 mils or Viton > 14 mils
- Chemical-resistant footwear plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY AND/OR POOR INSECT CONTROL.

NOT FOR SALE, SALE INTO, DISTRIBUTION AND/OR USE IN NASSAU AND SUFFOLK COUNTIES, NEW YORK.

PRODUCT INFORMATION

Trigard Insecticide is a 75% wettable powder. When this product is mixed with water and applied according to the directions for use to the specified crops on this label, Trigard Insecticide will control the specified insects in the crops that are listed on this label.

Resistance Management

For resistance-management, Trigard Insecticide contains a Group 17 insecticide (contains the active ingredient cyromazine). Any insect population may contain individuals naturally resistant to Trigard Insecticide or other Group 17 insecticides. The resistant individuals may dominate the insect population if these groups of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed. To reduce the potential for developing insect resistance, rotate to an insecticide with a different mode of action. Monitor treated pest populations for resistance development. Read the product label before applying any insecticide and follow label directions.

To delay insecticide resistance, take the following steps:

- Rotate the use of Trigard Insecticide or other Group 17 insecticides within a growing season, or among growing seasons, with different groups that control the same pests. Avoid application of more than the maximum seasonal use rate or the total number of consecutive sprays of Trigard Insecticide per season.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):

oIndividual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.

oMixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.

oWhen using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).

oMixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.

oThe insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.

- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.

- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.

- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.

- For further information or to report suspected resistance contact Gowan Company, LLC representatives at 1-800-883-1844.

MIXING AND APPLICATION INSTRUCTIONS

- 1) Thoroughly clean spray equipment before using this product.
- 2) Prepare no more spray mixture than is needed for the immediate operation.
- 3) Agitate spray mixture thoroughly before and during application. Good bypass agitation is sufficient to keep Trigard Insecticide thoroughly mixed.
- 4) Flush spray tank thoroughly with water daily after use and
- 5) Dispose of pesticide rinsate by application to an already treated area.

Instructions for Using Water Soluble Packages Directly into Spray tanks: Water Soluble Packages (WSPs) are designed to dissolve in water. Agitation may be used, if necessary, to help dissolve the WSP. Failure to follow handling and mixing instructions can increase your exposure to the pesticide products in WSPs. WSPs, when used properly, qualify as a closed mixing/loading system under the Agricultural Worker Protection Standard [40 CFR 170.607(d)].

Handling Instructions:

Follow these steps when handling pesticide products in WSPs.

1. Mix in spray tank only.
2. Handle WSP(s) in a manner that protects package from breakage and/or unintended release of contents. If package is broken, put on PPE required for clean-up and then continue with mixing instructions.
3. Keep the WSP(s) in outer packaging until just before use.

4. Keep the WSP dry prior to adding to the spray tank.
5. Handle with dry gloves and according to the label instructions for PPE.
6. Keep WSP intact. **DO NOT** cut or puncture WSP.
7. Reseal the WSP outer packaging to protect any unused WSP(s).

Mixing Instructions:

Follow the steps below when mixing this product, including if tank mixed with other pesticide products. If being tank mixed, the mixing directions 1 through 9 below take precedence over the mixing directions of the other tank mix products. WSPs may, in some cases, be mixed with other pesticide products so long as the directions for use of all mixed products do not conflict. **DO NOT** tank mix this product with products that prohibit tank mixing or have conflicting mixing directions.

1. If a basket or strainer is present in the tank hatch, remove prior to adding the WSP to the tank.
2. Fill tank with water to approximately one-third to one-half of the desired final volume of spray.
3. Stop adding water and stop any agitation.
4. Place intact/unopened WSP(s) into the tank.
5. **DO NOT** spray water from a hose or fill pipe to break or dissolve the WSP(s).
6. Start mechanical and recirculation agitation from the bottom of tank without using any overhead recirculation, if possible. If overhead recirculation cannot be turned off, close the hatch before starting agitation.
7. Dissolving the WSP(s) may take up to 5 minutes or longer, depending on water temperature, water hardness and intensity of agitation.
8. Stop agitation before tank lid is opened.
9. Open the lid to the tank, exercising caution to avoid contact with dusts or spray mix, to verify that the WSPs have fully dissolved and the contents have been thoroughly mixed into the solution.
10. **DO NOT** add other allowed products or complete filling the tank until the bags have fully dissolved and pesticide is thoroughly mixed.
11. Once the WSP have fully dissolved and any other products have been added to the tank, resume filling the tank with water to the desired level, close the tank lid, and resume agitation.
12. Use the spray solution when mixing is complete.
13. Maintain agitation of the diluted pesticide mix during transport and application.
14. It is unlawful to use any registered pesticide, including WSPs, in a manner inconsistent with its label.

Trigard Insecticide Alone:

- 1) Add 2/3 of the required amount of water to the spray or mixing tank.
- 2) With the agitator running, drop the required number of unopened soluble packets of Trigard Insecticide into the tank. (One 2.66 oz. water-soluble packet of Trigard Insecticide will treat one acre).
- 3) Continue agitation while adding the remainder of the water.

Spray Drift Requirements

Droplet Size

When making foliar applications of this product (aerial or ground), use spray nozzles that will deliver medium course or larger spray droplets as defined in the American Society Agricultural and Biological Engineers (ASABE) standard ANSI/ASAE S572.1 (March 2009). In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

CROP USE DIRECTIONS

Beans, Dried Varieties (Except Cowpeas and Soybeans) [*]:		
Adzuki beans, black-eyed peas, broad beans (dry), catjang, chick peas, Crowder peas, field beans, garbanzo beans, grain lupin, guar, kidney beans, lablab beans, lentils, lima beans (dry), moth beans, mung beans, navy beans, pinto beans, rice beans, southern peas, tepary beans, sweet lupin, urd beans, wax beans, white lupin, white sweet lupin		
Target Pests	Rates per Acre	Instructions
Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray when leafminers first appear. Repeat application up to 6 times to maintain control.
Use Restrictions for Beans, Dried Varieties (Except Cowpeas and Soybeans): <ul style="list-style-type: none"> • Pre-Harvest Interval: Allow a minimum of 7 days between the last application and harvest. • Application Method: Ground or aerial application equipment is permitted. • Amount of Water: Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage. • Retreatment Interval: Allow a minimum of 7 days between applications. • Maximum Amount: <ul style="list-style-type: none"> • DO NOT apply more than 0.125 lb a.i./A per 7-day interval. • DO NOT apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year. • DO NOT apply more than 6 times per crop season. <p>[*Not registered for use by California]</p>		

Beans, Succulent [*]: Bean (<i>Phaseolus</i> spp.) Broad bean, succulent; Lima bean, green; Runner bean; Snap bean (green bean); Wax bean Bean (<i>Vigna</i> spp.) Asparagus bean; Black-eyed pea; Chinese longbean; Cowpea; Jackbean; Moth bean; Southern pea; Yardlong bean		
Target Pests	Rates per Acre	Instructions
Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray when leafminers first appear. Repeat application up to 6 times to maintain control.
Use Restrictions for Beans, Succulent: <ul style="list-style-type: none"> • Pre-Harvest Interval: Allow a minimum of 7 days between the last application and harvest. • Application Method: Ground or aerial application equipment is permitted. • Amount of Water: Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage. • Retreatment Interval: Allow a minimum of 7 days between applications. • Maximum Amount: <ul style="list-style-type: none"> • DO NOT apply more than 0.125 lb a.i./A per 7-day interval. • DO NOT apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year. • DO NOT apply more than 6 times per crop season. [*Not registered for use by California]		

Pea, Succulent [*]: Edible podded pea legume vegetables: dwarf pea; edible podded pea; green pea; snap pea; snow pea; sugar snap pea; grass-pea; lentil; pigeon pea; chickpea; cultivars, varieties, and/or hybrids of these commodities. Succulent shelled peas: Chickpea; English pea; garden pea; green pea; pigeon pea; lentil; cultivars, varieties, and/or hybrids of these commodities.		
Target Pests	Rates per Acre	Instructions
Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray when leafminers first appear. Repeat application up to 6 times to maintain control.
Use Restrictions for Pea, Succulent: <ul style="list-style-type: none"> • Pre-Harvest Interval: Allow a minimum of 7 days between the last application and harvest. • Application Method: Ground or aerial application equipment is permitted. • Amount of Water: Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage. • Retreatment Interval: Allow a minimum of 7 days between applications. • Maximum Amount: <ul style="list-style-type: none"> • DO NOT apply more than 0.125 lb a.i./A per 7-day interval. • DO NOT apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year. • DO NOT apply more than 6 times per crop season. [*Not registered for use by California]		

Vegetable, Brassica, head and stem, group 5-16, except broccoli; Brassica, leafy greens, subgroup 4-16B [*]: Brussels sprouts; cabbage; cabbage, Chinese, napa; cauliflower; cultivars, varieties, and hybrids of these commodities. Arugula; broccoli, Chinese; broccoli raab; cabbage, abyssinian; cabbage, Chinese, bok choy; cabbage, seakale; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; watercress; cultivars, varieties, and hybrids of these commodities. Kohlrabi		
Target Pests	Rates per Acre	Instructions
Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray when leafminers first appear. Repeat application up to 6 times to maintain control.
Use Restrictions for Vegetable, Brassica, head and stem, group 5-16, except broccoli; and Brassica, leafy greens, subgroup 4-16B: <ul style="list-style-type: none"> • Pre-Harvest Interval: Allow a minimum of 7 days between the last application and harvest. • Application Method: Ground or aerial application equipment is permitted. • Amount of Water: Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage. • Retreatment Interval: Allow a minimum of 7 days between applications. • Maximum Amount: <ul style="list-style-type: none"> • DO NOT apply more than 0.125 lb a.i./A per 7-day interval. • DO NOT apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year. • DO NOT apply more than 6 times per crop season. • Cyromazine may not be applied when watercress is flooded. Water must be turned off 24 hours in advance of an application, and may be turned back on 24 hours after the application. [*Not registered for use by California]		

Broccoli [*]:		
Target Pests	Rates per Acre	Instructions
Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray when leafminers first appear. Repeat application up to 6 times to maintain control.
Use Restrictions for Broccoli: <ul style="list-style-type: none"> • Pre-Harvest Interval: Allow a minimum of 7 days between the last application and harvest. • Application Method: Ground or aerial application equipment is permitted. • Amount of Water: Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage. • Retreatment Interval: Allow a minimum of 7 days between applications. • Maximum Amount: <ul style="list-style-type: none"> • DO NOT apply more than 0.125 lb a.i./A per 7-day interval. • DO NOT apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year. • DO NOT apply more than 6 times per crop season. • Cyromazine may not be applied when watercress is flooded. Water must be turned off 24 hours in advance of an application, and may be turned back on 24 hours after the application. [*Not registered for use by California]		

Onion, bulb, subgroup 3-07A and Onion, green, subgroup 3-07B [*]:		
Daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, greatheaded, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these. Chive, fresh leaves; chive, Chinese, fresh leaves; elegans hosta; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.		
Target Pests	Rates per Acre	Instructions
Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray when leafminers first appear. Repeat application up to 6 times to maintain control.
Use Restrictions for Onion, bulb, subgroup 3-07A and Onion, green, subgroup 3-07B: <ul style="list-style-type: none"> • Pre-Harvest Interval: DO NOT make the last application within 7 days of harvest (or cutting). • Application Method: Ground or aerial application equipment is permitted. • Amount of Water: Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage. • Retreatment Interval: Allow a minimum of 7 days between applications. • Maximum Amount: <ul style="list-style-type: none"> • DO NOT make more than 6 applications to one crop. • DO NOT apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year. • DO NOT apply more than 6 times per crop season. [*Not registered for use by California]		

Cucurbit Vegetables Crop Group 9 [*]:		
Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); Momordica spp (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes cantaloupe); pumpkin; squash, summer; squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon.		
Target Pests	Rates per Acre	Instructions
Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray when leafminers first appear. Repeat application up to 6 times to maintain control.
Use Restrictions for Cucurbit Vegetables Crop Group 9: <ul style="list-style-type: none"> • Pre-Harvest Interval: Cucurbits may be harvested on the same day of the last application. (0-day PHI) • Application Method: Ground or aerial application equipment is permitted. • Amount of Water: Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage. • Retreatment Interval: Allow a minimum of 7 days between applications. • Maximum Amount: <ul style="list-style-type: none"> • DO NOT apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year. • DO NOT apply more than 6 times per crop season. [*Not registered for use by California]		

Leafy greens subgroup 4-16A and Leaf petiole vegetable subgroup 22B [*]:

Leafy greens subgroup including: Amaranth, Chinese; amaranth, leafy; aster, Indian; blackjack; cat's whiskers; cham-chwi; cham-na-mul; chervil, fresh leaves; chipilin; chrysanthemum, garland; cilantro, fresh leaves; corn salad; cosmos; dandelion, leaves; danggwi, leaves; dill-weed; dock; dol-nam-mul; ebolo; endive; escarole; fleamflower; feather cockscomb; good king henry; huauzontle; jute, leaves; lettuce, bitter; lettuce, head; lettuce, leaf; orach; parsley, fresh leaves; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; spinach; spinach, Malabar; spinach, New Zealand; spinach, tanier; Swiss chard; violet, Chinese, leaves; cultivars, varieties, and hybrids of these commodities.

Leafy greens subgroup including: Cardoon; celery; celery, Chinese; fuki; rhubarb; udo; zuiki; cultivars, varieties, and hybrids of these commodities. Also includes Celtuce, and Florence fennel.

Target Pests	Rates per Acre	Instructions
Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray when leafminers first appear. Repeat application up to 5 times to maintain control. May repeat application up to 6 times to maintain control in celery and head lettuce.

Use Restrictions for Leafy greens subgroup 4-16A and Leaf petiole vegetable subgroup 22B [*]:

- **Pre-Harvest Interval:** DO NOT make the last application within 7 days of harvest (or cutting).
- **Application Method:** Ground or aerial application equipment is permitted.
- **Amount of Water:** Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage.
- **Retreatment Interval:** Allow a minimum of 7 days between applications.
- **Maximum Amount:**
- **DO NOT** apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year.
- **DO NOT** make more than 6 applications per crop of celery or head lettuce.
- **DO NOT** apply more than 5 applications for all other leafy vegetable crops.

[*Not registered for use by California]

Pepper/Eggplant subgroup 8-10B [*]:

African eggplant; bell pepper; eggplant; martynia; non-bell pepper; okra; pea eggplant; pepino; roselle; scarlet eggplant; cultivars, varieties, and/or hybrids of these.

Target Pests	Rates per Acre	Instructions
Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray when leafminers first appear. Repeat application up to 6 times to maintain control.

Use Restrictions for Pepper/Eggplant subgroup 8-10B:

- **Pre-Harvest Interval:** Peppers may be harvested on the same day of the last application (0- day PHI)
- **Application Method:** Ground or aerial application equipment is permitted.
- **Amount of Water:** Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage.
- **Retreatment Interval:** Allow a minimum of 7 days between applications.
- **Maximum Amount:**
- **DO NOT** make more than 6 applications per crop.
- **DO NOT** apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year.

[*Not registered for use by California]

Vegetable, tuberous and corm, subgroup 1C [*]:

Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; potato; sweet potato; tanier; turmeric; yam bean; yam, true.

Target Pests	Rates per Acre	Instructions
Colorado potato beetle control	2.66 oz. (one packet) per A or 5.32 oz. (two packets) per A	Trigard Insecticide is an insect growth regulator and is most effective for control of first and second instar larvae of Colorado potato beetles. Although larger larvae can be affected, treatment of early instars will give best control. Affected larvae may remain on the plant for several days after treatment, but immediately stop feeding. • Trigard Insecticide does not control adult beetles. Apply 2.66 oz. (one packet) or 5.32 oz. (two packets) per acre as a foliar spray. The first application should be made just after egg hatch to early instar larvae. If a second application for the same generation is needed, apply 2.66 oz. (one packet) or 5.32 oz. (two packets) per acre 7-14 days later.
Leafminer	2.66 oz. (one packet) per A	Repeat application up to 7 times to maintain control.

Use Restrictions for Vegetable, tuberous and corm, subgroup 1C:

- **Pre-Harvest Interval:** Allow a minimum of 17 days between the last application and harvest (17-day PHI).
- **Application Method:** Ground or aerial application equipment is permitted.
- **Amount of Water:** Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage.
- **Retreatment Interval:** Allow a minimum of 7 days between applications.
- **Maximum Amount:**
- **DO NOT** apply more than 1lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year.

[*Not registered for use by California]

Tomato subgroup 8-10A [*]:

Bush tomato; cocona; currant tomato; garden huckleberry; goji berry; groundcherry; naranjilla; sunberry; tomatillo; tomato; tree tomato; cultivars, varieties, and/or hybrids of these.

Target Pests	Rates per Acre	Instructions
Colorado potato beetle control and Leafminer	2.66 oz. (one packet) per A	Apply as a foliar spray

Use Restrictions for Tomato subgroup 8-10A:

- **Pre-Harvest Interval:** Tomatoes may be harvested on the same day of the last application (0-day PHI).
- **Application Method:** Ground or aerial application equipment is permitted.
- **Amount of Water:** Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage.
- **Retreatment Interval:** Allow a minimum of 7 days between applications.
- **Maximum Amount:**
DO NOT apply more than 1 lb (0.75 lb. a.i./A) of Trigard Insecticide per acre per calendar year.
DO NOT make more than 6 applications to one crop of tomatoes.

[*Not registered for use by California]

ROTATIONAL CROPS

The following "plant-back intervals" must be observed when planting rotational crops after an application of Trigard Insecticide:

Rotational Crop	Planting Time From Last Application of Trigard Insecticide (Days)
All crops on the Trigard Insecticide label	0 days
Sweet corn, radishes, and cotton	30 days
Any other crop	300 days

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. **DO NOT** use in animal feeds.

PESTICIDE STORAGE: Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. **DO NOT** store this product under wet conditions. Handle outer container carefully to avoid breakage of water-soluble packets.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal is a violation of Federal law. To avoid harming aquatic organisms in rivers and other surface waters, do not pour spray mixture or rinse water into sanitary drains (for example; toilets, floor drains, and sinks) or into storm water sewers (for example; street drains). If pesticide, spray mixture, or rinsate cannot be disposed of by use according to the label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable Container (flexible-bag-all weights): Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Once all watersoluble packets are used according to label instructions, dispose of empty outer bag in a sanitary landfill or by other procedures allowed by State and local authorities.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC (800) 424-9300
For other product information, contact Gowan Company, LLC or See Safety Data Sheet

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User.

Gowan Company, LLC warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY, LLC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY, LLC'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY LLC'S SOLE DISCRETION.

Trigard® is a registered trademark of Gowan Company, LLC.

All other brands are registered trademarks of their respective owners.

[To EPA 06-23-2025, 09-24-2025, 10-14-2025]