

**PRECAUTIONARY STATEMENTS
DANGER**

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension or Service.

This Product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do Not apply where runoff is likely to occur. Do Not apply when weather conditions favor drift from areas treated. Do Not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and Crabs may be killed at application rates recommended on this label. Do Not apply where these are important resources. Apply this product only as specified on this label. **DO NOT APPLY** in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do Not use in serving areas while food is exposed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label. Use strictly in accordance with cautions, warnings and directions, and with applicable state and federal regulations. Timing may vary as a consequence of local weather or conditions; we recommend reference to Agricultural Experiment Stations on specific timing.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

PROKIL NALED is a liquid insecticide for the control of both sucking and crawling insects. It may be applied as a conventional diluted spray or as a concentrate spray according to directions. The normal requirements for the following application methods are as follows: Vegetable and Field Crops: Air 7-15 gals. per acre; Ground 20-200 gals. per acre. Small Fruits: Air 7-15 gals. per acre; Ground 20 to 200 gals. per acre. Tree Crops (mature): Air 10-20 gals. per acre; Ground 40-200 gals. per acre. Citrus (mature): Air 10-20 gals. per acre; Ground 100-1500 gals. per acre.

May be applied up to 4 days before harvest, branding or hand picking unless otherwise noted. Begin application at first sign of insects. Repeat as necessary. Do not use with highly alkaline materials such as lime or bordeaux. **MIXING DIRECTIONS:** Add to spray tank when 1/2 to 3/4 full with agitator running and maintain agitation until spraying is completed.

USE DESIGNATED RATE BY GROUND EQUIPMENT AS A THOROUGH COVERAGE SPRAY: ALFALFA (Seed): Lygus, Mite Suppression (1 application) 1 pt. per acre. Mites 1-2 pints per acre; 2 applications at 7 day intervals. Thrips 1 pt. per acre. Combine with recommended chemicals for residual control. **DO NOT** apply within 4 days of harvest.

BEANS (Dry or Field): SOYBEANS, LIMA BEANS AND PEAS for Processing: Alfalfa Looper, Aphids, Spider Mites, Lygus, Leaf Miner; **GREEN BEANS:** Aphids, Spider Mites,

**PROKIL
NALED INSECTICIDE**

DANGER

KEEP OUT OF REACH OF CHILDREN

DANGER

Concentrate causes skin and eye damage. Do not get on skin, eyes or clothing. Use water proof gloves and face shield or goggles when handling concentrate. May be fatal if swallowed or absorbed through the skin. Do not breathe spray mist. In case of contact immediately remove contaminated clothing and flush skin or eyes with plenty of water. For eyes, use special attention. Avoid contact with food, feed, milk, and water. **DO NOT USE FOR SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Use only in accordance with label directions. Liquidambar of China is the active ingredient.

NOTE TO PHYSICIAN: ATROPINE IS ANTIDOTAL. 2-PAM IS ALSO ANTIDOTAL AND MAY BE ADMINISTERED IN CONJUNCTION WITH ATROPINE.

Gowan Company

AGRICULTURAL CHEMICALS


12300 E. 8th Street

P.O. Box 5696

Telephone 783-8844

Tucson, AZ. 85364

EPA Reg. No. 10163-46 AA
EPA Est. No. 10163-AZ-1



Loopers, Stink Bugs, Leafhoppers, etc. Large Loopers, Stink Bugs, BROCCOLI, CABBAGE, CAULIFLOWERS, Imported Cabbageworm, Looper, 2 pts. per acre. May be a heavy insecticide. Sprouts and K. COTTON, Lygus, Leaf Perforator, Suppression of Mites, 1 pt. per acre. Heavy Insect or Egg Population of any for best results. Do not apply. CUCUMBERS, SUMMER SQUASH, Thrips, Armyworms 1 pt. per acre. Water to cover. Aerial Application up to 1 day of harvest. EGGPLANT, PEPPERS: Aphids, Whitefly 1 pt. per acre. May be applied. WALNUTS: Aphids 1 to 1 pt. apply within 10 days of harvest. GRAPES: Drosophila, Leafhopper spray per acre depending on size variety. Do not use on Italia variety. HOPS: Aphids, Spider Mites, Control within 4 days of harvest. CHERRY, SPINACH, CHARD: Marsh Caterpillars, Leaf Miners 50 total gals. water per acre on pines, total gals. water per acre. May be applied. MELONS (Cantaloupes, Honeydew, SQUASH: Loopers, Armyworms - if temperature is over 90° F. On Honeydew, Do not exceed 2 pints per acre. Do not prevent injury from spray applications. ORANGES, LEMONS, GRAPEFRUIT: Citrus Cutworms, Leaf Roller (1 or 2) 100 gals. water. Do not apply within 4 days of harvest in California and Arizona. PEACHES (Western States Only): Fruit Moth (California Only) - Do not exceed 2 pints per acre. Do not prevent fruit spotting, do not use over 90° F. RICE: Grasshoppers - 1 to 1 1/2 pints per acre. Do not use more than 3 applications. Do not use. ROSES (Field Grown): Aphids, Mites - water. STRAWBERRIES: Leaf Roller, Spider Mites - 1 pt. per acre. Northwest 1 pt. per acre. May be applied. SUGAR BEETS: Spider Mites, Leafhoppers. TOBACCO: Hornworm 1 pt. per acre. Flea Beetles 1 pt. per acre if necessary. TOMATOES (Field) Fruit Flies - 1 with ground sprayer, 2 to 10 gals. per acre and continue at 7 to 7 day intervals. Worm, Hornworms, Leafminers - 1 thoroughly. For Field Use up to 100° F. May be applied.