

ACCEPTED
 MAY 17 1973
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND ROBBITICIDE ACT
 FOR ECONOMIC POISON REGISTER
 ED UNDER NO. 10170-7
 TO ATTACHED COMMENTS

USE DIRECTIONS

Laundry: Use 1/2 ounce **SEPTICIDAL S-10** per 15 pounds of laundry in the Rinse Cycle.

Household: For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classroom, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce **SEPTICIDAL S-10** to 4 gallons of water. Disinfect with 3 ounces **SEPTICIDAL S-10** to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavy soiled or contaminated areas, a precleaning step is recommended.

Hospitals: For floors, walls and other hard surfaces. In Nursing Homes, Institutions. On sink tops, garbage pails, telephones and restrooms. For best results, first clean all surfaces with a suitable detergent, to remove gross filth and heavy soil deposits, then mop dry and treat with a solution of **SEPTICIDAL S-10** at a rate of 3 1/2 ounces per 5 gallons of water. Floors may be either mopped or scrubbed, other surfaces may be either wiped or scrubbed. At this level **SEPTICIDAL S-10** will be effective against Pseudomonas aeruginosa PRD-10.

Food Processing Equipment: For sanitizing of previously cleaned food processing equipment and food utensils dilute 1 ounce **SEPTICIDAL S-10** per 4 gallons of water to provide 200 ppm active sanitizer. At this level, a Potable Water Rinse is Not Required.

Dairies: To sanitize dairy equipment, such as tanks, lines, pails and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce **SEPTICIDAL S-10** to 4 gallons of water. At this level, a Potable Water Rinse is Not Required. Follow recommendations of local Health Board.

SEPTICIDAL S-10

DISINFECTANT • SANITIZER • DEODORIZER

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
 dimethyl benzyl ammonium chlorides 5%
 n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl ethylbenzyl
 ammonium chlorides 5%

INERT INGREDIENTS: 90%

SEPTICIDAL S-10 is a concentrate and should be diluted before using.

CONTENTS: 1 Gallon **EPA Reg. No. 10170-7**

DANGER: Keep Out of Reach of Children. See First Aid statement and other precautions on right panel.

Residual Bacteriostatic and Residual Self-Sanitizing Under Conditions of High Relative Humidity or Wet Contamination.

Fabric treated as directed will develop residual bacteriostatic activity and residual self-sanitizing activity against Gram positive (S. aureus) and Gram negative (E. coli) organisms, under conditions of high relative humidity or wet contamination.

SEPTICIDAL S-10 prevents growth of ammonia producing organisms (B. ammoniagenes) which act on urine in the diapers to produce ammonia.

Manufactured by
MIDWEST BIOCHEMICAL CORP.
 Milwaukee, Wisconsin 53205

Restaurant and Bar Rinse: To sanitize dishes, glassware, silverware, and cooking utensils, first wash with soap or synthetic detergent, then rinse thoroughly and immerse in a sanitizing solution containing 1 ounce **SEPTICIDAL S-10** to 4 gallons of water. No Terminal Potable Water Rinse Required At This Level.

Cold Disinfection: For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into a solution containing 1 1/2 ounces **SEPTICIDAL S-10** per gallon of water for 10 minutes.

DANGER - Corrosive: Keep Out of Reach of Children. Undiluted solutions causes severe eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear rubber gloves and eye mask when handling. Harmful or fatal if swallowed from container. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.