

DOCKTOR'S[®]

DOCKTOR'S[®]

LICE AND MITE SPRAY

DIRECTIONS: Shake gently before use. Spray parakeets, canaries and other caged birds as follows: For cage mite control, remove bird and spray the entire cage thoroughly. For bird lice, leave bird in cage and spray directly on bird holding unit at least two feet from bird. Spray lightly—one burst for 2 or 3 seconds. For control apply no more than two times per week. Remove food from cage before spraying. Bird cages should be cleaned in hot water on a weekly basis.

ACTIVE INGREDIENTS

Pyrethrins	0.090%
Technical Piperonyl Butoxide*	0.180%
N-Octyl bicycloheptene dicarboximide	0.300%
Petroleum Distillates	0.430%
Triethylene Glycol	0.075%
Propylene Glycol	0.075%
INERT INGREDIENTS	98.850%
	100.000%

*Consists of 0.144% (butylcarbityl 6-pylopyperonyl) ether and 0.036% related compounds.

USDA REG. NO. 10173-8

CAUTION: Contents under pressure. Do not puncture or incinerate can. Don't use near heat or store where temperature can exceed 120° F. Cover fish and turtle bowls when spraying.

ACCEPTED
 Feb-5-1960
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 10173-8



Net weight—7 oz. Avoir

C.G. 10/69

B216

Manufactured for
DOCKTOR'S LABORATORIES, INC.
 VETERINARY PRODUCTS DIVISION
 Philadelphia, Pennsylvania 19124
 Made in U.S.A.

To rid parakeets, canaries and all caged birds of lice and mites.

Caution: Keep out of Reach of Children
See back panel for additional cautions.

DOCKTOR'S®

DOCKTOR'S®

LICE AND MITE SPRAY

DIRECTIONS: Shake gently before use. Spray parakeets, canaries and other caged birds as follows: For cage mite control, remove bird and spray the entire cage thoroughly. For bird lice, leave bird in cage and spray directly on bird holding unit at least two feet from bird. Spray lightly—one burst for 2 or 3 seconds. For control apply no more than two times per week. Remove food from cage before spraying. Bird cages should be cleaned in hot water on a weekly basis.

ACTIVE INGREDIENTS

Pyrethrins	0.090%
Technical Piperonyl Butoxide*	0.180%
N-Octyl bicycloheptene dicarboximide	0.300%
Petroleum Distillates	0.430%
Triethylene Glycol	0.075%
Propylene Glycol	0.075%
INERT INGREDIENTS	98.850%
	100.000%

*Consists of 0.144% (butylcarbityl 6-pyropiperonyl) ether and 0.036% related compounds.

USDA REG. NO. 10173-8

CAUTION: Contents under pressure. Do not puncture or incinerate can. Don't use near heat or store where temperature can exceed 120° F. Cover fish and turtle bowls when spraying.

ACCEPTED
 Feb-5-1970
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 10173-8



Net weight—7 oz. Avoir

C.G. 10/69

B216

Manufactured for
DOCKTOR'S LABORATORIES, INC.
 VETERINARY PRODUCTS DIVISION
 Philadelphia, Pennsylvania 19154
 Made in U.S.A.

To rid parakeets, canaries and all caged birds of lice and mites.
Caution: Keep out of Reach of Children
 See back panel for additional cautions.