

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

**HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. MAY CAUSE EYE IRRITATION.**

Avoid contact with eyes, skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly after handling.

**STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED:** Do not induce vomiting. Call a physician. Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages via aspiration of the solvent. **IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Wash with plenty of soap and water.

For 24-hour emergency assistance call 302/575-3000.

**ENVIRONMENTAL HAZARDS:** This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, streams, ponds, tidal marshes or estuaries. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply AMBUSH or allow it to drift to crops or weeds on which bees are actively foraging. Additional information may be obtained from your Cooperative Extension Service. Apply this product only as specified on this label.

In case of a significant spill, call CHEMTREC 800/424-9300.

**PHYSICAL OR CHEMICAL HAZARDS:** Do not use or store near heat or open flame.

**STORAGE AND DISPOSAL**

Keep container closed when not in use. Do not store near food or feed. Protect from freezing.

In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area and dispose of in accordance with State and local regulations.

Do not reuse container. Triple rinse residue from empty container and add rinse to mixture in spray tank. Dispose of container in sanitary landfill in accordance with State and local regulations.

**NOTICE TO BUYER AND USER:** Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer and User assumes the risk of any such use. Seller disclaims all other warranties expressed or implied including any warranty of fitness or merchantability. Seller shall not be liable for consequential special or indirect damages resulting from the use or handling of this product and Seller's sole liability and Buyer's and User's exclusive remedy shall be limited to the refund of the purchase price.

**RESTRICTED USE PESTICIDE.**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

# Ambush<sup>®</sup>

## INSECTICIDE

**Keep Out  
of Reach  
of Children  
WARNING**

See Side Panel  
For Additional  
Precautionary  
Statements and  
Statement of  
Practical  
Treatment

**ACTIVE INGREDIENT:**

Permethrin—

(3-Phenoxyphenyl) methyl (±)-cis, trans-3-(2,2-dichloroethenyl)-  
2,2-dimethylcyclopropanecarboxylate\* ..... 25.6%

**INERT INGREDIENTS** ..... 74.4%

**Total** ..... 100%

\*Cis/trans ratio: Min. 35% (±) cis and max. 65% (±) trans.

AMBUSH contains 2 pounds active ingredient per gallon

US Patent No. 4,024,163



ICI Americas Inc.

Agricultural Chemicals Division

Wilmington, Delaware 19897

EPA Reg. No. 10182-18

EPA Est. No. 5905-AR-1

Net Contents:  
**ONE U.S. GALLON**  
(3.78 LITERS)

Batch No.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply AMBUSH to cotton as shown in the chart below, using sufficient spray TO OBTAIN FULL COVERAGE OF FOLIAGE. The higher rates of AMBUSH (0.15 to 0.2 lbs/acre) may be required under conditions of heavy lepidopterous worm infestation, or when boll weevils, or late instars of larvae are present. AMBUSH aids in the reduction of lepidopterous egg hatch by contact action. Timing and frequency of spraying should be based upon insect populations reaching local economic thresholds. Apply as required by scouting reports, usually at intervals of 5 to 7 days. Do not apply within 14 days of harvest.

### COTTON SPRAY RECOMMENDATIONS

Target Pests	Dosage Per Acre	Minimum Spray Volume Per Acre
Boll weevil*		
Boilworm		Minimum of
Tobacco Budworm	0.4	1.5 gal. by
Pink Bollworm	to	air or
Lygus Bugs	0.8	minimum of
Whitefly*	pint	5 gal. by
Cotton Aphid*		ground
Cabbage Looper		
Cotton Leafperforator		

### RATE CONVERSION CHART


Lb. Active/Acre	Fluid Ozs./Acre	Pints/Acre	Acres Treated/Gal.
0.10	6.4	0.4	20.0
0.125	8.0	0.5	16.0
0.15	9.6	0.6	13.3
0.175	11.2	0.7	11.4
0.20	12.8	0.8	10.0

#### Remarks

Do not graze livestock in treated areas.  
Do not apply more than 12 pints per acre per season (3 lbs ai/Acre).

Do not plant any crop except cotton within 60 days after last application. Any crop may be planted if it is not harvested or grazed.

\*Only suppression can be expected when utilized on a 5-7 day schedule.



2012