

10819  
PM25  
10/82-18

OCT 11 1984

BEST AVAILABLE COPY

ICI Americas Inc.  
Concord Pike & New Murphy Road  
Wilmington, DE 19877

Dear Mr. R.E. Ridsdale

Subject: Ambush Insecticide  
EPA Reg. No. 10182-18  
Your letter of September 29, 1984

Your proposed amendment to add a claim for use of the product on tomatoes is acceptable subject to the comment listed below and a stamped copy of the draft label is enclosed for your records. Submit five (5) copies of the finished printed label bearing the labeling restriction: "Only for Application to Tomatoes Grown in Florida for Final Marketing as Fresh Tomatoes" before the product is released for shipment with this amended labeling.

Sincerely

*fr*  
Timothy A. Gardner  
Product Manager 17  
Insecticide Rodenticide Branch  
Registration Division TS-767

IRB:CM2:TAG:JMT:pl: RM207:

10/15/84

ACCEPTED  
WITH COMMENTS  
in EPA Letter Dated:

OCT 15 1984

U.S. Environmental Protection Agency  
Registration Division  
Pesticide Division  
Washington, D.C. 20460  
10182-18

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. MAY CAUSE EYE IRRITATION.

Avoid contact with eyes, skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED: Do not induce vomiting. Call a physician. Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damage via aspiration of the solvent. IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists. IF ON SKIN: Wash with plenty of soap and water.

For 24-hour emergency assistance call 800/575-3000

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply AMBUSH, or allow it to drift to crops or weeds on which bees are actively foraging. Additional information may be obtained from your Cooperative Extension Service. Apply this product only as specified on this label.

In case of a significant spill call CHEMTREC 800/424-9300


PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.

## STORAGE AND DISPOSAL

Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, wash up with sand, earth, or synthetic absorbent. Remove to chemical waste area and dispose of in accordance with State and local regulations. Do not reuse container. Triple rinse residual from empty container and add rinse to mixture in spray, tank, pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures. Then after empty container for recycling or reconditioning, or purchase and dispose of in a sanitary landfill, or by other approved State and local procedures.

## RESTRICTED USE PESTICIDE

This pesticide is restricted to use only by certified applicators, or persons under their direct supervision, and only for those uses registered for this Certified Applicator's certification.



Keep Out  
of Reach  
of Children  
WARNING

See Side Panel For Additional  
Precautionary Statements and  
Statement of Practical  
Treatment

EPA Est. No. 5906-AR-1  
EPA Reg. No. 10102-10

Net Contents:  
ONE U.S. GALLON  
(3.78 LITERS)  
Batch No.

ICI Americas Inc.  
Pesticide Division

U.S. Patent No. 4,064,103

## ACTIVE INGREDIENT:

Permethrin-

(3-Phenoxycarbonyl) methyl (±)-cis, trans-3-(2,2-dichloroethoxy)-  
2,2-dimethylcyclopropanecarboxylate

INERT INGREDIENTS 74.4%  
TOTAL 100.0%

\*Containing either Min. 35% (±) cis and max. 65% (±) trans.  
AMBUSH contains 2 pounds active ingredient per gallon.

It is a violation of Federal law  
to apply AMBUSH\* as indicated  
on this Supplemental Pamphlet.

Pounds Active/ Gals	RAT
0.05	
0.10	
0.125	
0.15	
0.175	
0.20	

NOTICE TO BUYER:  
chemical description on the  
label in accordance with the  
order to the use of this  
product, or under which  
assumes the risk of any  
EXPRESSED OR IMPLIED  
ABILITY, SELLER SHALL IN  
DAMAGES RESULTING FROM  
SOLE LIABILITY AND BUYER  
THE REFUND OF THE PUR

BEST AVAILABLE COPY

Crop	Target Pest(s)	lb ai/A	Comments
Asparagus	Cutworm complex Asparagus beetle	0.05-0.1	Apply by ground equipment using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 4 applications. Do not apply within 3 days of harvest.
Asparagus ferns	Japanese beetle Asparagus beetle Lygus bugs	0.1	Apply to the fern stage of the asparagus plant after spear harvest when larval and adult stage of the asparagus beetle and lygus bugs and the adult stage of the Japanese beetle are present.

ACCEPTED

OCT 16 1984

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the purpose of  
registered under  
EPA Reg. No. 10182-18

ATVISC FEB 06 1984

**RESTRICTED USE PESTICIDE.**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.


**SUPPLEMENTAL LABEL FOR USE ON  
FIELD CORN/POPCORN****AMBUSH® Insecticide****EPA Reg. No: 10182-18****DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product  
in a manner inconsistent with its labeling**

Apply AMBUSH® Insecticide as shown in the following charts. Use sufficient water to obtain full coverage of foliage. Timing and frequency of applications should be based upon insect populations reaching economic thresholds. The higher rate may be required under conditions of heavy infestation.

**DIRECTIONS FOR APPLICATION**

Crop	Target Pest(s)	Rate (lbs. a.i./A)	Remarks
Field Corn/ Popcorn (Preemergent Use)	Armyworm Cutworms	0.1 - 0.2	Apply in the time period from 5 days prior to planting up to emergence of the crop. Apply as a broadcast spray in a minimum of 20 gallons of finished spray/acre with ground equipment. AMBUSH may be mixed with the herbicide Ortho Paraquat® CL or GRAMOXONE® Paraquat in accordance with label limitations and precautions. Label dosage rates should not be exceeded.

**OVER**

Armyworm

equipment.

AMBUSH may also be injected into overhead sprinkler irrigation water provided (1) an anti-backflow check valve is present between the injection point and the water source, (2) the injection system has a check valve in the line to prevent irrigation water from entering the chemical supply tank, and (3) the irrigation injection system must have interlocking on-off switches.

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

**This labeling must be in the possession of the user  
at the time of pesticide application**

Ortho Paraquat® CL Herbicide is a registered trademark of Chevron Chemical Co  
GRAMOXONE® Paraquat Herbicide is a registered trademark of ICI Americas Inc

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC  
ANIMALS  
WARNING**

**HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. MAY CAUSE EYE IRRITATION.**

Avoid contact with eyes, skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly after handling.

**STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED** Do not induce vomiting. Call a physician. Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages via aspiration of the solvent. **IF IN EYES** Flush with plenty of water. Get medical attention if irritation persists. **IF ON SKIN** Wash with plenty of soap and water.

For 24-hour emergency assistance call 302/575-3000

**ENVIRONMENTAL HAZARDS:** This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply AMBUSH, or allow it to drift to crops or weeds on which bees are actively foraging. Additional information may be obtained from your Cooperative Extension Service. Apply this product only as specified on this label. In case of a significant spill, call CHEMTREC 800/424-9300.

**PHYSICAL OR CHEMICAL HAZARDS:** Do not use or store near food or open flame.

**STORAGE AND DISPOSAL**

Keep containers closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area and dispose of in accordance with State and local regulations. Do not reuse container. Triple rinse residue from empty container and add rinse to mixture to spray tank. Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures. Then after empty container for recycling or reconditioning, or purchase and dispose of in accordance with applicable Federal, State, and local procedures.

**RESTRICTED USE PESTICIDE.**

This label shall be used only by Certified Applicators, or persons under their direct supervision, and only for those uses authorized by the Certified Applicator's certification.



**Keep Out  
of Reach  
of Children  
WARNING**

See Side Panel For Additional  
Precautionary Statements and  
Statement of Practical  
Treatment.

EPA Est. No. 5005-AR-1  
EPA Reg. No. 12182-18

Net Contents  
**ONE U.S. GALLON**  
(3.79 LITERS)  
Batch No.

ICI Americas Inc.  
Pesticide Division  
Shawnee, Kansas 66201

U.S. Patent No. 4,824,162

**ACTIVE INGREDIENT:**

Permethrin-

(3-Phenoxyphenyl methyl (±)-cis, trans-3-(2,2-dichloroethoxy)-  
2,2-dimethylcyclopropanecarboxylate

25.0%

**INERT INGREDIENTS**

74.4%

**TOTAL** 100.0%

\*Gross Weight: Min. 35% (±) and max. 65% (±) **trans**  
AMBUSH contains 2 pounds active ingredient per gallon

It is a violation  
Apply AMBUSH  
Supplemental

Pound Active Ingredient	
0.05	
0.10	
0.15	
0.20	

**NOTICE**  
chemicals  
used in  
operations  
may  
result in  
EXPOSURE  
ABILITY OF  
DAMAGES  
SOLE LIAB  
THE REFUN

ACC  
JUL 2  
EPA Reg. No.

days prior to planting up to emergence of the crop. Apply as a broadcast spray in a minimum of 20 gallons of finished spray per acre with ground equipment. AMBUSH may be mixed with the herbicide Ortho Paraquat CL or GRAMOXONE Paraquat in accordance with label limitations and precautions. Label dosage rates should not be exceeded.

Foliar Use: For control of Cutworms, European Corn Borer and Armyworm, apply at a rate of 0.1 to 0.2 pounds active prior to ear formation. Apply a minimum of 1 gallon of finished spray per acre by air and 10 gallons per acre with ground equipment. AMBUSH may also be injected into overhead sprinkler irrigation water provided (1) an anti-backflow check valve is present between the injection point and the water source, (2) the injection system has a check valve in the line to prevent irrigation water from entering the chemical supply tank and (3) the irrigation injection system must have interlocking on-off switches.

11

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC  
ANIMALS  
WARNING**

**HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. MAY CAUSE EYE IRRITATION.**

Avoid contact with eyes, skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly after handling.

**STATEMENT OF PRACTICAL TREATMENT:** IF SWALLOWED: Do not induce vomiting. Call a physician. Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages and aspiration of the solvent. IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists. IF ON SKIN: Wash with plenty of soap and water.

For 24 hour emergency assistance call 302/575-3000

**ENVIRONMENTAL HAZARDS:** This pesticide is extremely toxic to fish. Use with care when applying in or near aquatic or to any body of water. Do not apply directly to water. Do not apply when water conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply AMBUSH or allow it to drift to crops or weeds on which bees are actively foraging. Additional information may be obtained from your Cooperative Extension Service. Apply this product only as specified on this label.

In case of a significant spill, call CHEMTREC 800/424-9300

**PHYSICAL OR CHEMICAL HAZARDS:** Do not use or store near food or open flame.

**STORAGE AND DISPOSAL**

Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area and dispose of in accordance with State and local regulations. Do not reuse container. Triple rinse residue from empty container and add rinse to mixture in spray tank. Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures. Then offer empty container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved State and local procedures.

**RESTRICTED USE PESTICIDE.**



**Keep Out  
of Reach  
of Children  
WARNING**

See Side Panel for Additional  
Precautionary Statements and  
Statement of Practical  
Treatment

EPA Est. No. 5905 AR 1  
EPA Reg. No. 10182-18

Net Contents  
**ONE U.S. GALLON**  
(3.79 LITERS)

Batch No.

ICI Americas Inc.  
Newark, NJ 07102

U.S. Patent No. 4,824,163

**ACTIVE INGREDIENT:**

Permethrin-

(3-phenoxyphenyl methyl (1:1:5:6:1:2:2:3:12:2-dichloroethoxy))  
2,2-dimethyl-3-(2-sulfonyl-1,3-dioxane-5-carboxylate)

**INERT INGREDIENTS**

**TOTAL**

25.8%  
74.4%  
100.0%

100% active ratio. Min. 35% active and max. 65% active.  
AMBUSH contains 2 pounds active ingredient per gallon.


**DIRECTIONS FOR**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply AMBUSH® as indicated on the Supplemental Pesticide Supplemental Pesticide

**RATE CONVERSION**

Pounds Active/ Acre	Fluid Ounces/ Acre	Pounds Active/ Acre
0.05	3.2	0
0.10	6.4	0
0.125	8.0	0
0.15	9.6	0
0.175	11.2	0
0.20	12.8	0

**NOTICE TO BUYER AND USER:** Seller warrants chemical description on the label and is reasonably to be used in accordance with directions under normal conditions for the use of this product contrary to label conditions or under conditions not reasonably foreseeable. Seller disclaims any liability for damages resulting from the use or handling of this product. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. SELLER LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE THE RETURN OF THE PURCHASE PRICE.





Apply first application when the adults appear and repeat at 5-7 day intervals or as needed throughout the season. AMBUSH may be diluted in a nonvolatile vegetable oil or in water in a minimum of one gallon of finished spray per acre using equipment calibrated to give adequate coverage.

DIS

AMBUSH® Insecticide  
EPA Reg. No. 10182-18

The following changes are proposed under the "Directions for Use":

Under Apples changed the rate to read 0.1 to 0.4 lb. a.i./A.\*

Under Cotton added Cutworm to the list of insects controlled.

Under Peaches changed the rate to read 0.1 to 0.4 lb. a.i./A.\*

Under Spinach added Vegetable Leafminer to the list of insects controlled and aerial application.

Under Sweet Corn added Corn Rootworm Adults to the list of insects controlled. Added the previously approved LANNAT AMBUSH tank mix for cotton to sweet corn.

ACCEPTED

JUL 13 1984

10182-18

\* ICI in a letter of March 15, 1984, has previously requested that the rate be expressed in terms of lb. a.i./A rather than lb. a.i./100 gal.

1128 19

COPY A

0.1 - 0.4 lb. a.i./A

COPY B

Cutworm

COPY C

0.1 - 0.4 lb. a.i./A

COPY D

Vegetable Leafminer  
or air.

COPY E

Corn Rootworm Adults

AMBUSH may be mixed with LANNATE® (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded.


EST. 1128 19

**RESTRICTED USE PESTICIDE.**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**SUPPLEMENTAL LABEL FOR USE ON  
BELL PEPPERS**

AMBUSH® Insecticide  
EPA Reg. No: 10182-18



EST 1

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product  
in a manner inconsistent with its labeling

Apply AMBUSH® Insecticide as shown in the chart below. Use sufficient water to obtain full coverage of foliage. Timing and frequency of applications should be based upon insect populations reaching economic thresholds. The higher rates may be required under conditions of heavy infestation.

**DIRECTIONS FOR APPLICATION**

Crop	Target Pest(s)	Rate (lbs. a.i./A)	Remarks
Bell Peppers	Vegetable leafminer	0.1-0.2	Apply by ground or air equipment using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 8 sprays per season. Do not apply within 3 days of harvest.
	Cabbage looper		
	Flea beetle		
	Pepper weevil		
	European corn borer	0.2	

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed


This labeling must be in the possession of the user  
at the time of pesticide application

**RESTRICTED USE PESTICIDE.**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**SUPPLEMENTAL LABEL FOR USE ON  
RANGE GRASS****AMBUSH® Insecticide****EPA Reg. No: 10182-18****DIRECTIONS FOR USE**

It is a violation of Federal law to use this product  
in a manner inconsistent with its labeling

**DIRECTIONS FOR APPLICATION**

Crop	Range grass
Target Pest	Range caterpillar
Dosage	0.01 lb. a.i./A
Remarks	Apply by ground or air using sufficient volume to obtain uniform coverage. Do not apply more than once per year. Cattle may be present during application. Do not harvest or feed hay to livestock.

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

This labeling must be in the possession of the user  
at the time of pesticide application

**RESTRICTED USE PESTICIDE.**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**SUPPLEMENTAL LABEL FOR USE ON  
EGGPLANT****AMBUSH® Insecticide****EPA Reg. No: 10182-18****DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product  
in a manner inconsistent with its labeling**

Apply AMBUSH® Insecticide as shown in the chart below. Use sufficient water to obtain full coverage of foliage. Timing and frequency of applications should be based upon insect populations reaching economic thresholds. The higher rates may be required under conditions of heavy infestation.

**DIRECTIONS FOR APPLICATION**

Crop	Target Pest(s)	Rate (lbs. a.i./A)	Remarks
Eggplant	Colorado potato beetle	0.2	Apply by ground or air equipment using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 16 applications. Do not apply within 3 days of harvest.
	Vegetable leafminer	0.1-0.2	
	Cabbage looper		
	Flea beetle		

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

**This labeling must be in the possession of the user  
at the time of pesticide application**

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long-legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

BEST AVAILABLE COPY

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

## WARNING

Area treated with AMBUSH 2E on (insert date). Do not enter without appropriate protective clothing until sprays have dried (if more restrictive, insert State Department of Agriculture reentry interval).

In case of accidental exposure to pesticide spray, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. Get medical attention if needed.

COPY D

0.2 - 0.4

COPY E

Apply by ground equipment only, using 25-400 gallons of spray per acre. Spray to wet all foliage. Do not apply after petal fall.

COPY F

AMBUSH may also be applied by air diluted with a vegetable oil such as cottonseed oil or soybean oil. Agricultural mineral oils (crop oils) are not recommended. Apply using a minimum of 1 quart of finished spray per acre.

COPY G

0.2 - 0.4



11 :Y

COPY H

Apply by ground equipment using 25-400 gallons of spray per acre. Spray to wet all foliage. Can be applied up to 7 days before harvest. Do not exceed 8 applications per acre per season. Do not graze livestock in treated orchards.

COPY I

0.2 - 0.4

COPY J

Cabbage Looper

COPY K

Western Bean Cutworm

BJK/mah  
030984mah09

BEST AVAILABLE COPY

DIRECTIONS FOR APPLICATION

Crop	Target Pest	Dosage lb ai/A	Comments
Chrysanthemums (Florida only)	Vegetable Leafminer	0.1 - 0.2	Apply by ground.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC  
ANIMALS  
WARNING**

**HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. MAY CAUSE EYE IRRITATION.**

Avoid contact with eyes. Skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly after handling.

**STATEMENT OF PRACTICAL TREATMENT IF SWALLOWED:** Do not induce vomiting. Call a physician. Vomiting should be supervised by a physician or the professional staff. In case of the pesticide poisoning, do not give anything to the patient. IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists. IF ON SKIN: Wash with plenty of soap and water.

For 24-hour emergency assistance call 302-575-3000.

**ENVIRONMENTAL HAZARDS:** This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply AMBUSH or allow it to drift to crops or weeds on which bees are actively foraging. Additional information may be obtained from your Cooperative Extension Service. Apply this product only as specified on this label.

In case of a significant spill, call CHEMTREC 800-424-9300.

**PHYSICAL OR CHEMICAL HAZARDS:** Do not use or store near heat or open flame.

**STORAGE AND DISPOSAL**

Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on hard or paved surface, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area and dispose of in accordance with State and local regulations. Do not reuse container. Triple rinse residue from empty container and add rinse to mixture in spray tank. Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures. Then after empty container for recycling or reconditioning or purchase and dispose of in a sanitary landfill or by other approved State and local procedures.

**RESTRICTED USE PESTICIDE.**



**Keep Out  
of Reach  
of Children  
WARNING**

See Side Panel For Additional  
Precautionary Statements and  
Statement of Practical  
Treatment

EPA Est. No. 5905 AR-1  
EPA Reg. No. 10182-10

Net Contents  
**ONE U.S. GALLON**  
(12.9 LITERS)  
Batch No.

ICI Americas, Inc.  
P.O. Box 10000  
Greenville, SC 29615

U.S. Patent No. 4,024,163

**ACTIVE INGREDIENT**

**Permethrin-**

(3-Phenoxyphenyl methyl (1:1 cis, trans)-3-(2,2-dichloroethoxy)-2,2-dimethylcyclopropanecarboxylate)

**INERT INGREDIENTS**

**TOTAL**

Cis/trans ratio: Min. 35% (1:1 cis) and max. 65% (1:1 trans)  
AMBUSH contains 2 pounds active ingredient per gallon.

25.6%

74.4%

100.0%

**DIRECT**

It is a violation of Federal Law to use this  
Apply AMBUSH<sup>®</sup> as indicated in the Supp.  
Supporter or Pamphlet.

**RATE CONVERSION**

Pounds Active Acre	Fluid Ounces Acre
0.05	3.2
0.10	6.4
0.125	8.0
0.15	9.6
0.175	11.2
0.20	12.8

**NOTICE TO BUYER AND USER:**  
Chemical description on the label and all  
used in accordance with directions and  
extend to the use of this product con-  
ditions or under conditions not re-  
assumes the risk of any such use.  
EXPRESSED OR IMPLIED, INCLUDING  
ABILITY, SELLER SHALL NOT BE LIABLE  
DAMAGES RESULTING FROM THE USE  
SOFT LIABILITY AND BUYER'S AND W  
THE RETURN OF THE PURCHASE PR

AC  
JUL 1 1984  
E...