

PM-13

Reg # 10182-18

1925

RESTRICTED USE PESTICIDE
Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

ACCEPTED
APR - 8 1996
I hereby certify that the above information is true and correct for the product described herein.
10182-18

AMBUSH® Insecticide.

CONDITIONS OF SALE
AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

2925

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.

Made in U.S.A.
ZENECA Ag Products
ZENECA Inc.
Wilmington, DE 19850-5458

TABLE OF CONTENTS

Page

STATEMENT OF PRACTICAL TREATMENT

PRECAUTIONARY STATEMENTS

 Hazards to Humans and Domestic Animals

 Personal Protective Equipment

 Environmental Hazards

 Physical and Chemical Hazards

DIRECTIONS FOR USE

 Agricultural Use Requirements

 General Instructions

 Reentry Statements

 Chemigation

DIRECTIONS FOR APPLICATION

 Alfalfa

 Almonds

 Apples

 Artichokes

 Asparagus

 Asparagus Ferns

 Avocados

 Cantaloupes

 Celery

 Cherries

 Christmas Trees

 Chrysanthemums

 Cole Crops

 Broccoli

 Brussels Sprouts

 Cabbage

 Cauliflower

 - Chinese Broccoli

 Chinese Cabbage

 Collards

 Conifers

 Gotton

 Cucurbit Vegetables

 Eggplant

 Field Corn/Popcorn/Field Corn Grown for Seed (Preemergent Use)

 Field Corn/Popcorn/Field Corn Grown for Seed (Foliar Use)

 Filberts

 Horseradish

 Leafy Vegetables (except Brassica)

 Lettuce

 Mushrooms

 Onions (Dry Bulb and Garlic)

 Ornamentals Nursery Stock

 Papaya (Florida Only)

 Peaches

Pears (Dormant)

Pears (Pre-bloom/Seasonal)

Peppers (Bell)

Pine Seed Orchard

Pistachios

Potatoes

Pumpkins

Range Grass

Roses (field-grown)

Roses (greenhouse)

Soybeans

Spinach

Sweet Corn

Sweet Corn (Florida Only - ~~corn grown for the fresh market~~)

Tomatoes (FL and PR only)

Turnips (FL, GA, IL, IN, OK, SEC, TX, WA only)

Walnuts

STORAGE AND DISPOSAL

RATE CONVERSION CHART

5925

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

SI usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

ACTIVE INGREDIENT:

Permethrin (3-Phenoxyphenyl)methyl(±)- <i>cis, trans</i> -3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate*	25.6%
INERT INGREDIENTS	74.4%
TOTAL	100.0%

**Cis/trans* ratio: Min. 35% (±) *cis* and max. 65% (±) *trans*. AMBUSH contains 2 pounds active ingredient per gallon and is an emulsifiable concentrate.

EPA Reg. No. 10182-18

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Call a physician. Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages via aspiration of the solvent.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. MAY CAUSE EYE IRRITATION. Avoid contact with eyes, skin or clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber ≥ 14 mils or Nitrile Rubber ≥ 14 mils or Viton ≥ 14 mils.
- Shoes plus socks.
- Protective eyewear.

For Mushroom House Use:

- For exposures in enclosed areas, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
-

7 of 25

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

) Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
 - Chemical-resistant gloves, such as Barrier Laminate or Butril Rubber ≥ 14 mils or Nitrile Rubber ≥ 14 mils or Viton ≥ 14 mils.
 - Shoes plus socks.
 - Protective eyewear.
-

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Metal Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Glass Containers: Triple rinse (or equivalent). Then dispose of in a sanitary landfill, or by other approved state and local procedures.

FOR BULK AND MINI-BULK CONTAINERS:

Container Disposal: Reseal container and offer for reconditioning or triple rinse (or equivalent), and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

Container Precautions: Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.

REFILL ONLY WITH AMBUSH. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than AMBUSH will result in contamination and may weaken container. After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

GENERAL INSTRUCTIONS

Apply AMBUSH as shown in "Directions for Application". Use sufficient water to obtain full coverage of foliage. Timing and frequency of applications should be based upon insect populations reaching economic thresholds. The higher rates may be required under conditions of heavy infestation. **Do not apply more than 2.0 lbs ai per acre per season.** Crops in this label may be planted immediately after last treatment. Do not plant other crops within 60 days after last application. Any other crop may be planted if it is not harvested or grazed.

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your Local or State Agricultural Authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

CHEMIGATION

SPRINKLER IRRIGATION APPLICATION: Apply AMBUSH Insecticide at rates and timing described elsewhere in this label. Apply with center pivot or continuous-move equipment distributing 1/2 acre-inch or less during treatment. If stationary (solid sets, handlines or wheel lines other than continuous-move) are used, AMBUSH insecticide should be injected into no more than the last 20-30 minutes of the set. Do not apply when winds are greater than 10-15 mph to avoid drifts or wind skips. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

USE PRECAUTIONS - SPRINKLER IRRIGATION APPLICATION

- A. Apply this product only through (sprinkler including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.

- G. The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Do not apply through chemigation systems connected to public water systems.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
COLE CROPS - Brassica Leafy Vegetables			
Broccoli	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	by air or ground. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.
Brussels Sprouts	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	by air or ground. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.
Cabbage	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.2 (3.2-12.8)	by air or ground. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5-10 days or as needed. Do not apply more than 1.0 lb. a.i. per acre per season. Do not apply within 1 day of harvest.
Cauliflower	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	by air or ground. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.
Chinese Broccoli ^c (gai lan, white flowering)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	by air or ground. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.
Chinese Cabbage (Tight-heading varieties only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.2 (3.2-12.8)	by air or ground. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage foliage. Apply every 5-10 days or as needed. Do not apply more than 1.0 lb. a.i. per acre per season. Do not apply within 1 day of harvest.
Collards (NC, SC, GA, AZ, OK, TX, AR, IL only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^a	0.05-0.1 (3.2-6.4)	Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Spray using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 8 applications. Do not apply within 1 day of harvest. Do not graze treated areas or feed crop refuse to livestock.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
CUCURBIT VEGETABLES			
Cucurbit Vegetables+	Leafminers and Squash Bugs	0.2 (12.8)	Apply by air or ground equipment. Apply a minimum of 4 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 1.6 pounds active ingredient per acre per season. AMBUSH may be applied up to harvest.
	Cabbage Looper Cucumber Beetle (adults) Cutworms Melon Worm Pickleworm Plant Bugs (including Lygus and Stink Bugs) Rindworms and Squash Vine Borer	0.1-0.2 (6.4-12.8)	
FRUITING VEGETABLES			
Eggplant	Colorado Potato Beetle	0.2 (12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use using sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 3.0 lbs. a.i. per acre per season. Do not apply within 3 days of harvest.
	Vegetable Leafminer Cabbage Looper Flea Beetle	0.1-0.2 (6.4-12.8)	
Peppers (Bell)	Vegetable Leafminer Cabbage Looper Flea Beetle Pepper Weevil	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use using sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 1.6 lbs. a.i. per acre per season. Do not apply within 3 days of harvest.
	European Corn Borer	0.2 (12.8)	
Tomatoes (Only for application to tomatoes grown in Florida and Puerto Rico for final marketing as fresh tomatoes.)	Tomato Fruitworm Cabbage Looper Tomato Pinworm Vegetable Leafminer Beet Armyworm Southern Armyworm Colorado Potato Beetle Granulate Cutworm Hornworm	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Do not apply more than 1.2 lbs. a.i./acre per crop season. Can be applied up to the day of harvest. Do not apply as a band. Do not use on tomato varieties in which the mature tomatoes will be less than 1 inch in diameter (such as cherry tomatoes).

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
LEAFY VEGETABLES			
Leafy Vegetables* (except Brassica)	Vegetable Leafminer Cabbage Looper Cutworm Complex Fall Armyworm Beet Armyworm Corn Earworm Green Cloverworm Leafhoppers European Corn Borer	0.1-0.2 (6.4-12.8)	by air or ground. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 3-5 days or as needed. Do not apply more than 2.0 lbs. a.i. per acre per season. Can be applied up to 1 day of harvest. Spinach only. Do not apply more than 1.0 lbs. ai per acre per season.
ROOT, TUBER AND BULB VEGETABLES			
Horseradish	Imported Crucifer Weevil (<i>Bariis lepidii</i>) control	0.1% a.i. solution ^d in a spring preplant dip and 0.2 lb ai/A (12.8) as a foliar spray.	Soak sets for 30 min. and air-dry before planting; make up to 3 foliar applications as needed to control weevil adults during oviposition. Apply by ground equipment in a minimum spray volume of 20 gallons per acre. Do not apply within 22 30 days of harvest.
Onions (Dry Bulb and Garlic)	Onion Thrips Armyworms Cutworms Leafminers Onion Maggots (adult) Stink bugs	0.15-0.3 (9.6-19.2) 0.1-0.3 (6.4-19.2)	Apply by air or ground equipment. Apply in a minimum of 20 gallons of water per acre with ground equipment or in a minimum of 6 gallons per acre by air as foliar spray. Begin applications when pests appear and repeat as necessary to maintain control. Use the higher label rates as onion thrips population increases and avoid rescue situations. Do not apply more than 2.4 2.0 lbs. active per acre per season. Do not apply within 1 day of harvest. Do not graze livestock in treated areas or cut treated crops for feed.
Potatoes	Colorado Potato Beetle Potato Leafhopper Aster Leafhopper Potato Flea Beetle Potato Aphid Potato Tuberworm Potato Psyllid Cabbage Looper	0.05-0.2 (3.2-12.8)	by air or ground. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply as needed. Do not apply more than 2.4 1.6 lbs. a.i. per acre per season. Can be applied up to 7 14 days prior to harvest.
Turnips (FL, GA, IL, IN, OK, SC, TX, WA only) (TX, OK, IL, IN only) (SC, GA, FL, WA only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid*	0.05-0.1 (3.2-6.4)	Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Spray using Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply within 1 day of harvest. Do not graze treated areas or feed crop refuse to livestock. Do not exceed 8 applications. Do not exceed 4 applications.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
CEREAL GRAINS			
Field Corn/Popcorn/ Field Corn Grown for Seed (Preemergent Use)**	Armyworm Cutworms	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. Apply as a broadcast spray in a minimum of 10 gallons of finished spray/acre with ground equipment or 1 gallon/acre by air.** By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons water per acre. Apply in the time period from 5 days prior to planting up to emergence of the crop.
Field Corn/Popcorn/ Field Corn Grown for Seed (Foliar Use)**	Cutworms European Corn Borer Armyworm Southwestern Corn Borer Flea Beetle Corn Earworm*** Corn Rootworm Beetle*** Tobacco Budworm*** Stalk Borer	0.1-0.2 (6.4-12.8)	Apply prior to the brown silk stage. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply a minimum of 1 gallon of finished spray/acre by air and 10 gallons/acre with ground equipment. Do not apply more than 0.6 lbs ai per acre per season. Allow a minimum of 5 days between treatments. Do not apply within 30 days of harvest of grain or fodder (stover). Forage may be harvested on the day of application. When treating for stalk borer, AMBUSH must be applied when or shortly before the stalk borer larvae are moving into the corn from surrounding weeds and grasses. Mowing or burndown herbicide is suggested to initiate movement.
Sweet Corn	Corn Earworm European Corn Borer Fall Armyworm Corn Rootworm Adults Cutworms Aster Leafhopper Flea Beetle ^c	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply every 5 days or as needed. Do not apply more than 1.2 lbs. a.i. per acre per season. Do not apply within 1 day of harvest of ears or forage or livestock grazing. AMBUSH may be mixed with LANNATE [®] (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded.
Sweet Corn (Florida only --corn grown --for the fresh --market)	Corn Earworm European Corn Borer Fall Armyworm Southern Armyworm Corn Rootworm Adults Cutworms Aster Leafhopper	0.1-0.25 (6.4-16.0)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply every 5 days or as needed. Do not apply more than 3.25-2.0 lbs a.i. per acre per crop season. Do not graze livestock in treated areas or harvest for forage. AMBUSH may be mixed with LANNATE [®] (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
TREE CROPS			
Almonds	Navel Orangeworm Peach Twig Borer	0.2-0.4 (12.8-25.6)	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Can be applied up to 7 days before harvest. Do not apply more than 2.0 lbs. a.i. per acre per season. Do not graze livestock in treated orchards. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.
Apples	Apple Aphid Redbanded Leafroller Oblique banded Leafroller Plum Curculio White Apple Leafhopper Spotted Tentiform Leafminer Tarnished Plant Bug	0.1-0.4 (6.4-25.6)	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Use 25-400 gallons of spray per acre. Spray to wet all foliage. Do not apply after petal fall. Do not apply more than 0.6 lbs a.i. per acre per season.
Avocados	Thrips Leafhoppers Mirid bugs Lepidopterous larvae	0.2 (12.8) [†] <u>fl.oz/100 gal.</u> 12.8	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Apply when insects appear and repeat at 7-10 day intervals as needed to provide control. Do not make more than 6 applications per season. Do not apply within 7 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.
Cherries	Lesser Peachtree Borer Rose Chafer Green Fruitworm Redbanded Leafroller Plum Curculio Tarnished Plant Bug	Rate lbs ai/100 gal. 0.025-0.05 (1.6-3.2) 0.1-0.2 (6.4-12.8)	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Do not exceed 400 gallons per acre. Apply when insects appear. In the East, do not exceed 6 applications per season, with no more than 4 applications after petal fall. In the West, do not apply more than 4 applications per season, with no more than 3 applications after petal fall. Do not apply within 3 days of harvest. Do not graze livestock in treated orchards. Do not feed forage to livestock. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.
Filberts	Filbertworm Oblique Banded Leafroller	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. Apply in 25-400 gallons of water per acre. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply when insects appear. Do not apply more than 1.6 lbs a.i. per acre per season. Apply in a minimum of 15 gallons of finished spray per acre by aircraft or 20 gallons of finished spray per acre with ground equipment. Do not apply within 14 days of harvest. Do not graze livestock in treated orchards. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS AII/A (OZ/A)	REMARKS
Papaya (Florida only)	Fruit Fly	0.1 (12.8) ¹ <u>oz./100 gal.</u> (6.4)	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Apply when insects appear and repeat at 7-10 day intervals as needed to provide control. Do not make more than 6 applications per season. Do not apply within 7 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.
Peaches	Peach Twig Borer Oriental Fruit Moth Green Fruitworm Lesser Peachtree Borer Plum Curculio Tarnished Plantbug Rose Chafer	0.1-0.4 0.5 (6.4-25.6 19.2)	Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 25-400 gallons of water per acre. Spray to wet all foliage. Can be applied up to 7 8 days before harvest. Do not apply more than 1.5 3.0 lbs. a.i. per acre per season. Do not graze livestock in treated orchards. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.
Pears (Dormant)	Pear Psylla	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. Apply in 3 to 20 gallons of finished spray per acre by aircraft and 25 to 400 gallons per acre by ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply at a rate of 0.8 to 1.6 pints (0.2 to 0.4 lbs. active) per acre. May be combined with 2 to 8 gallons of spray oil per acre. Apply during the dormant through delayed dormant growth periods only. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not graze orchard floor. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.
Pears (Pre-bloom/Seasonal)	Pear Psylla Codling Moth Green Fruitworm	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. Apply in 10 to 20 gallons of finished spray per acre by aircraft and 25 to 400 gallons per acre by ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply two pre-bloom sprays followed by three seasonal sprays. Pre-bloom sprays can be applied from dormant through bud burst stages. Seasonal sprays can be applied from petal fall up to 14 days of harvest. Do not apply more than 2.0 0.8 lbs. a.i. per acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
Pistachios	Peach Twig Borer Naval Orangeworm Ch.nch Bug	0.2-0.4 (12.8-25.6)	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply at a rate of 0.8 to 1.6 pints (0.2-0.4 lb active) per acre. Do not apply more than 0.8 lbs a.i. per acre per season. Do not apply after 10 percent hull split. Nuts may be harvested on the day of application. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. Do not graze livestock in treated orchards.
Walnuts	Navel Orangeworm Codling Moth Walnut Husk Fly	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. Apply in a minimum of 20 gallons of finished spray per acre by aircraft or 100 gallons of finished sprays per acre with ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply when insects appear. Do not apply more than 1.6 lbs. a.i. per acre per season. Do not apply within 1 day of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. Do not graze livestock in treated orchards.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
OTHER CROPS			
Alfalfa, Alfalfa grown for seed	Alfalfa Caterpillar Armyworms Cutworms Loopers Webworms Potato Leafhopper Blue Alfalfa Aphid ^f Green Peach Aphid Pea Aphid ^f Green Cloverworm Spotted Alfalfa Aphid Velvetbean Caterpillar	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. Use higher recommended dosage for increased pest pressure or for increased residual pest control. Apply in a minimum of 4 gallons of finished spray per acre by aircraft or 10 gallons of finished spray per acre with ground equipment. Use higher gallonage, 5-10 gallons by air, or 20 by ground, finished spray per acre when foliage is dense and/or when pest populations are high. Do not apply more than 0.2 pound active per cutting. When rates 0.1 pound active per acre or less are used, application may be made on day of harvest. When rates greater than 0.1 pound active per acre are used, do not apply within 14 days of harvest. Avoid application when bees are actively foraging by applying during the early morning or during the evening hours.
	Alfalfa Weevil ^f Egyptian Alfalfa Weevil ^f Cucumber Beetle Meadow Spittlebug Plant Bugs including Lygus spp. ^f Stink Bugs	0.1-0.2 (6.4-12.8)	
Artichokes	Artichoke plume moth	0.1-0.3 (6.4-19.2)	Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Use using sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 1.5 lbs. a.i. per acre per season. Buds may be harvested on the day of application.
Asparagus	Cutworm Complex Asparagus Beetle	0.05-0.1 (3.2-6.4)	Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Use using sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 0.4 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
Cotton	Bollworm Tobacco Budworm Pink Bollworm Lygus Bugs Cabbage Looper Cotton Leafperforator Boll Weevil Cotton Fleahopper Saltmarsh Caterpillar Whitefly [®] Cotton Aphid [®] Cutworms	0.1-0.2 (6.4-12.8)	<p>Apply as required by scouting reports, usually at intervals of 5-7 days. Apply by air or ground. Use sufficient water to obtain full coverage of foliage. AMBUSH may also be applied by air diluted with a vegetable oil such as cottonseed oil or soybean oil. Agricultural oils (crop oils) are not recommended. Apply using a minimum of 1 quart of finished spray per acre. For control of boll weevils spray on a 3 to 5 day schedule or as needed. Do not apply within 14 days of harvest. The higher rates of AMBUSH (0.15 to 0.2 lbs. a.i. per acre) may be required under conditions of heavy lepidopterous worm infestation. AMBUSH aids in the reduction of lepidopterous egg hatch by contact action. Do not graze livestock in treated areas. Do not apply more than 2.0 lbs. a.i. per acre per season.</p> <p>Do not make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include AMBUSH[®] insecticide, Ammo[®] insecticide, Asana[®] XL insecticide, Baythroid[®] emulsifiable pyrethroid insecticide, Capture[®] insecticide/miticide, Danitol[®] 2.4 EG Spray insecticide/miticide, Fury[®] insecticide, KARATE[®] insecticide, Mustang[®] insecticide, Pounce[®] insecticide and Scout X-TRA[®] insecticide.</p> <p>All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.</p>

DIRECTIONS FOR APPLICATION

CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
Cotton (cont.)			<p>SPRAY DRIFT PRECAUTIONS: OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.</p> <p>Do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams; marshes, or natural ponds; estuaries and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultralow volume (ULV) application is made.</p> <p>For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.</p> <p>Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.</p> <p>Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.</p> <p>Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.</p> <p>Risk of exposure to aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.</p> <p>Do not cultivate within 10 feet of the aquatic area so as to allow growth of a vegetative filter strip.</p> <p>Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.</p> <p>Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.</p>

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
Mushrooms (Mushroom houses and adjacent premise areas)	Mushroom Flies (Sciariid and Phorid Adults)	Apply as a fogging or aerosol treatment at the rate of 3.2 to 4.0 ounces (0.05-0.0625 lb ai) per 30 ounces of water or suitable diluent. Use 1 quart of solution per standard double house (35,000 cu ft.; 8,000 sq. ft.)	Preparation of the building prior to fogging: (1) Close all doors, windows, and ventilators, (2) Lock or barricade all entrances, turn off pilot lights, post warning signs, and take precautions to prevent persons and animals from entering the area. Use prior to filling the house, during cool-down, during spawning, up to pinning and between breaks. Do not use when mushrooms are present. Treat once daily or as needed when flies appear. Do not make more than 20 applications prior to pinning of first break; apply no more than two applications between each break. Do not apply more than 30 applications total per crop of 5 breaks. Do not apply within 3 days of harvest. Length of exposure time should be limited to 1 hour, then ventilate the house. Use fans to ventilate in houses that do not have forced air circulation. Wear a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).
Range Grass	Range Caterpillar	0.01 (0.64)	Apply by ground or air. Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient volume to obtain uniform coverage. Apply as needed. Do not apply more than once per year. Cattle may be present during application. Do not harvest or feed hay to livestock.
Soybeans	Green Cloverworm Mexican Bean Beetle Bean Leaf Beetle Cabbage Looper Potato Leafhopper Velvetbean Caterpillar Saltmarsh Caterpillar Corn Earworm Beet Armyworm Japanese Beetle*** Soybean Looper*	0.05-0.1 (3.2-6.4) 0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Do not apply more than 0.4 lbs. a.i. per acre per season. Can be applied up to 60 days before harvest. Do not graze treated areas or harvest for forage or hay.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
ORNAMENTALS			
Asparagus Ferns	Japanese Beetle Asparagus Beetle Lygus Bugs	0.1 (6.4)	Apply to the fern stage of the asparagus plant after spear harvest when larval and adult stage of the asparagus beetle, lygus bugs and the adult stage of the Japanese beetle are present.
Christmas Trees ^c (container and field grown)	Nantucket Pine Tip Moth	0.1-0.2 (6.4-12.8)	Apply first application when adults appear and repeat at 5-7 day intervals or as needed throughout the season. Use sufficient water to obtain full coverage of foliage.
Chrysanthemums (Florida only)	Vegetable Leafminer	0.1-0.2 (6.4-12.8)	Apply by ground.
Conifers (container and field grown)	Nantucket Pine Tip Moth	0.1-0.2 (6.4-12.8)	Apply first application when the adults appear and repeat at 5-7 day intervals or as needed throughout the season. AMBUSH may be diluted in a nonvolatile vegetable oil or in water in a minimum of one gallon of finished spray per acre using equipment calibrated to give adequate coverage.
Ornamentals Nursery Stock ^c (field-grown)	Beet Armyworm Cabbage Looper Citrus Thrip Heliothis Spp. Leafhoppers Leafminers Whiteflies Cutworms	0.1-0.2 (lbs ai/100 gal)	Marginal leaf burn may occur on Salvia, Dieffenbachia and Pteris Fern. Application to blooming plants may cause browning of petals. Not all species and varieties have been tested. Before treating large numbers of plants of a particular variety, treat a few plants and observe prior to full scale application. Apply to nonbearing plants of fruiting species.
Pine Seed Orchard	Coneworms Seed bugs	13 fl oz/ 100 gals water (0.025% dilution by wt) for high volume sprayers; 68 fl oz/ 100 gals water (0.125% dilution by wt) for low volume sprayers; 48 fl oz/A application in not less than 10 gals of water for aerial applications	Apply first application within 1 week of female flower closure or peak pollen flight for webbing coneworm control. For other coneworms and seed bugs, apply first application within 30 days following female flower closure. Repeat applications at intervals of 4 weeks, but do not apply more than 6 applications. Apply approximately 5-10 gals of the 0.025% dilution per tree with high volume sprayers. With low volume sprayers, apply 100 gallons of the 0.125% dilution per acre. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. Do not graze or harvest cover crop. Avoid contact with open water
Roses ^c (field grown)	Heliothis Spp	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. Apply in a minimum of 12 gallons/acre by air in and 10.5 gallons of finished spray per acre with ground equipment

DIRECTIONS FOR APPLICATION			
CRCP	TARGET PESTS	RATE LBS A/A (OZ/A)	REMARKS
Roses ^c (greenhouse)	Beet Armyworm Cabbage Looper Omnivorous Leafroller	8 fl oz/100 gals water	AMBUSH is not phytotoxic to the following varieties of greenhouse roses: Ballena, Bettina, Cara Mia, Coquette, Excitement, Forever Yours, G. Wave, Jack Frost, Jr. Bridesmaid, Matador, Paul's Pink, Samantha, Seventeen, Town Crier, Tropicana and Visa. Other varieties may vary in their sensitivity to AMBUSH, and a small number of plants should be treated under local conditions to determine plant safety prior to commercial use.

- * ~~LEAF VEGETABLES (Cress, Brassica)~~ Amaranth (leafy amaranth, Chinese spinach, tampala); arrugla (Roquette); celery, celtuce; chervil; corn salad; chrysanthemum, edible-leaved; chrysanthemum, garland; cress, garden; cress upland (yellow rocket, winter cress); dandelion; dock (sorrel); endive (escarole); fennel, Florence; lettuce (head and leaf); orach; parsley; purslane, garden; purslane, winter; rhubarb; spinach; spinach, fine; spinach, new Zealand; Swiss chard.
- + ~~CUCURBIT VEGETABLES~~ Balsam pear (Bitter Melon); Chinese Waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourds, edible; melons, including hybrids such as cantaloupe, casaba, crenshaw, honeydew melons, honey balls, mango melon, muskmelon, Persian melon; pumpkin; squash, summer and winter; watermelon, including hybrids.
- ** AMBUSH may be applied in tank-mixtures with the following herbicides in accordance with label limitations and precautions. Label dosage rates should not be exceeded. GRAMOXONE[®] SUPER; GRAMOXONE[®] EXTRA, BICEP[®], LASSO[®]/Atrazine Premix; LASSO[®] 4E & ME; DUAL[®] 8E; Atrazine 4L & 80W; BLADEx[®] 80W & 4L; PRINCEP[®]; PRINCEP[®] CALIBER 90[®]; 2,4-D; ROUNDUP[®]; BRONCO[®]; BANVEL[®]; EXTRAZINE[®]
- *** Not for control of this insect in California.
- Aids in suppression.
- Only suppression can be expected when utilized on a 5-7 day schedule.
- Currently not registered for use in California.
- 0.1% ai solution = 3 1/3 pt formulation per 100 gallons of water.
- Maximum use rate in California is 0.1 lbs. a.i./Acre.
- Under heavy insect pressure use 0.2 lbs a.i./Acre. A second application of alternate chemistry may be necessary to reduce populations below economic thresholds.
- ~~Do not use more than 12.8 oz. AMBUSH/A. The final spray should be no more concentrated than 12.8 oz./100 gallons for avocados and 6.4 oz./100 gallons for papayas. Since tree size and spacing will affect the amount of spray needed for dilute sprays, the applicator must know the number of gallons of water/A needed to spray the trees to the point of drip. If you do not know this gallonage, you should conduct a test to determine it. If you do not know how to conduct such a test with your equipment, you should ask for assistance from your equipment dealer. For dilute sprays (i.e., sprayed to the point of runoff) 12.8 oz. AMBUSH should be dilute with the gallonage determined by this test.~~

RATE CONVERSION CHART			
POUNDS ACTIVE/ACRE	FLUID OUNCES/ACRE	PINTS/ACRE	ACRES TREATED/ GALLON
0.05	3.2	0.2	40.0
0.10	6.4	0.4	20.0
0.125	8.0	0.5	16.0
0.15	9.6	0.6	13.3
0.175	11.2	0.7	11.4
0.20	12.8	0.8	10.0

) AMBUSH®, GRAMOXONE®, and KARATE® are trademarks of ZENECA Group Companies.
 Ammo®, Capture®, Fury™, Mustang® and Pounce® are trademarks of FMC Corporation.
 Baythroid® is a trademark of Mobay Corporation.
 Bronco®, Lasso®, and Roundup® are trademarks of Monsanto Company.
 Bicep®, Caliber®, Dual®, and Princep® are trademarks of Ciba-Geigy Group.
 Asana®, Bladex®, Extrazine® and Lannate® are trademarks of E.I. duPont de Nemours & Co.
 Banvel® is a trademark of Sandoz Crop Protection Corp.
 Danitol® is a trademark of Sumitomo Chemical Company, LTD.
 Scout X-TRA™ is a trademark of Hoechst-Roussel Agri-Vet Company.