PM D3 /	0.182-)8NOTIFICA			Approvel expires 05-31-98
Enviro	United States Onmental Protection Age Washington, DC 20460 Handeman To nonostes Late to a		Registration Amendment Other <u>A storro</u> t	OPP Identifier Number 248706 138 140 1471 1
ىرىيىلىم بەركىيىكىنى بىر بەركىك ئايامانى بەركىكى بەركىكىكى بەركىكىكىكىكىكىكىكىكىكىكىكىكىكىكىكىكىكىكى	Application for	Pesticide - Section	La reserve de la conserve	i na rižno set mienieni
1. Company/Product Number	A antoster isterprovid . 4 (f	2. EPA Product Manager George LaRocca	temes -	
	addentifier work is an anderlight a	9 - 13 - 2 m Ou S m	deuryoser "wi lisi d	None Restricted
1800 Concord Pike PO Box 15458 Wilmington, DE 19850-5458	Tenten Hyliky o d Bo lgeska for y	(b)(i), my product is simi	ar or identical in con 9005 Statestant (TT) 1005 Statestant 1005 Statestant	nposition and labeling
Check if this is a new	/ address	Product Name		
Amendment - Explain below./	n war not envised and an inter Sec Beaulie - Beau Brand an inter Sec 2 - 1 - Envised Presentation - Sec Agency letter dated <u>Strates of opport</u>	Final printed labels Agency letter date	in response to stilles d tionrest rest, and	 และ (การ์ตออง เขาสารสุดที่สะวิวิ มารายสะ 2003 เขาสารสารสารสารสาร อาการ (ค.ศ. 1012) (การรายสารสาร การรายสารสารสารสารสารสารสารสารสารสารสารสารสารส
Explanation: Use additional page(s Final printed labels reflec	s) if necessary. (For section I and S ting EPA approval use direct	Beaton II.)		
	Sec	tion - III	NOITA	
1. Material This Product Will Be Pack			tis was america to as a state of	ان از ها مخطقها کار ۲۹۹۹ مار ملایات مهناز ها مخطقها کار ۲۹۹۹ مار ملایات
Yes* No * Certification must	Yes No No. per If "Ye ckaging wgt. container Packa	No Soluble Packaging Yes No S [*] No. per Container	2. Type of Container Metal Plastic Glass Paper Other (S	
3. Location of Net Contents Informati			ation of Label Directio	
Label and and the Container	es e lo phái i i anaigst am bhanna i Is '		On Labeling accom	panying product
6. Manner in Which Cabel is Affixed t	o Product Paper glued Stenciled	Other	931. 491. 1813 (1898 187 	ers for er stad for any or estations
miseur goist" ;"	o mainecel : laimealined and LSec	tion - IV and to zeeq , ene	a to notibbe shi" 28 i	ราย มีระบบที่เพื่อเรื่อง เป็น
1: Contact Point (Complete items di	-	vidual to be contacted, if nece		application.)
Deborah Touzell	Regulato	ry Information Coordina	tor #302-880	No. (Include Area Code)
i acknowledge that any knowir both under applicable law.	iave made on this torm, and all artic ngly false or misleading statement m ng tabe or misleading statement m	number our tetel statisticter: tines in which product will be upper, out, ore objective training in, to enit, yd alderiging, days an of the use directions for yo statistic product is	rate and comparisons of contract of contract of the second comparison of the second contract of the second contrac	6. Data Application Received for Lossian J. S. Microsoft of Market Strength Microsoft of Market Strength Strength of Strength Strengt
2. Signatur		ory,Information_Coordi		A State Son VI MOITOSE
4. Typed Name Deborah Touzell	5. Date Septent	per 9, 1997	•	1-5. Soil-oxplanator. 3.5. EPA Use Only.
PA Form 8570-1 (Rev. 8-94) Previou	s editions are obsolete.	White - EP	A File Copy (original)	Yellow - Applicant Copy

À

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

ZENECA Ag Products



ACTIVE INGREDIENT: Permethrin

[3-Phenoxyphenyi]methyl(±)-cis, trans-3-{2,2-dichloroethenyi]-2,2dimethylcyclopropanecarboxylate* 25.6% INERT INGREDIENTS 74,4%

EPA Reg. No. 10182-18 EPA Est. No. 10182-AR-1



KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE FOR STATEMENT OF PRACTICAL TREATMENT PRECAUTIONARY STATEMENTS AND COMPLETE DIRECTIONS FOR USE

Contents: 1 gal (3.79 L)

0400XXXX/069608 Rev. 9/97

ג. נ נ

د د د ر د د

, , , , , , , , , , , , ,

NOTIFICATION

≠ 23 1997 Jotts

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ZENECA or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ZENECA and Seller harmless for any claims relating to such factors.

ZENECA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ZENECA, and Buyer and User assume the risk of any such use. ZENECA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall ZENECA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ZENECA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ZENECA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ZENECA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ZENECA.




TABLE OF CONTENTS

يردر المردوم م in the the state of the s . 1997 Hors \$76 5. A. J.

<u> NOE</u>

.....

. . .

.

		Pag
STATEMENT OF PRACTICAL TREATMENT		
PRECAUTIONARY STATEMENTS.	et genne en	
Hazards to Humans and Domestic Animals		
Personal Protective Equipment		
Environmental Hazards	-	
Physical and Chemical Hazards	· · ·	5
DIRECTIONS FOR USE		
Agricultural Use Requirements		
Storage and Disposal		
General Instructions		
Chemigation		
Use Precautions - Sprinkler Irrigation Application		
DIRECTIONS FOR APPLICATION		
COLE CROPS - Brassica Leafy Vegetables		
Broccoli		8
Brussels Sprouts	•	
Cabbace		
Cauliflower		
Chinese Broccoli		
Chinese Cabbaoe.		
Collards		-
CUCURBIT VEGETABLES	*******************************	
Cucurbit Vegetables		0
FRUITING VEGETABLES	*************************	
Eggplant		0
Peppers (Bell)		
Tomatoes		10
LEAFY VEGETABLES		
Leafy Vegetables (Except Brassica)	•••••••••••••••••	
ROOT, TUBER AND BULB VEGETABLES		
Horseradish		
Onions		
Potatoes		
Turnips		

<u>د د د د د</u> ננ ג ג גוזני ***** د ددد د د

>

40fus

CEREAL GRAINS	
Field Corn/Popcom/Field Com Grown for Seed (Preemergent Use)	12
Field Corn/Popcorn/Field Com Grown for Seed (Foliar Use)	12
Sweet Com	12
Sweet Com (Horida Only)	12
TREE CROPS	
Almonds	13
Apples	13
Avocados	13
Cherries	13
Filberts	14
Papaya (Florida Oniy)	14
Peaches	
Pears (Dormant)	14
Pears (Pre-bloom)	
Pistachios	
Walnuts	15
OTHER CROPS	
Alfalfa, Alfalfa Grown for Seed	
Artichokes	
Asparagus	
Mushrooms	
Range Grass	17
Soybeans	18
ORNAMENTALS	
Asparagus Ferns	
Christmas Trees	
Chrysanthemums (Florida Only)	
Conifers	
Ornamentals Nursery Stock	
Pine Seed Orchard	
Roses (Field Grown)	
Roses (Greenhouse)	
RATE CONVERSION CHART	21

•

,

۰.

50f35

• • • • • • • • • • • •

STATEMENT OF PRACTICAL TREATMENT

60425

IF SWALLOWED: Do not induce vomiting. Call a physician. Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages via aspiration of the solvent.

IF IN EVES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of scap and water.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633). FOR CHEMICAL EMERGENCY: Spill, look, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. MAY CAUSE EYE IRRITATION. Avoid contact with eyes, skin or clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.
 Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber ≥14 mils or Nitrile Rubber ≥ 14 mils or Viton ≥ 14 mils.

Shoes plus socks.

Protective evewear.

For Mushroom House Use:

 For exposures in enclosed areas, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NI0SH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NI0SH approval number prefix TC-14G).

number prenx 10-146J. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170-240kd)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

. WE

- ers should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. This product is highly toxic to bese exposed to direct treatment or residues on blooming crops or weeds. Bo not apply this product is a drift to drift to drift to drift the drift of t

or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

5

. .

DIRECTIONS FOR USE

idd S

,

ر مالين در الماني

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

AGRICULTURAL USE REDUREMENTS Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with enything that has been treated, such as plants, soil, or water, is:

Coveralits.

Chemical-resistant gloves, such as Barrier Laminate or Butril Rubber ≥ 14 mils or Nitrile Rubber ≥ 14 mils or Viton ≥ 14 mils. Shoes plus socks. Protective eyewear

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a

violation of Federal law. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by inclineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INSTRUCTIONS

Apply AMBUSH as shown in "Directions for Application". Use sufficient water to obtain full coverage of foliage. Timing and frequency of applications should be based upon insect populations reaching economic thresholds. The higher rates may be required under conditions of heavy infestation. Do not apply more than 2.0 lbs ai per acre per season. Crops in this label may be planted immediately after last treatment. Do not plant other crops within 60 days after last application. Any other crop may be planted if it is not harvested or grazed.

80H25

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your Local or State Agricultural Authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

CHEMIGATION

SPRINKLER IRRIGATION APPLICATION: Apply AMBUSH Insecticide at rates and timing described elsewhere in this label. Apply with center pivot or continuous-move equipment distributing 1/2 acre-inch or less during treatment. If stationary (solid sets, handlines or wheel lines other than continuous-move) are used, AMBUSH insecticide should be injected into no more than the last 20-30 minutes of the set. Do not apply when winds are greater than 10-15 mph to avoid drifts or wind skips. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period. USE PRECAUTIONS - SPRINKLER IRRIGATION APPLICATION

- A Apply this product only through (sprinkler including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check value, vacuum relief value, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- G. The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The inigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L Do not apply when wind speed favors drift beyond the area intended for treatment. M. Do not apply through chemigation systems connected to public water systems.



•••••

	DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS AI/A OZ/A}	REMARKS	
Broccoli	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid*	0.05-0.1 (3.2-6.4)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre. per season. Do not apply within 1 day of harvest.	
Brussels Sprouts	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid®	0.05-0.1 (3.2-6.4)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.	
Cabbage	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid"	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5-10 days or as needed. Do not apply more than 1.0 lb. ai. per acre per season. Do not apply within 1 day of harvest.	
Cauliflower	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid*	0.05-0.1 (3.2-6.4)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.	
Chinese Broccoli ^b (gai lon, white flowering)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid*	0.05-0.1 (3.2-6.4)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5 days or as needed. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not apply within 1 day of harvest.	

90f25

درز د د د د د د د

	DIREC	TIONS FOR API	PLICATION	
CROP	TARGET PESTS	RATE LBS AL/A OZ/A)	REMARKS fy Vegetables	
Chinese Cabbage (Tight-heading varieties only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ^e	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 5-10 days or as needed. Do not apply more than 1.0 lb. a.i. per acre per season. Do not apply within 1 day of harvest.	
Collards (NC, SC, GA, AZ, OK, TX, AR, IL only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ⁴	0.05-0.1 (3.2-6.4)	Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Spray using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 8 applications. Do not apply within 1 day of harvest.	
·	and the states and the second	CUCURBIT VEGETA	BLES	
Cucurbit Vegetables+	Leafminers and Squash Bugs Cabbage Looper Cucumber Beetle (adults) Cutworms Melon Worm Pickleworm Plant Bugs (including Lygus and Stink Bugs) Rindworms and Squash Vine Borer	02 (12.8) 0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. Apply a minimum of 4 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 1.6 pounds active ingredient per acre per season. AMBUSH may be applied up to harvest.	
· -	FRUITING VEGETABLES			
Eggplant	Colorado Potato Beetle Vegetable Leafminer Cabbage Looper Fiea Beetle	0.2 (12.8) 0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 20 lbs. a.i. per acre per season. Do not apply within 3 days of harvest.	

.

Dof25

1

• •

• ,

.

.

•.

9

,

110705

	DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS AI/A OZ/A)	REMARKS	
فللمساء للتعلقات المتخد العايدة		FRUITING VEGETA	BLES CONTRACTOR DE LA CONT	
Peppers (Bell)	Vegetable Leafminer Cabbage Looper Flea Beetle Pepper Weevil European Corn Borer	0.1-0.2 (6.4-12.8) 0.2 (12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 1.6 lbs, a.i. per acre per season. Do not apply within 3 days of harvest.	
Tomatoes	Tomato Fruitworm Cabbage Looper Tomato Pinworm Vegetable Leafminer Beet Armyworm Southern Armyworm Colorado Potato Beetle Granulate Cutworm Hornworm	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Do not apply more than 12 lbs. ai/acre per crop season. Can be applied up to the day of harvest. Do not apply as a band. Do not use on tomato varieties in which the mature tomatos will be less than 1 inch in diameter (such as cherry tomatoes).	
	a la contrata de la c	LEAFY VEGETAB	LES	
Leaty Vegetables* (except Brassica)	Vegetable Leafminer Cabbage Looper Cutworm Complex Fall Armyworm Beet Armyworm Com Earworm Green Cloverworm Leafhoppers European Com Borer	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply every 3-5 days or as needed. Do not apply more than 2.0 lbs. a.i. per acre per season. Can be applied up to 1 day of harvest. Spinach only - Do not apply more than 1.0 lb a.i. per acre per season.	
	And a standard a ROOT, T	UBER AND BULB	VEGETABLES	
Horseradish	Imported Crucifer Weevil (Baris Iepidii) control	0.1% a.i. solution ^e in a spring preplant dip and 0.2 lb ai/A (12.8) as a foliar spray.	Soak sets for 30 min. and air-dry before planting; make up to 3 foliar applications as needed to control weevil adults during oviposition. Apply by ground equipment in a minimum spray volume of 20 gallons per acre. Do not apply within 30 days of harvest.	

봁

,,

Dafts

در (زدور دود ود

))))))))))

. :

12-3

4 : 12

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS AVA OZ/A)	REMARKS
and the state of the	NOOT, T	UBER AND BULB	VEGETABLES
Onions (Dry Bulb and Garlic)	Onion Thrips Armyworms Cutworms	0.15-0.3 (9.6-19.2) 0.1-0.3	Apply by air or ground equipment. Apply in a minimum of 20 gallons of water per acre with ground equipment or in a minimum of 6 gallons per acre by air as foliar spray. Begin applications when pests appear and repeat as necessary to maintain control. Use the
	Leafminers Onion Maggots (adult) Stink bugs	(6.4-19.2)	higher label rates as onion thrips population increases and avoid rescue situations. Do not apply more than 2.0 lbs. active per acre per season. Do not apply within 1 day of harvest.
Potatoes	Colorado Potato Beetle Potato Leafhopper Aster Leafhopper Potato Flea Beetle Potato Aphid Potato Aphid Potato Poyllid Cabbage Looper	0.05-0.2 (3.2-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply as needed. Do not apply more than 1.6 lbs. a.i. per acre per season. Can be applied up to 14 days prior to harvest.
Turnips (FL, GA, IL, IN, OK, SC, TX, WA only) (TX, OK, IL, IN only)	Cabbage Looper Imported Cabbageworm Diamondback Moth Cabbage Aphid ⁴	0.05-0.1 (3.2-6.4)	Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply within 1 day of harvest. Do not graze treated areas or feed crop refuse to livestock. Do not exceed 8 applications.
(SC, GA, FL, WA only)			Do not exceed 4 applications.

۰,

2.

٦,

쁿

11

5 a 35 a ...

	DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS AI/A OZ/A}	REMARKS	
فكالمجمر بالمعاد والمعاد والمعادي				
Field Corn/Popcom/ Field Corn Grown for Seed (Preemergent Use)**	Armyworm Cutworms	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in minimum of 2 gallons of water per acre. By groun apply in a minimum of 10 gallons water per acre. Apply in the time period from 5 days prior to plantin up to emergence of the crop.	
Field Corn/Popcorn/ Field Corn Grown for Seed (Foliar Use)**	Cutworms European Com Borer Armyworm Southwestern Corn Borer Flea Beetle Com Earworm*** Com Rootworm Beetle*** Tobacco Budworm*** Stalk Borer	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in minimum of 2 gallons of water per acre. By ground apply in a minimum of 10 gallons of water per acre. "Do not apply more than 0.6 ibs ai per acre per season Allow a minimum of 6 days between treatments. D not apply within 30 days of harvest of grain or fodde (stover). Forage may be harvested on the day of application." When treating for stalk borer, AMBUSH must b applied when or shortly before the stalk borer larva are moving into the corn from surrounding weeds an grasses. Moving or burndown herbicide is suggeste to initiate movement.	
Sweet Corn	Com Earworm European Com Borer Fail Armyworm Com Rootworm Adults Cutworms Aster Leafhopper Flea Beetle ⁶	0.1-0.2 (6.4-12.8)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Apply every 5 days or as needed. Do not apply more than 1.2 lbs. a.i. per acre per season. Do not apply within 1 day of harvest of ears or forage or livestock grazing. AMBUSH may be mixed with LANNATE® (methomyl) in accordance with label limitations and precautions. Label dosage rates should not be exceeded.	
Sweet Corn (Florida only)	Com Earworm European Com Borer Fall Annyworm Southern Armyworm Com Rootworm Adults Cutworms Aster Leafhopper	0.1-0.25 (6.4-16.0)	Apply by air or ground equipment. By air, apply in minimum of 2 gallons of water per acre. By groun apply in a minimum of 10 gallons of water per acr Apply every 5 days or as needed. Do not apply moo than 2.0 lbs. a.i. per acre per crop season. AMBUS may be mixed with LANNATE® (methomyl) in accordance with label limitations and precaution: Label dosage rates should not be exceeded.	

•

•

13qf25

.

1

÷

.

.

• • • •

•-

12

.

,

DIRECTIONS FOR APPLICATION RATE LBS AVA TARGET CROP PESTS 0Z/A) REMARKS TREE CROPS Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Can be applied up to 7 days before harvest. Do not apply more than 2.0 lbs. a.i. per Navel Orangeworm 0.2-0.4 (12.8-25.6) Almonds Peach Twig Borer acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Do not apply after petal fall. Do not apply more than 0.6 lbs a.i, per acre per season. 0.1-0.4 (6.4-25.6) Apples Apple Aphid **Redbanded Leafroller** Oblique banded Leafroller Plum Curculio White Apple Leafhopper Spotted Tentiform Leafminer Tarnished Plant Bug Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Apply when insects appear and repeat at 7-10 day intervals as needed to provide control. Avocados Thrips 0.2 (12.8) Leathoppers Mirid bugs Lepidopterous larvae Do not make more than 6 applications per season. Do not apply within 7 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. 0.1-0.2 (6.4-12.8) Lesser Peachtree Borer Cherries Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Apply when insects appear. In the East, do not exceed 6 applications per **Rose Chafer** Green Fruitworm Redbanded Leafroller season, with no more than 4 applications after petal fall, in the West, do not apply more than 4 applications per season, with no more than 3 applications after petal fall. Do not apply within 3 days of harvest. Do not Plum Curculio Tarnished Plant Bug graze livestock in treated orchards. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.

140f35

13

팚

	DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS AI/A OZ/A)	REMARKS	
Filberts	Filbertworm Oblique Banded Leafroller	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply when insects appear. Do not apply more than 1.6 lbs ai. per acre per season. Do not apply within 14 days of harvest. Do not graze livestock in treated areas to livestock.	
Papaya (Florida only)	Fruit Fly	0.1 (6.4)	Apply by ground equipment only. Apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Apply when insects appear and repeat at 7-10 day intervals as needed to provide control. Do not make more than 6 applications per season. Do not apply within 7 days of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.	
Peaches	Peach Twig Borer Oriental Fruit Moth Green Fruitworm Lesser Peachtree Borer Plum Curculio Tarnished Plantbug Rose Chafer	0.1-0.3 (6.4-19.2)	Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Spray to wet all foliage. Can be applied up to 14 days before harvest. Do not apply more than 5 applications or 1.5 lbs. a.i. per acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.	
Pears (Dormant)	Pear Psylla	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply at a rate of 0.8 to 1.6 pints (0.2 to 0.4 lbs. active) per acre. May be combined with 2 to 8 gallons of spray oil per acre. Apply during the dormant through delayed dormant growth periods only. Do not apply more than 0.8 lbs. a.i. per acre per season. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.	

.

•••••

5aps

؛ د در د د د د

۲۲۲۲ ۲۲۲۲

د ددد د د

> > > > > > > > > > >

.

DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS AI/A OZ/A)	REMARKS
Pears (Pre-bloom)	Pear Psylla Coolling Moth Green Fruitworm	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply two pre-bloom sprays. Pre-bloom sprays can be applied from dormant through bud burst stages. Do not apply more than 0.8 lbs. a.i. per acre per sesson. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock.
Pistachios	Peach Twig Borer Naval Orangeworm Chinch Bug	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain full coverage of foliage. Apply at a rate of 0.8 to 1.6 pints (0.2-0.4 lb active) per acre. Do not apply more than 0.8 lbs a.i. per acre per season. Do not apply after 10 percent hull split. Nuts may be harvested on the day of application. Do not feed cover crops from treated areas to livestock.
Wainuts	Navel Orangeworm Codling Moth Walnut Husk Fly	0.2-0.4 (12.8-25.6)	Apply by air or ground equipment. By air, apply in a minimum of 10 gallons of water per acre. By ground, apply in a minimum of 25 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply when insects appear. Do not apply more than 1.6 lbs. a.i. per acre per season. Do not apply within 1 day of harvest. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to fivestock.

.

,

د د د د د د د د

 * * * * * *

 * * * *

 * * * *

 * * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

 * * *

. • • •

,

•••

1601-25

15

. .

- Alexandre - A

	DIRECTIONS FOR APPLICATION			
CROP	TARGET PESTS	RATE LBS AI/A OZ/A)	REMARKS	
a second of the		CTHER CROPS		
Alfalfa, Alfalfa grown for seed	Affalfa Caterpillar Armyworms Cutworms Loopers Webworms Potato Leafhopper Blue Alfalfa Aphid* Green Peach Aphid Pea Aphid* Green Cloverworm Spotted Alfalfa Aphid Velvetbean Caterpillar Alfalfa Weevil* Egyptian Alfalfa Weevil* Cucumber Beetle Meadow Spittlebug Plant Bugs including Lygus Spp.*	0.05-0.2 {3.2-12.8} 0.1-0.2 {6.4-12.8}	Apply by air or ground equipment. Use higher recommended dosage for increased pest pressure or for increased residual pest control. Apply in a minimum of 2 gallons of finished spray per acre by aircraft or 10 gallons of finished spray per acre with ground equipment. Use higher gallonage, 5-10 gallons by air, or 20 by ground, finished spray per acre when foliage is dense and/or when pest populations are high. Do not apply more than 0.2 lbs, ai. per cutting. When rates 0.1 lb. ai. per acre or less are used, application may be made on day of harvest. When rates greater than 0.1 lb. ai. per acre are used, do not apply within 14 days of harvest. Avoid application when bees are actively foraging by applying during the early morning or during the evening hours.	
Anichokes	Artichoke plume moth	0.1-0.3 (6.4-19.2)	Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 1.5 lbs. a.i. per acre per season. Buds may be harvested on the day of application.	
Asparagus	Cutworm Complex Asparagus Beetle	0.05-0.1 (3.2-6.4)	Apply by ground equipment only. Apply in a minimum of 10 gallons of water per acre. Use sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 0.4 lbs. a.i, per acre per season. Do not apply within 1 day of harvest.	

•••

, ..

170425

: .

`.

	DIRECTIONS FOR APPLICATION				
CROP	TARGET PESTS	RATE LBS AI/A OZ/A)	REMARKS		
Mushrooms (Mushroom houses and adjacent premise areas)	Mushroom Files (Sciarid and Phond Adults)	Apply as a fogging or aerosol treatment at the rate of 3.2 to 4.0 ounces (0.05-0.0625 tb ai) per 30 ounces of water or suitable diluent. Use 1 quart of solution per standard double house (35,000 cu ft.: 8,000 sq. ft.)	Preparation of the building prior to fogging: (1) Close all doors, windows, and ventilators, (2) Lock or barricade all entrances, turn off pilot lights, post warning signs, and take precautions to prevent persons and animals from entering the area. Use prior to filling the house, during cool-down, during spawning, up to pinning and between breaks. Do not use when mushrooms are present. Treat once daily or as needed when files appear. Do not make more than 20 applications prior to pinning of first break; apply no more than two applications between each break. Do not apply more than 30 applications total per crop of 5 breaks. Do not apply within 3 days of harvest. Length of exposure time should be limited to 1 hour; then ventilate the house. Use fans to ventilate in houses that do not have forced air circulation. Wear a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-146).		
Range Grass	Range Caterpillar	0.01. (0.64)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground, apply in a minimum of 10 gallons of water per acre. Use sufficient volume to obtain uniform coverage. Apply as needed. Do not apply more than once per year. Cattle may be present during application. Do not harvest or feed hay to livestock.		

٠,

۰.

180425

د د د د د د د

ود د ددد د ددد د د د د د د د د

 >
 >
 >
 >

 >
 >
 >
 >

 >
 >
 >
 >

 >
 >
 >
 >

 >
 >
 >
 >

 >
 >
 >
 >

 >
 >
 >
 >

 >
 >
 >
 >

DIRECTIONS FOR APPLICATION				
CROP	TARGET PESTS	RATE LBS AVA OZ/A)	REMARKS	
أحقارها الأخار المرادين	List and the state of the second	OTHER CROPS		
Soybeans	Green Cloverworm Mexican Bean Beetle Bean Leaf Beetle Cabbage Looper Potato Leafhopper Velvetbean Caterpillar Salimarsh Caterpillar	0.05-0.1 (3.2-6.4)	Apply by air or ground equipment. By air, apply in a minimum of 2 gallons of water per acre. By ground apply in a minimum of 10 gallons of water per acre. Do not apply more than 0.4 lbs. a.i. per acre per season. Can be applied up to 60 days before harvest. Do not graze treated areas or harvest for forage or hay.	
	Com Earworm Beet Armyworm Japanese Beetle*** Soybean Looper ⁴	0.1-0.2 (6.4-12.8)		
		··· ORNAMENTAL	S State Stat	
Asparagus Ferns	Japanese Beetle Asparagus Beetle Lygus Bugs	0.1 (6.4)	Apply to the fern stage of the asparagus plant after spear harvest when larval and adult stage of the asparagus beetle, lygus bugs and the adult stage of the Japanese beetle are present.	
Christmas Trees ^b (container and field grown)	Nantucket Pine Tip Moth	0.1-0.2 (6.4-12.8)	Apply first application when adults appear and repeat at 5-7 day intervals or as needed throughout the season. Use sufficient water to obtain full coverage of foliage.	
Chrysanthemums (Florida only)	Vegetable Leafminer	0.1-0.2 (6.4-12.8)	Apply by ground.	
Conifers (container and field grown)	Nantucket Pine Tip Moth	0.1-0.2 {6.4-12.8}	Apply first application when the adults appear and repeat at 5-7 day intervals or as needed throughout the season. AMBUSH may be diluted in a nonvolatile vegetable oil or in water in a minimum of one gallon of finished spray per acre using equipment calibrated to give adequate coverage.	

•

۰.

...

190425

.

د در در د

د د د د د د د د د د د

DIRECTIONS FOR APPLICATION					
CROP	TARGET PESTS	RATE LBS AI/A OZ/A)	REMARKS		
ORNAMENTALS					
Ornamentals Nursery Stock ^b (field-grown)	Beet Armyworm Cabbage Looper Citrus Thrip Heliothis Spp. Leafhoppers Leafminers Whiteflies Cutworms	0.1-0.2 (lbs ai/100 gal)	Marginal leaf burn may occur on Salvia, Dieffenbachia and Pteris Fern. Application to blooming plants may cause browning of petals. Not all species and varieties have been tested. Before treating large numbers of plants of a particular variety, treat a few plants and observe prior to full scale application. Apply to nonbearing plants of fruiting species.		
Pine Seed Orchard	Coneworms Seed bugs	13 fl oz/ 100 gals water (0.025% dilution by wi) for high volume sprayers; 68 fl oz/ 100 gals water (0.125% dilution by wi) for low volume sprayers; 48 fl oz/A application in not less than 10 gals of water for aerial applications	Apply first application within 1 week of female flower closure or peak pollen flight for webbing coneworm control. For other coneworms and seed bugs, apply first application within 30 days following female flower closure. Repeat applications at intervals of 4 weeks, but do not apply more than 6 applications. Apply approximately 5-10 gals of the 0.025% dilution per tree with high volume sprayers. With low volume sprayers, apply 100 gallons of the 0.125% dilution per acre. Do not graze livestock in treated areas. Do not feed cover crops from treated areas to livestock. Avoid contact with open water.		
Roses ⁶ (field-grown)	Heliothis Spp.	0.1-0.2 {6.4-12.8}	Apply by air or ground equipment. Apply in a minimum of 2 gallons/acre by air and 10 gallons of finished spray per acre with ground equipment.		
Roses ⁶ (greenhouse)	Beet Armyworm Cabbage Looper Omnivorous Leafroller	8 fl oz/100 gals water	AMBUSH is not phytotoxic to the following varieties of greenhouse roses: Ballena, Bettina, Cara Mia, Coquette, Excitement, Forever Yours, G. Wave, Jack Frost, Jr. Bridesmaid, Matador, Paul's Pink, Samantha, Seventeen, Town Crier, Tropicana and Visa. Other varieties may vary in their sensitivity to AMBUSH, and a small number of plants should be meated under local conditions to determine plant safety prior to commercial use.		

ŧ

< ÷_

÷

.

. .

_

•

۰.

19

•

200f25

LEAFY VEGETABLES (Except Brassica): Amaranth (leafy amaranth, Chinese spinach, tampala); arrugla (Roquette); Cardoon; celery; Chinese celery; celtuce; chervit; com salad; chrysanthemum, edible-leaved; chrysanthemum, garland; cress, garden; cress upland (yellow rocket, winter cress); dandetion; dock (sorrel); endive (escarole); fennel, Florence; lettuce (head and leaf); orach; parsley; purslane, garden; purslane, winter; radiochio (red chicory); rhubarb; spinach; spinach, vine (Malabar spinach, Indian spinach); spinach, new Zealand; Swiss chard.

2/07-35

- + CUCURBIT VEGETABLES: Chayote (fruit); Chinese Waxgourd (Chinese preserving meton); citron meton; cucumber; gherkin; gourds, edible (includes hyotan, cucuzza, hechima, Chinese okra); Momordica spp. (includes balsam apple, balsam pear, bitter meton, Chinese cucumber); muskmetons, including hybrids such as true cantaloupe, cantaloupe, casaba, crenshaw, golden pershaw meton, honeydew metons, honey balls, mango meton, Persian meton, pineapple meton, Santa Claus meton, snake meton; pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini, and winter (butternut squash, calabaza, hubbard squash, acom squash, spaghetti squash; watermeton, includion bybrids) including hybrids).
- AMBUSH may be applied in tank-mixtures with the following herbicides in accordance with label limitations and precautions. Label dosage rates should not be exceeded. GRAMOXONE⁵ SUPER; GRAMOXONE⁶ EXTRA, BICEP⁵, LASSO⁵/Atrazine Premix; LASSO⁵ 4E & ME; DUAL⁶ 8E; Atrazine 4L & 80W; BLADEX⁶ 80W & 4L; PRINCEP⁵; PRINCEP⁵ CALIBER 90⁵; 2.4-D; ROUNDUP⁵; BRONCO⁵; BANVEL⁵; EXTRAZINE⁶

*** Not for control of this insect in California.

Aids in suppression.

Currently not registered for use in California.
0.1% ai solution = 3½ pt formulation per 100 gallons of water.
Maximum use rate in California is 0.1 lbs. a.i./Acre.

Under heavy insect pressure use 0.2 lbs a.i./Acre. A second application of alternate chemistry may be necessary to reduce populations below economic thresholds.



...

RATE CONVERSION CHART ACRES TREATED/ POUNDS FLUID ACTIVE/ACRE OUNCES/ACRE PINTS/ACRE GALLON 0.05 3.2 0.2 40.0 6.4 0.4 20.0 0.10 0.125 8.0 0.5 16.0 0.15 9.6 0.6 13.3 0.175 11.2 8.7 11.4 0.20 12.8 0.8 10.0

,) 220125

 $\label{eq:ambulk} \begin{array}{l} \mathsf{AMBUSH}^{\mathtt{s}}, \mathsf{GRAM0XONE}^{\mathtt{s}}, \mathsf{and} \; \mathsf{KARATE}^{\mathtt{s}} \; \mathsf{are} \; \mathsf{trademarks} \; \mathsf{of} \; \mathtt{s} \; \mathsf{ZENECA} \; \mathsf{Group} \; \mathsf{Company}, \\ \mathsf{Ammo}^{\mathtt{s}}, \; \mathsf{Capture}^{\mathtt{s}}, \; \mathsf{Fury}^{\mathtt{m}}, \; \mathsf{Mustang}^{\mathtt{s}} \; \mathsf{and} \; \mathsf{Pounce}^{\mathtt{s}} \; \mathsf{are} \; \mathsf{trademarks} \; \mathsf{of} \; \mathsf{FMC} \; \mathsf{Corporation}, \\ \end{array}$ Annual Capitale , Fuly , Nuslang and Founce are trademarks of PMC Corporator. Baythroid is a trademark of Mobay Corporation. Bronco⁶, Lasso⁶, and Roundup⁶ are trademarks of Monsanto Company. Bicep⁶, Caliber⁴, Dual⁶, and Princep⁶ are trademarks of Ciba-Geigy Group. Asana⁸, Bladex⁸, Extrazine⁸ and Lannate⁸ are trademarks of E.I. duPont de Nemours & Co. Banvel⁶ is a trademark of Sandoz Crop Protection Corp.

Danitoi⁸ is a trademark of Sumitomo Chemical Company, LTD.

Scout X-TRA™ is a trademark of Hoechst-Roussel Agri-Vet Company. ©1995 ZENECA, Inc.

NOTES

.

--

24/4/25

 r_{k}

NOTES

....

.

••

د ه

د و د د د د د د د د د د د د

Fals

ZENECA Inc. Wilmington, Delaware 19850-5458

.

.

· 1

.

~

0400XXXX 069608 Rev. 9/97

>))))

، ، ، ، ، ، ، ب ، ب

د ر د ر ۰.

к кл Г 5 7 Элэк

}

, **,** ,

.

د د د د د د د د د د د د

. .

_ ._