

Talon™

RODENTICIDE BAIT PACK (MINI-PELLETS)

For Effective Indoor Control of Commensal Mice and Rats

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY BE HARMFUL IF EATEN. IT IS A LETHAL TOXIC POWDER TO HUMANS AND DOMESTIC ANIMALS. AND PETS. WASH HANDS AFTER HANDLING.

KEEP OUT OF REACH OF CHILDREN. KEEP THIS PRODUCT FROM CONTACT WITH ALL OTHER HUMANS AND DOMESTIC ANIMALS.

IF THIS PRODUCT IS EATEN IMMEDIATELY INDUCE VOMITING AND TAKE IMMEDIATE MEDICAL ATTENTION.

NOTE TO PHYSICIAN

This product may reduce the clotting ability of the blood and a severe hemorrhagic syndrome, intraventricular and intracranial, may be combined with blood transfusions are indicated. Also, the product may be aggravated by warfarin, heparin, benzhydrocycoumarin, Vitamin K₁ and other blood clotting agents.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not use in or near water.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed. Storage in original container is prohibited. Do not reuse empty container.

DISPOSAL: Bait that cannot be used should be disposed of in a landfill or other suitable location or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers. For more information, consult Federal, State or local disposal authorities. Do not reuse after disposal unless such is indicated on label.

Kills warfarin-resistant Norway Rats
Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins

Active Ingredient

1,3-Dichloro-5-(4-chlorophenyl)-4-isopropyl-2-imidazolepropanolamine Hydrochloride (90% active ingredient)

Inert Ingredients

Total

0.005%

99.995%

100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

Mix with animal feed or water
Best Adaptation: Prevention of infestation



ICI Americas Inc.

Wilmington Delaware 19897

EPA Reg. No. 101-111-01

FPA Reg. No. 101-111-01

Net Content: 100 Pellets

Net Weight: 100 g

DIRECTIONS FOR USE

Apply to all areas where rodents are known to be present.

RECOMMENDATIONS

Apply to all areas where rodents are known to be present. Apply to all areas where rodents are known to be present. Apply to all areas where rodents are known to be present.

SELECTION OF TREATMENT AREA

Select areas where rodents are known to be present. Select areas where rodents are known to be present. Select areas where rodents are known to be present.

APPLICATION DIRECTIONS

Norway and Roof Rats

Apply 4 to 5 pellets per 100 sq ft of area. Apply 4 to 5 pellets per 100 sq ft of area. Apply 4 to 5 pellets per 100 sq ft of area.

House Mice

Apply 1 to 2 pellets per 100 sq ft of area. Apply 1 to 2 pellets per 100 sq ft of area. Apply 1 to 2 pellets per 100 sq ft of area.

Rats & Mice

Apply 1 to 2 pellets per 100 sq ft of area. Apply 1 to 2 pellets per 100 sq ft of area. Apply 1 to 2 pellets per 100 sq ft of area.

BEST DOCUMENT AVAILABLE

ACCEPTED