

Talon-G

RODENTICIDE MINI-PELLETS IN MOUSE BOX
For Effective Control of House Mice

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS AND PETS. WASH HANDS AFTER HANDLING BAIT. IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE.

NOTE TO PHYSICIAN OR VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. If this occurs, intravenous and oral administration of vitamin K₁ combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg) and should be given and repeated as necessary (based on monitoring of prothrombin times).

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

STORAGE AND DISPOSAL

Do not reuse container. Wrap container and put in trash collection.

Kills warfarin-resistant house mice.

House mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

Active ingredient: 3-[3-(4'-bromo[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one ... 99.995%
Inert ingredients ... 0.005%
Total ... 100.000%

**KEEP OUT OF
REACH OF
CHILDREN
CAUTION**

May Be Harmful or Fatal if Swallowed. Read Additional Precautionary Statements on Side Panel.



ICI Americas Inc.

Wilmington, Delaware 19897

EPA Reg No 10182-25

EPA Est No 523-1L-1

Net Contents 4 ounces

Batch Number

ACCEPTED

10182-25

MAR 8 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 10182-25

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of House Mice in homes, industrial and commercial buildings. For indoor use only. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Place Mouse Boxes wherever you see signs of mice. Place bait along walls, in corners and concealed places.

APPLICATION DIRECTIONS: Place Mouse Boxes at points of mouse activity. Replace containers as they are consumed. Two or more containers should be used where there is extremely high mouse activity. Boxes not touched after 3 days should be moved to a new location where mouse activity is evident. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.