

17 JAN 1986

164,984

300
17

Barbara Kaminski
ICI Americas, Inc.
Agricultural Chemicals Division
Wilmington, DE 19897

Dear Ms. Kaminski:

Subject: Talon-G Rodenticide
EPA Registration No. 10182-39
Your Application Dated October 28, 1985


The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(A) is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

Delete "General Classification."
3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,


William H. Miller
Product Manager (16)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure
86774:Palmateer/Miller;CBI;85/02;KENCO;1/8/86;Del.1/17/86;eg:VO

CONCURRENCES

SYMBOL ▶								
SURNAME ▶								
DATE ▶								

Talon®-G

RODENTICIDE BAIT PACK (PELLETS)

For Effective Control of Commensal Rats

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT.

IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI Americas Inc. 302/575-3080.

IF BAIT IS EATEN BY ANIMALS OR PETS, call University of Illinois Animal Poison Control Center 217/333-3611 or ICI Americas Inc. 302/575-3000 for 24-hour emergency assistance.

In case of significant spill, call CHEMTREC 800/424-9300.

NOTE TO PHYSICIAN OR VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated as in poisoning from overdose of bishydroxy coumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

NOTICE TO BUYER AND USER: Seller warrants that the product conforms to the chemical description thereon and is conformant to the purposes stated on the label. This product may be used under the following conditions: (1) For the purposes stated on the label; (2) The use of this product shall be limited to the use of this product as a rodenticide for the control of commensal rats; (3) This product shall not be used for any other purpose; (4) This product shall not be used in any area where it may be injurious to humans, animals, birds, fish, or wildlife; (5) This product shall not be used in any area where it may be injurious to crops, plants, or other vegetation; (6) This product shall not be used in any area where it may be injurious to the environment; (7) This product shall not be used in any area where it may be injurious to the health of humans, animals, birds, fish, or wildlife; (8) This product shall not be used in any area where it may be injurious to the health of crops, plants, or other vegetation; (9) This product shall not be used in any area where it may be injurious to the health of the environment; (10) This product shall not be used in any area where it may be injurious to the health of the community. The user shall be responsible for the safe handling, use, and disposal of this product. The user shall be responsible for the safe handling, use, and disposal of this product. The user shall be responsible for the safe handling, use, and disposal of this product.

KRM warfarin-resistant Norway Rats.

Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

Active Ingredient:

Brodifacoum-
 3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-
 1,2,3,4-tetrahydro-1-naphthalenyl]-4-
 hydroxy-2H-1-benzopyran-2-one

Inert Ingredients

Total

0.005%
 99.995%
 100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

May Be Harmful or Fatal if Swallowed
 Read Additional Precautionary Statements
 on Side Panel



ICI Americas Inc.

Wilmington, Delaware 19897

EPA Reg. No. 10182-39

EPA Est. No. 2393 WI-1

Net Contents 4/100 x 50 gm (1.76 oz) bait pack boxes

Batch Number

Do not contaminate water container except for the Storage: Store in original and pets
 Pesticide Disposal: Waste of on site or at an approved
 Cashflow Disposal: Dispose if allowed by state and

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of federal labeling

USE RESTRICTIONS: For homes, industrial, commercial use in and around terminal buildings. Do not proof bait boxes or in local wildlife. Do not place bait on surfaces that come in


Authorized by the USE processing plants

SELECTION OF TREATMENT: find and consume the bait openings, in or beside burrow walls, or in locations where much food as possible

APPLICATION DIRECTIONS

Norway and Roof Rats: Apply per placement. Maintain signs of rat activity cease at intervals of 8 to 12 feet needed at points of very fresh bait for 15 days or contaminated or spoiled and unconsumed bait or eliminate food, water, and occur, repeat treatment, fish permanent bait station

31/51 259, 1794 11




PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS—CAUTION MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS, AND PETS. WASH HANDS AFTER HANDLING BAIT. IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call ICI American Inc. 302.575-3000. IF BAIT IS EATEN BY ANIMALS OR PETS, CALL University of Illinois Animal Poison Control Center 217.333-3611 or ICI American Inc. 302.575-3000 for 24-hour emergency assistance. In case of significant spill, call CHEMTREC 800-424-9300.

NOTE TO PHYSICIAN OR VETERINARIAN: This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and oral administration of vitamin K₁ are indicated, as in poisoning from overdose of bishydroxycoumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

STORAGE AND DISPOSAL: Store away from children and pets. Wrap empty or used container and put in trash.




Talon-G
RODENTICIDE BAIT PACK (PELLETS)
For Effective Control of Commercial Rats

Kills warlike-resistant Norway Rats. Rodents may consume lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT:
Brodifacoum 0.005%
3-[3-(4-bromo-1,1-biphenyl-4-yl)-1,2,4-tetrazolo-1-imidazolyl]-4-hydroxy-2H-1-benzopyran-2-one 99.995%
INERT INGREDIENTS 100.000%
Total

KEEP OUT OF REACH OF CHILDREN
CAUTION

May Be Harmful or Fatal if Swallowed
Read Additional Precautionary Statements on Side Panel



ICI American Inc.
Agricultural Chemical Division
Wilmington, Delaware 19802

EPA Reg. No. 10182-39
EPA Est. No. 2393 WI-1
Net Contents 50 gms (1.76 oz)
Batch Number
ACD 790-236R2 1/E

DIRECTIONS FOR USE General Classification: It is a violation of federal law to use this product in a manner inconsistent with its labeling. **Use Restrictions:** For control of Norway Rats and Roof rats in and around homes, industrial, commercial, agricultural and public buildings. TALON may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. This product must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Authorized by the USDA for use in federally inspected meat, poultry and egg processing plants. **Selection of Treatment Areas:** Determine areas where rats will most likely feed and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS Norway and Roof Rats: Apply 2 to 8 bait packs (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease. **Mice:** Open pack and apply 1/4 to 1/2 ounce of bait at intervals of 8 to 12 feet per placement. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease. **Rats and Mice:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph. To prevent reinfestation, eliminate food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

10182-39

Talon-G

RODENTICIDE BAIT PACK PELLETS

For Effective Control of Commercial Rats

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. KEEP AWAY FROM HUMANS, DOMESTIC ANIMALS AND PETS. WASH HANDS AFTER HANDLING BAIT.

IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE. For 24-hour emergency assistance call 302/575-3000.

IF BAIT IS EATEN BY ANIMALS OR PETS, CALL 217/333-3611 or 302/575-3000 for 24-hour emergency assistance.

In case of significant spill, call CHEMTREC 800/474-9300.

NOTE TO PHYSICIAN OR VETERINARIAN

This product may reduce the clotting ability of the blood and cause hemorrhaging. If poisoning occurs, intramuscular and/or oral administration of vitamin K₁ are indicated as is poisoning from overdose of bishydroxycoumarin. For human cases, vitamin K₁ is antidotal at doses of 10-20 mg (not mg/kg). For animal cases, vitamin K₁ is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Keep out of any body of water.

NOTICE TO BUYER AND USER: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller and Buyer and User assumes the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

ACD 790 218R1 14

High water-resistant Barony Bait. Rodents may consume a baited area in one feeding with first feed pellets appearing lost or few days after treatment begins.

Active Ingredient

Brodifacoum

3-[2-(4-bromo-1,1'-biphenyl)-4-yl]-

1,2,3,4-tetrahydro-1-naphthalenyl]-4-

hydroxy-2H-1-benzopyran-2-one

Inert Ingredients

0.000%

99.999%

100.000%

Total

KEEP OUT OF REACH OF CHILDREN CAUTION

May Be Harmful or Fatal if Swallowed
Read Additional Precautionary Statements on Side Panel



ICI Americas Inc.

Wilmington Delaware 19877

EPA Reg. No. 10182-39

EPA Est. No. 2393-WI-1

Net Contents: 100.50gm (1.76oz) bait packs

Batch Number

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse or recycle rodenticide.

Storage: Store in original container only in a dry place inaccessible to children.

Portable Storage: Wastes resulting from the use of this product may be disposed of in a sanitary landfill or by incineration.

Container Disposal: Dispose of bait container in a sanitary landfill or by incineration.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with the USE RESTRICTIONS. For control of Norway Rats and Roof rats in and around buildings, TALON may also be used in and around transport vehicles (ships, trains, use in sewers. This product must be placed in tamper-proof bait boxes or in locations inaccessible to wildlife. Do not place bait in areas where there is a possibility of contamination.

Authorized by the USDA for use in federally inspected meat, poultry and other animal products. **SELECTION OF TREATMENT AREAS:** Determine areas where rats will most likely enter, by gnawed openings, in or beside burrows, in corners and concealed rodents or their signs have been observed. Remove as much bait as possible.

APPLICATION DIRECTIONS

Barony and Roof Rats: Apply 2 to 8 bait packs (usually at intervals of 15 to 30 feet) per bait for 10 days or until signs of rat activity cease. Remove bait packs and replace with fresh bait for 15 days or until signs of mouse activity cease. Rats and Roof Rats and dispose of all dead animals and unconsumed bait according to "Disposal" paragraph and haulage as much as possible. If reinfestation does occur, repeat treatment and establish permanent bait stations and replenish bait as needed.

ACCEPTED with COMMENT in EPA L. C. R. 14

JAN 2 1988

Under the Fungicide, and as amended, for registered under EPA

10182-3